

$SO(1,4)$ flux-balance laws of de Sitter spacetime at quadrupolar orderGeoffrey Compère^{1,*}, Sk Jahanur Hoque^{1,2,3,†} and Emine Seyma Kutluk^{4,5,‡}¹*Université Libre de Bruxelles, International Solvay Institutes, CP 231, B-1050 Brussels, Belgium*²*Birla Institute of Technology and Science, Pilani, Hyderabad Campus, Jawaharnagar, Hyderabad 500 078, India*³*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 180 00 Prague 8, Czech Republic*⁴*Scuola Normale Superiore, Piazza dei Cavalieri 7, Pisa 56126, Italy*⁵*INFN Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy*

(Received 6 December 2024; accepted 21 February 2025; published 11 March 2025)

The linear solution for quadrupolar perturbations around de Sitter spacetime was recently constructed. In this paper, we provide the flux-balance laws for each background symmetry (dilatations, rotations, spatial translations and cosmological boosts) in terms of source moments at quadrupolar order. We write the dilatation flux-balance law in two distinct ways, which allows to contrast two distinct proposals for the negative definite energy flux. The standard Poincaré flux-balance laws at future null infinity are recovered in the flat limit of the $SO(1,4)$ flux-balance laws.

DOI: [10.1103/PhysRevD.111.064039](https://doi.org/10.1103/PhysRevD.111.064039)**I. INTRODUCTION**

According to the standard model of cosmology, our Universe admits a positive cosmological constant $\Lambda \approx 3 \times 10^{-122}$ in Planck units [1,2]. Though tiny, the effect of the cosmological constant dominates in the infrared and leads to an asymptotically de Sitter spacetime structure at late times, which overarch in particular the definition of asymptotically conserved quantities. Besides, gravitational wave observations have provided within the last decade new insights into astrophysics and cosmology [3]. These observations have led to a large theoretical effort to define gravitational waves emitted from localized sources in a cosmological background, see e.g. [4–60].

Many simple theoretical questions remain unanswered. In this article, we will concentrate our focus on two questions: What is the role of the $SO(1,4)$ de Sitter symmetry regarding gravitational radiation? What is the definition of energy in asymptotically de Sitter spacetime?

It was demonstrated in [36,39] that even though gravitational waves change the leading components of the metric at future infinity, a boundary gauge exists such that the asymptotic symmetry algebra of asymptotically de Sitter spacetime is the infinite-dimensional Λ -BMS algebra, which reduces to the BMS algebra [61,62] in the flat limit. The structure constants of the Λ -BMS algebra generically depend upon the radiation field. However, at quadratic order in perturbation theory around de Sitter, one can isolate a

universal $SO(1,4)$ algebra of background symmetries, as we will review in Sec. II, simply because the fluxes are already quadratic in the fields once the symmetry generators are taken to be the background Killing vectors of de Sitter. Our first objective is to derive the set of flux-balance laws associated with the background $SO(1,4)$ symmetry and evaluate them in the physical phase space. We will perform this computation at the level of the quadrupolar truncation of the linear spectrum [57], which has been demonstrated to match [58] with the spectrum of perturbations obtained from independent derivations [55,58].

Several formulas for the energy flux in asymptotically de Sitter spacetime have been proposed [9,12,13,21–23,29–31,34,38,41,50,54,60,63–66]. Our second objective is to identify from the dilatation flux-balance law how one can identify a charge and a flux such that (i) the flux reduces to the standard energy flux in the flat limit, (ii) the flux is manifestly non-negative at quadratic order. We will contrast our expressions for the energy loss and angular momentum flux with part of the literature. The linear momentum and boost charge loss formulas that we will derive have not yet appeared in that form in the literature to the best of our knowledge.

The rest of the paper is organized as follows. We first define the $SO(1,4)$ flux-balance laws in perturbation theory around de Sitter spacetime from first principles in Sec. II. We then evaluate them on the solution space of even and odd quadratic perturbations of de Sitter in Sec. III and we prove that the standard flat spacetime limit is obtained. We compare our expressions of the energy loss and angular momentum loss with the literature in Sec. IV. We finally conclude in Sec. V.

*Contact author: geoffrey.compere@ulb.be†Contact author: jahanur.hoque@hyderabad.bits-pilani.ac.in‡Contact author: emine.kutluk@sns.it

II. CHARGES AND FLUXES ON DE SITTER SPACETIME

The future region \mathcal{I}^+ of asymptotically de Sitter spacetimes with cosmological constant $\Lambda = 3H^2$ admits a Starobinsky expansion [67] of the form

$$ds^2 = -\frac{d\tau^2}{H^2\tau^2} + \frac{1}{\tau^2}(g_{ab}^{(0)} + \tau^2 g_{ab}^{(2)} + \tau^3 g_{ab}^{(3)} + O(\tau^4))dx^a dx^b, \quad (2.1)$$

where $\tau < 0$ denotes time and x^a are the other coordinates. In terms of the conformal completion, $\tau = 0$ is the future boundary \mathcal{I}^+ of asymptotically de Sitter spacetimes and x^a are the coordinates on \mathcal{I}^+ . Latin letters a, b, \dots will always refer to three-dimensional indices of fields defined on \mathcal{I}^+ . The Starobinsky expansion is related to the Fefferman-Graham expansion of asymptotically anti-de Sitter spacetimes by an analytical continuation. We call $g_{ab}^{(0)}$ the three-dimensional metric on the future boundary of asymptotically de Sitter \mathcal{I}^+ , $g_{(0)}^{ab}$ its inverse and we will denote as $D_a^{(0)}$ its metric-compatible covariant derivative. The holographic stress-energy tensor defined as $T_{ab} = \frac{3H}{16\pi G} g_{ab}^{(3)}$ is trace-free and divergence-free with respect to the boundary metric $g_{ab}^{(0)}$ outside of sources:

$$D_a^{(0)} T^{ab} = 0, \quad g_{(0)}^{ab} T_{ab} = 0, \quad (2.2)$$

where T^{ab} is defined as T_{ab} with indices raised with the inverse metric $g_{(0)}^{ab}$.

In the presence of gravitational radiation, the boundary metric $g_{ab}^{(0)}$ generically admits no symmetry. In perturbation theory where $g_{\mu\nu} = \bar{g}_{\mu\nu} + \delta g_{\mu\nu}$ (where μ, ν denote 4-dimensional indices), we can however write

$$g_{ab}^{(0)} = \bar{g}_{ab}^{(0)} + \delta g_{ab}^{(0)}, \quad (2.3)$$

where the background boundary metric $\bar{g}_{ab}^{(0)}$, which describes the future boundary of de Sitter $\mathcal{I}_{\text{dS}}^+$, admits symmetries. Motivated by the Bondi framework and by the description of spatially compact binary systems that reach future infinity \mathcal{I}^+ at a single point $u = +\infty$, we choose $\bar{g}_{ab}^{(0)}$ to be the metric on $\mathbb{R} \times S^2$

$$\bar{g}_{ab}^{(0)} dx^a dx^b = H^2 du^2 + \hat{q}_{AB}(x^C) dx^A dx^B. \quad (2.4)$$

Here \hat{q}_{AB} is the unit metric on S^2 and u ranges over the real line. These coordinates do not cover the entire $\mathcal{I}_{\text{dS}}^+$ of de Sitter, which has topology S^3 , but $\mathcal{I}_{\text{dS}}^+$ minus two points. The topology of \mathcal{I}^+ depends upon the number of massive bodies. Here, we consider at least two such points removed

at $u = \pm\infty$. We also gauge fix the perturbation to so-called Λ -BMS gauge [36] such that

$$g_{ab}^{(0)} dx^a dx^b = H^2 du^2 + q_{AB}(u, x^C) dx^A dx^B, \quad (2.5)$$

and $\sqrt{q} = \sqrt{\hat{q}}$. The inverse of q_{AB} is denoted q^{AB} . The shear is defined as $C_{AB} = H^{-2} \partial_u q_{AB}$.

The background boundary metric admits 10 conformal Killing vectors $\bar{\xi}^a(u, x^A)$ which are in one-to-one correspondence with the Killing symmetries of the de Sitter background and define the $SO(1, 4)$ group. By definition they obey

$$\bar{g}_{(0)c(a} \bar{D}_{(0)b)} \bar{\xi}^c = 0, \quad (2.6)$$

where $\bar{D}_{(0)a}$ is the covariant derivative compatible with the background boundary metric $\bar{g}_{(0)ab}$ and $\langle \cdot \rangle$ denote the symmetric trace-free part. The 10 conformal Killing vectors consist of the dilatations and rotations (which are Killing vectors of the background boundary metric) and the spatial translations and cosmological boosts (which are proper conformal Killing vectors of the background boundary metric). They are given by the following table:

Charge	Name	Generator
E	Dilatation	$\bar{\xi}^u = 1, \bar{\xi}^A = 0$
P^i	Spatial translation	$\bar{\xi}_{(i)}^u = n_i \exp(Hu),$ $\bar{\xi}_{(i)}^A = -H \hat{D}^A n_i \exp(Hu)$
L^i	Rotation	$\bar{\xi}_{(i)}^u = 0, \bar{\xi}_{(i)}^A = -\epsilon^{AB} \hat{D}_B n_i$
K^i	Cosmological boost	$\bar{\xi}_{(i)}^u = n_i \exp(-Hu),$ $\bar{\xi}_{(i)}^A = H \hat{D}^A n_i \exp(-Hu)$

Here n_i is the unit normal to the round sphere embedded in \mathbb{R}^3 , and \hat{D}_A is the covariant derivative compatible with the unit sphere metric.

For an arbitrary vector ξ^a , one has the identity

$$\begin{aligned} & \int_M dud^2\Omega \sqrt{\bar{q}} g_{(0)}^{ac} D_{(0)c}(T_{ab} \xi^b) \\ &= \int_M dud^2\Omega \sqrt{\bar{q}} T^{ab} g_{(0)c(a} D_{(0)b)} \xi^c \end{aligned} \quad (2.7)$$

for any region M of \mathcal{I}^+ outside of sources thanks to Eq. (2.2). In perturbation theory, T_{ab} admits linear and quadratic contributions in the perturbation $\delta g_{\mu\nu}$, $T_{ab} = \delta T_{ab} + \delta^2 T_{ab} + O((\delta g)^3)$. Let us now set $\xi^a = \bar{\xi}^a$ a background conformal Killing vector. Then, for any choice of spheres S_1 and S_2 such that $\partial M = S_1 \cup S_2$ (with opposite relative orientation), one has the equalities at linear and quadratic order, respectively,

$$\int_{S_1} d^2\Omega \sqrt{\bar{q}} n^a \delta T_{ab} \bar{\xi}^b - \int_{S_2} d^2\Omega \sqrt{\bar{q}} n^a \delta T_{ab} \bar{\xi}^b = 0, \quad (2.8)$$

$$\begin{aligned} & \int_{S_1} d^2\Omega \sqrt{\bar{q}} n^a \delta^2 T_{ab} \bar{\xi}^b - \int_{S_2} d^2\Omega \sqrt{\bar{q}} n^a \delta^2 T_{ab} \bar{\xi}^b \\ &= \int_M du d^2\Omega \sqrt{\bar{q}} \delta T^{ab} \mathcal{L}_{\bar{\xi}} \delta g_{ab}^{(0)}, \end{aligned} \quad (2.9)$$

where n^a is the unit normal to S_1 and S_2 defined as $n^a \partial_a = H^{-1} \partial_u$. The right-hand side of Eq. (2.8) is zero thanks to the conformal Killing equation (2.6).

Finally, we derived that on a given section of \mathcal{I}^+ the charge defined from the holographic stress-energy tensor as

$$Q_{\bar{\xi}}^T(u) \equiv - \int_{S^2} d^2\Omega \sqrt{\bar{q}} n^a T_{ab} \bar{\xi}^b \quad (2.10)$$

is a constant in linear theory, i.e. $Q_{\bar{\xi}}^T(u_1) = Q_{\bar{\xi}}^T(u_2)$ for any u_1, u_2 . At quadratic order, the presence of radiation will induce a change in $Q_{\bar{\xi}}^T(u)$ given by the flux-balance formula

$$Q_{\bar{\xi}}^T(u_2) - Q_{\bar{\xi}}^T(u_1) = \int_{u_1}^{u_2} du \dot{Q}_{\bar{\xi}}^T(u), \quad (2.11)$$

$$\dot{Q}_{\bar{\xi}}^T(u) \equiv \int_S d^2\Omega \sqrt{\bar{q}} \delta T_{ab} \bar{g}_{(0)}^{ac} \bar{g}_{(0)}^{bd} \mathcal{L}_{\bar{\xi}} \delta g_{cd}^{(0)} + O((\delta g)^3). \quad (2.12)$$

Instead of considering the background symmetry generator $\bar{\xi}^a$ we could consider an arbitrary perturbation of the vector $\xi^a = \bar{\xi}^a + \delta \xi^a$. Assuming that the Λ -BMS gauge is obeyed in the full theory, the generic infinitesimal diffeomorphism of the (4-dimensional) metric is parametrized by a vector ξ^a of the boundary metric which obeys the equations

$$\partial_u \xi^u = \frac{1}{2\sqrt{\bar{q}}} \partial_A (\sqrt{\bar{q}} \xi^A), \quad (2.13a)$$

$$\partial_u \xi^A = -H^2 q^{AB} \partial_B \xi^u. \quad (2.13b)$$

Let us denote as ξ^μ the 4-dimensional vector associated with ξ^a in Λ -BMS gauge. A linearized diffeomorphism of the 4-dimensional metric is simply the Lie derivative of the background metric with respect to ξ^μ , i.e. linearized gravitational fields are defined up to a gauge transformation $\delta g_{\mu\nu} \rightarrow \delta g_{\mu\nu} + \mathcal{L}_\xi \bar{g}_{\mu\nu}$. We have $\mathcal{L}_\xi \bar{g}_{\mu\nu} = \mathcal{L}_{\delta \xi} \bar{g}_{\mu\nu}$ where the equality follows from the Killing equation $\mathcal{L}_{\bar{\xi}} \bar{g}_{\mu\nu} = 0$. Therefore, at linear order in perturbation theory and in Λ -BMS gauge, a perturbed vector $\delta \xi^a$ obeys the linearization of Eq. (2.13) where q_{AB} is replaced by the background \bar{q}_{AB} :

$$\partial_u \delta \xi^u = \frac{1}{2\sqrt{\bar{q}}} \partial_A (\sqrt{\bar{q}} \delta \xi^A), \quad (2.14a)$$

$$\partial_u \delta \xi^A = -H^2 \bar{q}^{AB} \partial_B \delta \xi^u. \quad (2.14b)$$

For dilatations and rotations, we have $\partial_A \bar{\xi}^u = 0$ and we can fix $\xi^a = \bar{\xi}^a$ in the nonlinear theory because the right-hand side of Eq. (2.13b) is zero. We can therefore define universal (i.e. field independent) time translation and rotations that form a $\mathbb{R} \times SO(3)$ subalgebra of the Λ -BMS algebra in the nonlinear theory. This subalgebra is part of the larger $\mathbb{R} \times \text{ADiff}(S^2)$ universal subalgebra of the Λ -BMS algebra which consists of time translations and area-preserving diffeomorphism [with generators $\xi^u = 0$, $\xi^A = \epsilon^{AB} \partial_B \Psi(\theta, \phi)$]. The universal $\mathbb{R} \times SO(3)$ subalgebra of asymptotic symmetries was also identified in [55] in a different boundary gauge. The background spatial translations and cosmological boosts are generically deformed in the nonlinear theory. Yet, in the linear theory, the deformation vector is a background Λ -BMS generator [that obeys by definition (2.14)], which we can conventionally fix to be vanishing. With this convention, all background symmetries are undeformed in the linear theory.

We define the cosmological Bondi mass aspect $M^{(\Lambda)}$, angular momentum aspect $N_A^{(\Lambda)}$ and higher Bondi aspect $J_{AB}^{(\Lambda)}$ from the decomposition of the stress-energy tensor [see Eq. (3.15) of [36]]

$$T_{ab} = \frac{3H}{16\pi G} \begin{bmatrix} -\frac{4}{3} M^{(\Lambda)} & -\frac{2}{3} N_B^{(\Lambda)} \\ -\frac{2}{3} N_B^{(\Lambda)} & J_{AB}^{(\Lambda)} + \frac{2}{\Lambda} M^{(\Lambda)} q_{AB} \end{bmatrix}. \quad (2.15)$$

Here $J_{AB}^{(\Lambda)}$ is traceless: $q^{AB} J_{AB}^{(\Lambda)} = 0$. The Bondi mass aspect M and Bondi angular momentum aspect N_A defined in the flat limit $H \mapsto 0$ are related to the cosmological quantities as [see Eqs. (2.38) and (2.52) of [36]]

$$M^{(\Lambda)} = M + \frac{1}{16} \partial_u (C_{CD} C^{CD}), \quad (2.16a)$$

$$N_A^{(\Lambda)} = N_A - \frac{1}{2H^2} D^B N_{AB} + \frac{9}{32} \partial_A (C_{CD} C^{CD}). \quad (2.16b)$$

The charges (for the vector ξ^a) can be rewritten as

$$Q_\xi^T = \frac{1}{4\pi G} \int_{S^2} d^2\Omega \sqrt{\bar{q}} \left[M^{(\Lambda)} \xi^u + \frac{1}{2} N_A^{(\Lambda)} \xi^A \right]. \quad (2.17)$$

In terms of Bondi aspects, the conservation of the holographic stress-energy tensor (2.2) is equivalent to

$$\partial_u M^{(\Lambda)} + \frac{H^2}{2} D^A N_A^{(\Lambda)} + \frac{3H^4}{8} C_{AB} J^{(\Lambda)AB} = 0, \quad (2.18a)$$

$$\partial_u N_A^{(\Lambda)} - \partial_A M^{(\Lambda)} - \frac{3H^2}{2} D^B J_{AB}^{(\Lambda)} = 0. \quad (2.18b)$$

Here D_A is the covariant derivative compatible with q_{AB} . Using Eqs. (2.18) and (2.13), the charge flux is then equivalently written as

$$\dot{Q}_\xi^T = -\frac{3H^2}{32\pi G} \int d^2\Omega \sqrt{q} (H^2 \xi^u C_{AB} + 2D_A \xi_B) J^{(\Lambda)AB}, \quad (2.19)$$

$$= -\frac{3H^2}{32\pi G} \int d^2\Omega \sqrt{q} (\xi^u \partial_u q_{AB} + \mathcal{L}_{\dot{\xi}} q_{AB}) J^{(\Lambda)AB}, \quad (2.20)$$

where the Lie derivative is here understood to act on the S^2 manifold using the pullback vector $\dot{\xi} = \xi^A \partial_A$.

The flux is quadratic in the metric perturbation. Upon expanding $\xi^a = \bar{\xi}^a + \delta\xi^a + O(\delta^2)$ [as well as its 2-dimensional pullback $\dot{\xi}^A = \bar{\dot{\xi}}^A + \delta\dot{\xi}^A + O(\delta^2)$], $q_{AB} = \bar{q}_{AB} + \delta q_{AB} + O(\delta^2)$, $J_{AB}^{(\Lambda)} = \delta J_{AB}^{(\Lambda)} + \delta^2 J_{AB}^{(\Lambda)} + O(\delta^3)$ we have

$$\delta J_{AB}^{(\Lambda)} \bar{q}^{AB} = 0, \quad (2.21)$$

$$\delta^2 J_{AB}^{(\Lambda)} \bar{q}^{AB} = \delta J_{AB}^{(\Lambda)} \bar{q}^{AC} \bar{q}^{BD} \delta q_{CD}, \quad (2.22)$$

as a consequence of the tracelessness of $J^{(\Lambda)AB}$, as well as

$$\begin{aligned} & 2\delta(D^A \bar{\xi}^B) \delta J_{AB}^{(\Lambda)} + 2\delta^2 J^{(\Lambda)AB} \bar{D}_B \bar{\xi}_A \\ &= (-\mathcal{L}_{\dot{\xi}} \delta q^{AB} + \bar{D}_C \bar{\xi}^C \bar{q}^{AC} \bar{q}^{BD} \delta q_{CD}) \delta J_{AB}^{(\Lambda)} \\ &= (\mathcal{L}_{\dot{\xi}} \delta q_{AB} - \bar{D}_C \bar{\xi}^C \delta q_{AB}) \bar{q}^{AC} \bar{q}^{BD} \delta J_{CD}^{(\Lambda)}. \end{aligned} \quad (2.23)$$

The flux formula therefore reads at quadratic level as

$$\begin{aligned} \dot{Q}_\xi^T &= -\frac{3H^2}{32\pi G} \int d^2\Omega \sqrt{q} \bar{q}^{AC} \bar{q}^{BD} (\bar{\xi}^u \partial_u \delta q_{AB} + \mathcal{L}_{\dot{\xi}} \delta q_{AB} \\ &+ \mathcal{L}_{\delta\dot{\xi}} \bar{q}_{AB} - \bar{D}_E \bar{\xi}^E \delta q_{AB}) \delta J_{CD}^{(\Lambda)} + O(\delta^3). \end{aligned} \quad (2.24)$$

Here $\delta\xi^a$ obeys to Eq. (2.14). We choose the solution $\delta\xi^a = 0$, i.e. we do not act with any further Λ -BMS symmetries but only with the background symmetry. In that sense, there are flux-balance laws associated with $SO(1,4)$ symmetry at quadratic order in perturbation theory.

In the flat limit, using the conventions of [68], the generators associated with time translations, spatial translations and rotations are simply the limit $H \mapsto 0$ of E , P^i and L^i , respectively. Lorentz boosts are associated with the limit $H \mapsto 0$ of the generator $\frac{1}{2H}(K^i - P^i)$ with components $\bar{\xi}_{(i)}^u = -\frac{1}{H} n_i \sinh(Hu)$, $\bar{\xi}_{(i)}^A = \bar{D}^A n_i \cosh(Hu)$. Using these relationships we notice that the charges Q_ξ^T do not have the standard flat limit, see Eq. (3.2) of [69]. In order to recover the standard flat limit, we will define instead the charges as

$$Q_\xi(u) = Q_\xi^T(u) + \Delta Q_\xi(u), \quad (2.25)$$

where the shifting term $\Delta Q_\xi = O((\delta g)^2)$ will be defined in the next section. There is an ambiguity in defining this shift. We will partially fix this ambiguity by requiring that the energy is positive definite.

III. THE $SO(1,4)$ FLUX-BALANCE LAWS

In this section we will analyze the flux formulas at quadratic order in the quadrupolar approximation. We will use the quadrupolar solution derived in [57]. In Bondi coordinates and Λ -BMS gauge, the metric reads as

$$\begin{aligned} ds^2 &= \left(\frac{\Lambda r^2}{3} - \left(\frac{1}{2} R[q] + \frac{\Lambda}{12} C_{AB} C^{AB} \right) + \frac{2M}{r} + O(r^{-2}) \right) du^2 \\ &+ 2 \left(-1 + \frac{1}{r^2} \frac{C_{AB} C^{AB}}{16} + \frac{1}{r^4} \frac{3}{16} \left(C^{AB} E_{AB} - \frac{3}{32} (C_{AB} C^{AB})^2 \right) + O(r^{-6}) \right) dudr \\ &+ 2 \left(\frac{1}{2} D^B C_{BA} + \frac{1}{r} \left(\frac{2}{3} N_A + \frac{1}{6} C_A^F D^E C_{EF} \right) + O(r^{-2}) \right) dx^A du \\ &+ \left(r^2 q_{AB} + r C_{AB} + \frac{1}{4} q_{AB} C_{CD} C^{CD} + \frac{E_{AB}}{r} + O(r^{-2}) \right) dx^A dx^B. \end{aligned} \quad (3.1)$$

At linear order, we can expand the boundary metric q_{AB} , the shear C_{AB} and the higher Bondi moment E_{AB} as

$$q_{AB} = \bar{q}_{AB} + \delta q_{AB}, \quad C_{AB} = \delta C_{AB}, \quad E_{AB} = \delta E_{AB}. \quad (3.2)$$

For convenience we introduce the notations

$$\delta q_{AB} \equiv e_{(A}^i e_{B)}^j \delta q_{ij}, \quad \delta C_{AB} \equiv e_{(A}^i e_{B)}^j \delta C_{ij}, \quad \delta E_{AB} \equiv e_{(A}^i e_{B)}^j \delta E_{ij}. \quad (3.3)$$

Here r is the radial coordinate and $\theta^A = (\theta, \phi)$ are the angular coordinates. The unit normal vector is denoted as $n^i = x^i/r$. We employ the natural basis on the unit 2-sphere $e_A = \frac{\partial}{\partial \theta^A}$ embedded in \mathbb{R}^3 with components $e_A^i = \partial n^i / \partial \theta^A$. Given the unit sphere metric $\hat{q}_{AB} = \text{diag}(1, \sin^2 \theta)$, we have $n^i e_A^i = 0$, $\hat{q}_{AB} = \delta_{ij} e_A^i e_B^j$, and $\hat{q}^{AB} e_A^i e_B^j = \perp^{ij}$, where $\perp^{ij} = \delta^{ij} - n^i n^j$ is the projector operator. Using the covariant derivative \hat{D}_A compatible with the unit sphere metric, $\hat{D}_A \hat{q}_{BC} = 0$, we have $\hat{D}_A e_B^i = \hat{D}_B e_A^i = \hat{D}_A \hat{D}_B n^i = -\hat{q}_{AB} n^i$. The transverse-traceless projector is denoted $\perp_{tt}^{ijkl} = \perp^{k(i} \perp^{j)l} - \frac{1}{2} \perp^{ij} \perp^{kl}$. The tt projected part of any symmetric tensor T_{ij} will be defined as $T_{ij}^{tt} = \perp_{tt}^{ijkl} T_{kl}$. We also use the notation $e_{(A}^i e_{B)}^j = e_{(A}^i e_{B)}^j - \frac{1}{2} \hat{q}_{AB} \perp^{ij}$ for the trace-free product of the basis vectors.

At quadrupolar order in the perturbation we can express the Bondi fields in terms of the even and odd source quadrupolar moments $Q_{ij}^{(\rho+p)}$ and J_{ij} as [57]

$$G^{-1} \delta q_{ij} = \partial_u \zeta_{ij} + 2H^2 \partial_u Q_{ij}^{(\rho+p)} + 2H^2 n_k \epsilon_{kl(i} \left(K_{j)l} + H \int^u du' K_{j)l}(u') \right), \quad (3.4)$$

$$G^{-1} \delta C_{ij} = 3\zeta_{ij} + 2(\partial_u^2 - H^2) Q_{ij}^{(\rho+p)} + 2n_k \epsilon_{kl(i} (\partial_u + H) K_{j)l}, \quad (3.5)$$

$$G^{-1} \delta E_{ij} = 2Q_{ij}^{(\rho+p)} + 2n_k \epsilon_{kl(i} J_{j)l}. \quad (3.6)$$

Here $K_{ij} = -(\partial_u - H)J_{ij}$ and ζ_{ij} obeys the differential equation

$$\partial_u^2 \zeta_{ij} - 3H^2 \zeta_{ij} = -2GH^4 Q_{ij}^{(\rho+p)}. \quad (3.7)$$

We further have

$$G^{-1} \delta M = Q^{(\rho)} - HP_{i|i} - 3n_i (P_i - HQ_i^{(\rho)} - H^2 P_{i|kk}) + (3n_i n_j - \delta_{ij}) (\partial_u^2 Q_{ij}^{(\rho+p)} - H^2 Q_{ij}^{(\rho+p)}), \quad (3.8)$$

$$G^{-1} \delta N_i = Q_i^{(\rho)} + HP_{i|kk} + n^j (\epsilon_{ijk} J_k + 2\partial_u Q_{ij}^{(\rho+p)}) - 2\epsilon_{ijk} n_j n_l (K_{kl} - HJ_{kl}), \quad (3.9)$$

where the linear Bondi mass aspect has been simplified as Eq. (5.35) of [58].

A. Energy loss

From Eq. (2.24), the flux of the charge associated with dilatation reads at quadratic order in the perturbations as

$$\dot{Q}_D^T = -\frac{3H^4}{8G} \oint_S \delta C_{MN} \delta J_{AB}^{(\Lambda)} \hat{q}^{AM} \hat{q}^{BN}. \quad (3.10)$$

Here we defined $\oint_S = \frac{1}{4\pi} \int_{S^2} d^2\Omega \sqrt{\hat{q}}$. The expression for $J_{AB}^{(\Lambda)}$ reads to linear order as [36]

$$3H^4 \delta J_{AB}^{(\Lambda)} = -\partial_u \delta N_{AB} - H^2 \left(\hat{D}_{(A} \hat{D}^C \delta C_{B)C} - \frac{1}{2} \hat{q}_{AB} \hat{D}^C \hat{D}^D \delta C_{CD} - \delta C_{AB} \right) - 3H^4 \delta E_{AB}. \quad (3.11)$$

Using the substitution (2.16), we can rewrite the flux-balance law of the dilatation (3.10) as

$$\begin{aligned} \partial_u \oint_S \delta^2 M &= -\frac{1}{8} \oint_S [\delta N_{AB} \delta N_{CD} \\ &\quad + H^2 \delta C_{AB} (\delta C_{CD} - \hat{D}_C \hat{D}^E \delta C_{DE}) \\ &\quad - 3H^4 \delta C_{AB} \delta E_{CD}] \hat{q}^{AC} \hat{q}^{BD}. \end{aligned} \quad (3.12)$$

This formula also matches with the quadratic part of Eq. (2.52) of [36] after integrating over 2-sphere. Here we used that the linear energy $\oint_S \delta M$ is separately conserved, $\partial_u \oint_S \delta M = 0$. From Eq. (3.8), the terms proportional to n_i and $3n_i n_j - \delta_{ij}$ integrate to zero over the sphere while the term independent of n_i is conserved as a consequence of the conservation of the stress-energy tensor, $\partial_u Q^{(\rho)} = -HQ^{(\rho)} = -HS_{ii} = H\partial_u P_{i|i}$ [see also Eq. (5.37) of [58]]. Therefore, the energy loss only involves the quadratic perturbation $\delta^2 M$.

1. Even parity energy loss

We are now ready to evaluate the flux-balance law of dilatations for quadratic modes. Let us first consider the even parity modes. The shear C_{ij} is then independent of the angles. We can derive

$$C_{AB} C^{AB} = \perp_{tt}^{ijkl} C_{ij} C_{kl} = C_{ij}^{tt} C_{ij}, \quad (3.13)$$

$$\hat{D}_A \hat{D}^C C_{BC} = -2(e_{(A}^i e_{B)}^j - n^i n^j \hat{q}_{AB}) C_{ij}, \quad (3.14)$$

$$C^{AB} \hat{D}_A \hat{D}^C C_{BC} = -2C_{ij}^{tt} C_{ij}. \quad (3.15)$$

Equation (3.11) simplifies to

$$\delta J_{AB}^{(\Lambda)} = -\frac{1}{3H^4} \partial_u \delta N_{AB} + \frac{1}{H^2} \delta C_{AB} - \delta E_{AB}. \quad (3.16)$$

Using the identities (3.13), (3.14), (3.15), the energy loss formula (3.12) becomes

$$\partial_u \oint_S \delta^2 M = -\frac{1}{8G} \oint_S (\partial_u \delta C_{ij}^{\text{tt}} \partial_u \delta C_{ij} + 3H^2 \delta C_{ij}^{\text{tt}} \delta C_{ij} - 3H^4 \delta C_{ij}^{\text{tt}} \delta E_{ij}). \quad (3.17)$$

Using the even-parity moments from (3.5), (3.6), the integrand of (3.17) can be expanded as

$$\begin{aligned} & G^{-2} (\partial_u \delta C_{ij}^{\text{tt}} \partial_u \delta C_{ij} + 3H^2 \delta C_{ij}^{\text{tt}} \delta C_{ij} - 3H^4 \delta C_{ij}^{\text{tt}} \delta E_{ij}) \\ &= \partial_u (9 \zeta_{ij}^{\text{tt}} \dot{\zeta}_{ij} + 12 \ddot{Q}_{ij}^{(\rho+p)} \zeta_{ij}^{\text{tt}} - 12 H^2 Q_{ij}^{(\rho+p)} \dot{\zeta}_{ij}^{\text{tt}} - 8 H^2 \ddot{Q}_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}} - 12 H^4 Q_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}}) \\ & \quad + 4 (\ddot{Q}_{ij}^{(\rho+p)} \ddot{Q}_{ij}^{(\rho+p)\text{tt}} + 5 H^2 \ddot{Q}_{ij}^{(\rho+p)} \ddot{Q}_{ij}^{(\rho+p)\text{tt}} + 4 H^4 \dot{Q}_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}}). \end{aligned}$$

We can finally rewrite the flux-balance law of dilatations as

$$\partial_u Q_D^e(u) = -\frac{G}{2} \oint_S (\ddot{Q}_{ij}^{(\rho+p)} \ddot{Q}_{ij}^{(\rho+p)\text{tt}} + 5 H^2 \ddot{Q}_{ij}^{(\rho+p)} \ddot{Q}_{ij}^{(\rho+p)\text{tt}} + 4 H^4 \dot{Q}_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}}), \quad (3.18)$$

where the dilatation charge in the even sector is identified as

$$Q_D^e = G \oint_S \left(\frac{1}{G} \delta M + \frac{1}{G} \delta^2 M + \frac{9}{8} \zeta_{ij}^{\text{tt}} \dot{\zeta}_{ij} + \frac{3}{2} \ddot{Q}_{ij}^{(\rho+p)} \zeta_{ij}^{\text{tt}} - \frac{3}{2} H^2 Q_{ij}^{(\rho+p)} \dot{\zeta}_{ij}^{\text{tt}} - H^2 \ddot{Q}_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}} - \frac{3}{2} H^4 Q_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}} \right). \quad (3.19)$$

The flux of energy is negative definite as it should be. Using the identity

$$\oint \perp_{\text{tt}}^{ijkl} = \frac{2}{15} (3 \delta^{k(i} \delta^{j)l} - \delta^{ij} \delta^{kl}), \quad (3.20)$$

the flux-balance law in (3.18) becomes

$$\partial_u Q_D^e(u) = -\frac{G}{5} \left((\ddot{Q}_{ij}^{(\rho+p)} - \frac{1}{3} \delta_{ij} \ddot{Q}^{(\rho+p)})^2 + 5 H^2 \left(\ddot{Q}_{ij}^{(\rho+p)} - \frac{1}{3} \delta_{ij} \ddot{Q}^{(\rho+p)} \right)^2 + 4 H^4 \left(\dot{Q}_{ij}^{(\rho+p)} - \frac{1}{3} \delta_{ij} \dot{Q}^{(\rho+p)} \right)^2 \right). \quad (3.21)$$

2. Odd-parity energy loss

Let us now consider the odd-parity modes. Denoting $C_{AB} = e^i_{\langle A} e^j_{B \rangle} C_{ij}$ with $C_{ij} = \epsilon_{ikl} n_k T_{jl}$, we now have

$$C_{AB} C^{AB} = \perp_{\text{tt}}^{ijkl} C_{ij} C_{kl} = C_{ij}^{\text{tt}} C_{ij}, \quad (3.22)$$

$$\dot{D}_A \dot{D}^C C_{BC} = (4 n^i e^j_{[A} e^k_{B]} - 2 n_k e^i_{(A} e^j_{B)}) \epsilon_{ikl} T_{jl}, \quad (3.23)$$

$$\dot{D}_{(A} \dot{D}^C C_{B)C} = -2 C_{AB}, \quad (3.24)$$

$$C^{AB} \dot{D}_A \dot{D}^C C_{BC} = -2 C_{ij}^{\text{tt}} C_{ij}. \quad (3.25)$$

Equation (3.11) again simplifies to Eq. (3.16) and the energy loss formula (3.12) becomes again (3.17). Using the odd-parity moments from (3.5), (3.6), the integrand of (3.17) becomes

$$\begin{aligned} & G^{-2} (\partial_u \delta C_{ij}^{\text{tt}} \partial_u \delta C_{ij} + 3H^2 \delta C_{ij}^{\text{tt}} \delta C_{ij} - 3H^4 \delta C_{ij}^{\text{tt}} \delta E_{ij}) \\ &= -8H^2 \partial_u (\dot{J}_{ij}^{\text{tt}} \dot{J}_{ij}) - 12H^4 \partial_u (J_{ij}^{\text{tt}} \dot{J}_{ij}) \\ & \quad + 4 (\ddot{J}_{ij}^{\text{tt}} \ddot{J}_{ij} + 5H^2 \dot{J}_{ij}^{\text{tt}} \dot{J}_{ij} + 4H^4 J_{ij}^{\text{tt}} \dot{J}_{ij}). \end{aligned}$$

We can finally rewrite the flux-balance law of dilatations in the odd sector as

$$\partial_u Q_D^o(u) = -\frac{G}{2} \oint_S (\ddot{J}_{ij}^{\text{tt}} \ddot{J}_{ij} + 5H^2 \dot{J}_{ij}^{\text{tt}} \dot{J}_{ij} + 4H^4 J_{ij}^{\text{tt}} \dot{J}_{ij}), \quad (3.26)$$

where the dilatation charge is identified as

$$Q_D^o = \oint_S \left(\frac{1}{G} \delta^2 M - GH^2 \dot{J}_{ij}^{\text{tt}} \dot{J}_{ij} - \frac{3GH^4}{2} J_{ij}^{\text{tt}} \dot{J}_{ij} \right). \quad (3.27)$$

Finally, the total dilation charge Q_D is the sum of the even and odd sectors:

$$\begin{aligned} Q_D = G \oint_S \left(\frac{1}{G} \delta M + \frac{1}{G} \delta^2 M + \frac{9}{8} \zeta_{ij}^{\text{tt}} \zeta_{ij} + \frac{3}{2} \ddot{Q}_{ij}^{(\rho+p)} \zeta_{ij}^{\text{tt}} - \frac{3}{2} H^2 Q_{ij}^{(\rho+p)} \zeta_{ij}^{\text{tt}} - H^2 \ddot{Q}_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}} \right. \\ \left. - \frac{3}{2} H^4 Q_{ij}^{(\rho+p)} \dot{Q}_{ij}^{(\rho+p)\text{tt}} - GH^2 \ddot{J}_{ij}^{\text{tt}} J_{ij} - \frac{3GH^4}{2} J_{ij}^{\text{tt}} J_{ij} \right). \end{aligned} \quad (3.28)$$

It obeys the dilatation flux balance law

$$\dot{Q}_D = \partial_u Q_D^e + \partial_u Q_D^o, \quad (3.29)$$

where the even and odd energy fluxes are given in Eqs. (3.18), (3.26).

B. Angular momentum loss

From (2.24), the quadratic part of the angular momentum loss formula becomes

$$\dot{Q}_{L^i}^T = -\frac{3H^2}{8G} \oint_S \dot{q}^{AM} \dot{q}^{BN} \mathcal{L}_{\xi(i)}^{\circ} \delta q_{MN} \delta J_{AB}^{(\Lambda)}. \quad (3.30)$$

Using Eq. (3.16), the angular momentum loss formula can be written as

$$\begin{aligned} \partial_u \left(Q_{L^i}^T - \frac{1}{8GH^2} \oint_S (\mathcal{L}_{\xi(i)}^{\circ} \delta q_{MN}) \partial_u \delta C_{AB} q^{\circ AM} q^{\circ BN} \right) \\ = -\frac{1}{8G} \oint_S [(\mathcal{L}_{\xi(i)}^{\circ} \delta C_{MN}) \partial_u \delta C_{AB} \\ + 3(\mathcal{L}_{\xi(i)}^{\circ} \delta q_{MN}) \delta C_{AB} - 3H^2 (\mathcal{L}_{\xi(i)}^{\circ} \delta q_{MN}) \delta E_{AB}] q^{\circ AM} q^{\circ BN}. \end{aligned} \quad (3.31)$$

Let us first discuss the even-parity modes. We note the identities $e^{AB} = e_i^A e_j^B n_k \epsilon_{ijk}$, $\oint_S 3n_i n_j = \delta_{ij}$ and $\oint_S n_i n_j n_k n_l = \frac{1}{15} (\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk})$. After performing the integral over the sphere, the right-hand side of Eq. (3.31) becomes

$$-\frac{1}{10G} \epsilon_i^{mn} (\delta C_{mk} \partial_u \delta C_{kn} + 3\delta q_{mk} \delta C_{kn} - 3H^2 \delta q_{mk} \delta E_{kn}). \quad (3.32)$$

After some algebra we can rewrite the latter expression as

$$\begin{aligned} -\frac{G}{10} \epsilon_i^{mn} \partial_u (6\zeta_{km} \ddot{Q}_{kn}^{(\rho+p)} - 6H^2 \zeta_{km} Q_{kn}^{(\rho+p)} + 6\dot{\zeta}_{kn} \dot{Q}_{km}^{(\rho+p)} - 4H^2 Q_{km} \ddot{Q}_{kn}^{(\rho+p)}) \\ -\frac{2G}{5} \epsilon_i^{mn} (\ddot{Q}_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 5H^2 \dot{Q}_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)} + 4H^4 Q_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)}). \end{aligned} \quad (3.33)$$

Let us now consider the odd-parity modes. After performing the integral over the sphere and reorganizing the terms, the right-hand side of Eq. (3.31) becomes

$$-\frac{G}{10} \epsilon_i^{mn} \left[\partial_u \left(-4H^2 J_{km} \ddot{J}_{kn} - 12H^4 \int^u J_{mk} \dot{J}_{kn} \right) + 4(\ddot{J}_{km} \ddot{J}_{kn} + 5H^2 \dot{J}_{km} \dot{J}_{kn} + 4H^4 J_{km} \dot{J}_{kn}) \right]. \quad (3.34)$$

Combining both even- and odd-parity sectors, the flux-balance law of angular momentum finally reads as

$$\partial_u Q_{L^i} = -\frac{2G}{5} \epsilon_{imn} (\ddot{Q}_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 5H^2 \dot{Q}_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)} + 4H^4 Q_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)} + \ddot{J}_{km} \ddot{J}_{kn} + 5H^2 \dot{J}_{km} \dot{J}_{kn} + 4H^4 J_{km} \dot{J}_{kn}), \quad (3.35)$$

where the final angular momentum charge (in both parity sectors) is

$$\begin{aligned} Q_{L^i} = Q_{L^i}^T - \frac{1}{8GH^2} \oint_S (\mathcal{L}_{\xi(i)}^{\circ} \delta q_{MN}) \partial_u \delta C_{AB} q^{\circ AM} q^{\circ BN} + \frac{G}{10} \epsilon_{imn} \left(6\zeta_{km} \ddot{Q}_{kn}^{(\rho+p)} - 6H^2 \zeta_{km} Q_{kn}^{(\rho+p)} + 6\dot{\zeta}_{kn} \dot{Q}_{km}^{(\rho+p)} - 4H^2 Q_{km} \ddot{Q}_{kn}^{(\rho+p)} \right. \\ \left. - 4H^2 J_{km} \ddot{J}_{kn} - 12H^4 \int^u J_{km} \dot{J}_{kn} \right). \end{aligned} \quad (3.36)$$

C. Linear momentum and boost charge losses

From Eq. (2.24) with $\delta\xi^a = 0$, the quadratic part of the linear momentum loss and cosmological boost formula becomes

$$\dot{Q}_{\xi(i)}^T = -\frac{3H^2}{8G} \oint_S (H^2 \bar{\xi}_{(i)}^u \delta C_{MN} + \mathcal{L}_{\xi}^i \delta q_{MN} - \dot{D}_C \bar{\xi}_{(i)}^C \delta q_{MN}) \delta J_{AB}^{(\Lambda)} \hat{q}^{AM} \hat{q}^{BN}, \quad (3.37)$$

where $\bar{\xi}_{(i)}^u = n_i \exp(\varepsilon Hu)$; $\bar{\xi}_{(i)}^A = -\varepsilon H \exp(\varepsilon Hu) \dot{D}_A n_i$. For linear momenta $\varepsilon = 1$ while for cosmological boosts $\varepsilon = -1$. We simplify the expression as

$$\dot{Q}_{\xi(i)}^T = -\frac{3H^2}{8G} \oint_S \exp(\varepsilon Hu) (H^2 n_i \delta C_{MN} - \varepsilon H (\dot{D}^C n_i) (\dot{D}_C \delta q_{MN})) \delta J_{AB}^{(\Lambda)} \hat{q}^{AM} \hat{q}^{BN}. \quad (3.38)$$

In order to perform the angular integrals, we note that for any symmetric trace-free tensors V_{AB} , W_{AB} that can be decomposed into even and odd parts as

$$V_{AB} = e_{(A}^i e_{B)}^j (V_{ij}^e + \varepsilon_{ikl} n_k V_{lj}^o), \quad (3.39)$$

$$W_{AB} = e_{(A}^i e_{B)}^j (W_{ij}^e + \varepsilon_{ikl} n_k W_{lj}^o), \quad (3.40)$$

we have the integrals

$$\oint_S n^i V_{AB} W^{AB} = -\frac{4}{15} \varepsilon_{imt} (V_{ml}^e W_{il}^o + W_{ml}^e V_{il}^o), \quad (3.41)$$

$$\oint_S e_i^C \dot{D}_C V_{AB} W^{AB} = -\frac{4}{15} \varepsilon_{imt} (V_{ml}^e W_{il}^o + W_{ml}^e V_{il}^o). \quad (3.42)$$

Therefore,

$$\dot{Q}_{\xi(i)}^T = \frac{H^2}{10G} \exp(\varepsilon Hu) \varepsilon_{imt} [(H^2 \delta C_{ml}^e - \varepsilon H \delta q_{ml}^e) \delta J_{il}^{(\Lambda)o} + (H^2 \delta C_{il}^o - \varepsilon H \delta q_{il}^o) \delta J_{ml}^{(\Lambda)e}]. \quad (3.43)$$

We now substitute $\delta J_{ij}^{(\Lambda)}$ using Eq. (3.16) and integrate by parts to obtain

$$\begin{aligned} \dot{Q}_{\xi(i)}^{\text{int}} = & \frac{\exp(\varepsilon Hu)}{10G} \varepsilon_{imt} \left[\frac{1}{3} (2\delta N_{ml}^e \delta N_{il}^o + 6H^2 \delta C_{ml}^e \delta C_{il}^o - 3H^4 \delta C_{ml}^e \delta E_{il}^o - 3H^4 \delta E_{ml}^e \delta C_{il}^o) \right. \\ & \left. - \frac{\varepsilon H}{3} \delta q_{ml}^e (3\delta C_{il}^o - 3H^2 \delta E_{il}^o) - \frac{\varepsilon H}{3} \delta q_{il}^o (3\delta C_{ml}^e - 3H^2 \delta E_{ml}^e) - \frac{1}{3} (\delta q_{ml}^e \delta N_{il}^o + \delta N_{ml}^e \delta q_{il}^o) \right], \end{aligned} \quad (3.44)$$

where the charge (defined as an intermediate quantity for now, hence the superscript) is

$$Q_{\xi(i)}^{\text{int}} = Q_{\xi(i)}^T + \frac{1}{30G} \exp(\varepsilon Hu) \varepsilon_{imt} \left[\delta C_{ml}^e \delta N_{il}^o + \delta N_{ml}^e \delta C_{il}^o - \frac{\varepsilon}{H} (\delta q_{ml}^e \delta N_{il}^o + \delta N_{ml}^e \delta q_{il}^o) \right]. \quad (3.45)$$

Substituting all Bondi fields in terms of multipolar moments, we rewrite the flux-balance law as

$$\dot{Q}_{\xi(i)} = -\frac{4Ge^{\varepsilon Hu}}{15} \varepsilon_{imt} (3\varepsilon H^3 \zeta_{ml} \dot{J}_{lt} + 2H^6 Q_{ml} J_{lt} + 6H^4 \dot{Q}_{ml} \dot{J}_{lt} + 6H^2 \ddot{Q}_{ml} \ddot{J}_{lt} + \ddot{\ddot{Q}}_{ml} \ddot{\ddot{J}}_{lt}), \quad (3.46)$$

where the final charge is defined as

$$\begin{aligned} Q_{\xi(i)} = & Q_{\xi(i)}^{\text{int}} + \frac{e^{\varepsilon Hu}}{15} \varepsilon_{imt} \left[6H^2 \partial_u (\dot{Q}_{ml} \dot{J}_{lt}) + 8H^4 \partial_u (Q_{ml} J_{lt}) - 4\varepsilon H^5 \partial_u \left(Q_{ml} \int^u J_{lt} \right) \right. \\ & \left. - 5\zeta_{ml} \dot{J}_{lt} + 3H^2 (\zeta_{ml} J_{lt} - \zeta_{ml} \dot{J}_{lt}) + 8\varepsilon H \zeta_{ml} \dot{J}_{lt} - (2H^4 \ddot{Q}_{ml} - 6H^6 Q_{ml} - 3\varepsilon H^3 \zeta_{ml}) \int^u J_{lt} \right]. \end{aligned} \quad (3.47)$$

D. Flat spacetime limit

Let us now prove that we recover the standard Poincaré flux-balance laws in the quadrupolar approximation upon taking the flat spacetime limit. In the quadrupolar truncation, these laws read as [70]¹ (original derivations can be found in [71–75])

$$\partial_u \mathcal{E} = -\frac{G}{5} \ddot{I}_{ij}^{\text{PM}} \ddot{I}_{ij}^{\text{PM}} - \frac{16G}{45} \ddot{J}_{ij}^{\text{PM}} \ddot{J}_{ij}^{\text{PM}}, \quad (3.48)$$

$$\partial_u \mathcal{P}_i = -\frac{16G}{45} \epsilon_{ijk} \ddot{I}_{jl}^{\text{PM}} \ddot{J}_{kl}^{\text{PM}}, \quad (3.49)$$

$$\partial_u \mathcal{J}_i = -\frac{2G}{5} \epsilon_{ijk} \ddot{I}_{jl}^{\text{PM}} \ddot{I}_{kl}^{\text{PM}} - \frac{32G}{45} \epsilon_{ijk} \ddot{J}_{jl}^{\text{PM}} \ddot{J}_{kl}^{\text{PM}}, \quad (3.50)$$

$$\partial_u \mathcal{N}_i = \mathcal{P}_i, \quad (3.51)$$

where the flat tensors I_{ij}^{PM} , J_{ij}^{PM} are symmetric and trace-free. The charges are expressed as the flat limit of the SO(1,4) generators as $\mathcal{E} = \mathcal{Q}_D$, $\mathcal{P}_i = \mathcal{Q}_{P^i}$, $\mathcal{J}_i = \mathcal{Q}_{L^i}$ and $\mathcal{K}_i = \lim_{H \rightarrow 0} \frac{1}{2H} (\mathcal{Q}_{K^i} - \mathcal{Q}_{P^i})$ with $\mathcal{N}_i = \mathcal{K}_i + u\mathcal{P}_i$.

The flat limit of Eqs. (3.21), (3.26), (3.35) reproduce the flat flux-balance laws of energy and angular momentum with the dictionary

$$I_{ij}^{\text{PM}} = \mathcal{Q}_{ij}^{(\rho+p)} - \frac{1}{3} \delta_{ij} \mathcal{Q}_{kk}^{(\rho+p)}, \quad (3.52)$$

$$J_{ij}^{\text{PM}} = \frac{3}{4} J_{ij}. \quad (3.53)$$

The flat limit of Eq. (3.46) reproduces the flat flux-balance law for the momentum. For Lorentz boosts we have to extract the $O(H)$ term in Eq. (3.44). Note that $\delta q_{ab} = O(H^2)$, therefore, only the $\exp(\epsilon H u) \delta N_{mi}^e \delta N_{il}^o$ term in Eq. (3.44) will contribute to the boost charge. From Eq. (3.46) we obtain

$$\dot{\mathcal{K}}_i = \lim_{H \rightarrow 0} \frac{1}{2H} (\dot{\mathcal{Q}}_{K^i} - \dot{\mathcal{Q}}_{P^i}) = \frac{4Gu}{15} \epsilon_{iml} \ddot{\mathcal{Q}}_{mj} \ddot{\mathcal{J}}_{jl}. \quad (3.54)$$

We can finally shift the charge as

$$\mathcal{N}_i = \mathcal{K}_i - \frac{4Gu}{15} \int^u \epsilon_{iml} (\ddot{\mathcal{Q}}_{mj} \ddot{\mathcal{J}}_{jl}) \quad (3.55)$$

to obtain the correct flux-balance law (3.51).

IV. COMPARISON WITH THE LITERATURE

The first proposed quadrupolar formula for the energy flux for even-parity modes was derived in [7], see also [17]. In our earlier work [57], we pointed out that there is an inconsistent quadrupolar truncation in [7,17], which leads to incorrect

¹Note that \mathcal{G}_i in [70] is \mathcal{N}_i in our conventions, see Sec. 9 of [68].

terms proportional to $Q_{ij}^{(p)}$ even in the even-parity sector. Performing the quadrupolar truncation using the methods of [57], let us now upgrade the flux formula of [7] to include the $Q_{ij}^{(p)}$ terms. We first note the algebraic identity

$$\begin{aligned} \ddot{\mathcal{Q}}_{ij}^{(\rho+p)} \ddot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)} + 5H^2 \ddot{\mathcal{Q}}_{ij}^{(\rho+p)} \ddot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)} + 4H^4 \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)} \\ = \mathcal{R}_{ij}^e \mathcal{R}_{ij}^{\text{tt}} - \partial_u \delta E^e \end{aligned},$$

where we defined

$$\mathcal{R}_{ij}^e = \ddot{\mathcal{Q}}_{ij}^{(\rho+p)} - 3H \dot{\mathcal{Q}}_{ij}^{(\rho+p)} + 2H^2 \dot{\mathcal{Q}}_{ij}^{(\rho+p)}, \quad (4.1)$$

$$\begin{aligned} \delta E^e = -3H \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)} + 4H^2 \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)} \\ - 6H^3 \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{\text{tt}(\rho+p)}. \end{aligned} \quad (4.2)$$

The same identity holds in the odd sector with $Q_{ij}^{(\rho+p)}$ replaced by J_{ij} , namely,

$$\ddot{J}_{ij} \ddot{J}_{ij}^{\text{tt}} + 5H^2 \dot{J}_{ij} \dot{J}_{ij}^{\text{tt}} + 4H^4 J_{ij} J_{ij}^{\text{tt}} = \mathcal{R}_{ij}^o \mathcal{R}_{ij}^{\text{tt}} - \partial_u \delta E^o,$$

where we defined

$$\mathcal{R}_{ij}^o = \ddot{J}_{ij} - 3H \dot{J}_{ij} + 2H^2 J_{ij}, \quad (4.3)$$

$$\delta E^o = -3H \dot{J}_{ij} \dot{J}_{ij}^{\text{tt}} + 4H^2 \dot{J}_{ij} \dot{J}_{ij}^{\text{tt}} - 6H^3 J_{ij} J_{ij}^{\text{tt}}. \quad (4.4)$$

The energy flux-balance law (3.29) can therefore be equivalently defined as

$$\begin{aligned} \partial_u \left(\mathcal{Q}_D + \frac{G}{2} \oint_S \delta(E^e + E^o) \right) \\ = -\frac{G}{2} \oint_S (\mathcal{R}_{ij}^e \mathcal{R}_{ij}^{\text{tt}} + \mathcal{R}_{ij}^o \mathcal{R}_{ij}^{\text{tt}}) \\ = -\frac{G}{5} (\mathcal{R}_{ij}^e \mathcal{R}_{ij}^e + \mathcal{R}_{ij}^o \mathcal{R}_{ij}^o). \end{aligned} \quad (4.5)$$

The right-hand side is negative definite. It depends upon the combination $Q_{ij}^{(\rho+p)}$, consistently with the Bondi fields (3.4), (3.5), (3.6), (3.8). The energy flux (4.5) is our upgrade of the formula proposed in [7].²

²Note that the power radiated quadrupolar formula in [7] was given in terms of the transverse-traceless (TT) part of \mathcal{R}_{ij} field. Let us recall that any symmetric rank-2 tensor can be decomposed as $Q_{ij} = \frac{1}{3} \delta_{ij} \delta^{kl} Q_{kl} + (\partial_i \partial_j - \frac{1}{3} \delta_{ij} \nabla^2) B + \partial_i B_j^T + \partial_j B_i^T + Q_{ij}^{\text{TT}}$ where Q_{ij}^{TT} is the TT part of Q_{ij} , which satisfies $\partial^j Q_{ij}^{\text{TT}} = 0 = \delta^{ij} Q_{ij}^{\text{TT}}$. The two notions, tt projection and TT part of a rank-2 symmetric tensor are therefore generically distinct. However, this distinction disappears after integrating over the two sphere at infinity [60,76]. Note in addition that in [7], \dot{Q}_{ij} denotes the Lie derivative with respect to dilatation Killing vector field (T) in conformal coordinates of de Sitter, i.e. $\dot{Q}_{ij} = \mathcal{L}_T Q_{ij}$. After conversion to Bondi frame, it becomes $\mathcal{L}_T Q_{ij} = (\partial_u - 2H) Q_{ij}$.

We can also upgrade the angular momentum flux proposed in [7]. Again we note the identity

$$\ddot{Q}_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 5H^2 \dot{Q}_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 4H^4 Q_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)} = \mathcal{L}_{km}^e \mathcal{R}_{kn}^e - \partial_u \delta L^e, \quad (4.6)$$

where we defined

$$\mathcal{L}_{km}^e = \ddot{Q}_{km}^{(\rho+p)} - 3H \dot{Q}_{km}^{(\rho+p)} + 2H^2 Q_{km}^{(\rho+p)}, \quad (4.7)$$

$$\delta L^e = -3H \dot{Q}_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 2H^2 Q_{km}^{(\rho+p)} \ddot{Q}_{kn}^{(\rho+p)} + 2H^2 \dot{Q}_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)} - 6H^3 Q_{km}^{(\rho+p)} \dot{Q}_{kn}^{(\rho+p)}. \quad (4.8)$$

The same identity holds in the odd sector with $Q_{ij}^{(\rho+p)}$ replaced by J_{ij} ,

$$\ddot{J}_{km} \ddot{J}_{kn} + 5H^2 \dot{J}_{km} \ddot{J}_{kn} + 4H^4 J_{km} \dot{J}_{kn} = \mathcal{L}_{km}^o \mathcal{R}_{kn}^o - \partial_u \delta L_T^o, \quad (4.9)$$

where we defined

$$\mathcal{L}_{km}^o = \ddot{J}_{km} - 3H \dot{J}_{km} + 2H^2 J_{km}, \quad (4.10)$$

$$\delta L^o = -3H \dot{J}_{km} \ddot{J}_{kn} + 2H^2 J_{km} \ddot{J}_{kn} + 2H^2 \dot{J}_{km} \dot{J}_{kn} - 6H^3 J_{km} \dot{J}_{kn}. \quad (4.11)$$

Hence the angular momentum flux-balance law (3.35) can also be equivalently defined as

$$\begin{aligned} \partial_u \left(Q_{Li} - \frac{2G}{5} \epsilon_{imn} (\delta L^e + \delta L_T^o) \right) \\ = -\frac{2G}{5} \epsilon_{imn} (\mathcal{L}_{km}^e \mathcal{R}_{kn}^e + \mathcal{L}_{km}^o \mathcal{R}_{kn}^o). \end{aligned} \quad (4.12)$$

This is our upgrade of the angular momentum flux proposed in [7].

In anti-de Sitter spacetime with Dirichlet boundary conditions, the $SO(3,2)$ conserved quantities are defined using the holographic stress-energy tensor [77]. More precisely, they are defined from the analytic continuation to negative cosmological constant of the charge formula (2.10). We started from this charge formula as initial ansatz in the case of positive cosmological constant in this paper. When rewritten in Bondi variables, the charge can be expressed in terms of the Λ -corrected Bondi mass aspects $M^{(\Lambda)}$ and Bondi angular momentum aspects $N_A^{(\Lambda)}$ while the flux is proportional to the field $J_{AB}^{(\Lambda)}$ [36,39], see Eq. (2.19). However, for leaky boundary conditions, where the boundary metric is allowed to vary, as it is the case here, the holographic stress tensor does not allow by itself to define the quasiconserved quantities. Indeed, computing the flux

of $M^{(\Lambda)}$ (2.16) using Eq. (3.17) shows that the “would be energy flux” associated with the Bondi mass is not manifestly negative definite. It cannot therefore lead to a proposal for a definition of energy in the presence of a positive cosmological constant.

Bonga, Bunster, and Pérez (BBP) [55] studied several aspects of gravitational waves in de Sitter for both linear and the nonlinear theory. For the study of linearized solutions they focused on solving the homogeneous Einstein equations in a partial wave expansion to the gravitational field. One of the results of their paper is the derivation of an energy flux formula for $l=2$ mode of linearized perturbations around de Sitter. Starting from the energy loss formula in a different boundary gauge fixing introduced in [38,44], they obtained a quadratic energy flux formula for linearized perturbations in de Sitter background. In [58], an energy flux formula for de Sitter Teukolsky waves has also been obtained. With the correct identification of the variables, the energy flux formula for BBP quadrupolar waves exactly matches with that of de Sitter Teukolsky waves. In this section, we wish to compare our quadrupole formula with the energy flux formula for BBP quadrupolar waves in de Sitter [55].

For $l=2$ even-parity linearized solutions of [55], the energy flux is given by

$$\begin{aligned} E^{(E)} = -\frac{3}{8\pi G} \sum_{m=-2}^{m=+2} \int_{u=-\infty}^{u=+\infty} [|\ddot{a}_m|^2 + 5H^2 |\dot{a}_m|^2 \\ + 4H^4 |a_m|^2] du, \end{aligned} \quad (4.13)$$

where $a_m = a_m(u)$ is a function of retarded time u . The function $a_m(u)$ can also be understood as the coefficient of the $l=2$ even parity solution of homogeneous linearized Einstein’s equation. The relationship between the variables a_m used in [55] and our even parity quadrupole moments are obtained from [58], where a relation between the BBP wave solutions and $l=2$ mode quadrupolar truncated solutions [57] have been achieved via de Sitter Teukolsky waves. Using Eqs. (3.55), (5.79) of [58], we obtain

$$a_m = \frac{H^2}{2} q_m - \frac{1}{6} \ddot{q}_m, \quad (4.14)$$

where q_m can be written in terms of $l=2$ spherical harmonics $Y^{2m}(\theta, \phi) = \mathcal{Y}_{ij}^{2m} n_i n_j$,

$$q_m = \frac{1}{H^4} \frac{24\pi}{15} (\mathcal{Y}_{ij}^{2m})^* \zeta_{ij}(u). \quad (4.15)$$

An explicit expression for \mathcal{Y}_{ij}^{2m} can be found in Eq. (3.218) of [78]. Therefore using (4.14), and the identity (3.7), we have

$$a_m = \frac{8\pi G}{15} (\mathcal{Y}_{ij}^{2m})^* \mathcal{Q}_{ij}^{(\rho+p)}. \quad (4.16)$$

where we have used the identity,

Therefore,

$$\sum_{m=-2}^{m=+2} |\ddot{a}_m|^2 = \frac{8\pi G^2}{15} \left(\ddot{\mathcal{Q}}_{ij}^{(\rho+p)} \ddot{\mathcal{Q}}_{ij}^{(\rho+p)} - \frac{1}{3} \ddot{\mathcal{Q}}_{ij}^{(\rho+p)} \ddot{\mathcal{Q}}_{kl}^{(\rho+p)} \delta_{ij} \delta_{kl} \right), \quad (4.17)$$

$$\sum_{m=-2}^{m=+2} (\mathcal{Y}_{ij}^{2m})^* (\mathcal{Y}_{kl}^{2m}) = \frac{15}{16\pi} \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{3} \delta_{ij} \delta_{kl} \right). \quad (4.19)$$

$$= \frac{8\pi G^2}{15} \left(\ddot{\mathcal{Q}}_{ij}^{(\rho+p)} - \frac{1}{3} \ddot{\mathcal{Q}}^{(\rho+p)} \delta_{ij} \right)^2, \quad (4.18)$$

Hence the energy flux in terms of even-parity quadrupole moments becomes

$$E^{(E)} = -\frac{G}{5} \int_{-\infty}^{+\infty} du \left(\left(\ddot{\mathcal{Q}}_{ij}^{(\rho+p)} - \frac{1}{3} \ddot{\mathcal{Q}}^{(\rho+p)} \delta_{ij} \right)^2 + 5H^2 \left(\dot{\mathcal{Q}}_{ij}^{(\rho+p)} - \frac{1}{3} \dot{\mathcal{Q}}^{(\rho+p)} \delta_{ij} \right)^2 + 4H^4 \left(\dot{\mathcal{Q}}_{ij}^{(\rho+p)} - \frac{1}{3} \dot{\mathcal{Q}}^{(\rho+p)} \delta_{ij} \right)^2 \right). \quad (4.20)$$

This can also be written as

$$E^{(E)} = -\frac{G}{8\pi} \int_{-\infty}^{+\infty} du \int_{S^2} d\Omega \left(\ddot{\mathcal{Q}}_{ij}^{(\rho+p)} \ddot{\mathcal{Q}}_{ij}^{(\rho+p)tt} + 5H^2 \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{(\rho+p)tt} + 4H^4 \dot{\mathcal{Q}}_{ij}^{(\rho+p)} \dot{\mathcal{Q}}_{ij}^{(\rho+p)tt} \right). \quad (4.21)$$

Considering now the $l = 2$ odd-parity linearized solution, the energy flux is given by

$$E^{(B)} = -\frac{3}{8\pi G} \sum_{m=-2}^{m=+2} \int_{u=-\infty}^{u=+\infty} [|\ddot{b}_m|^2 + 5H^2 |\dot{b}_m|^2 + 4H^4 |\dot{b}_m|^2] du, \quad (4.22)$$

where $b_m = b_m(u)$ is a function of retarded time u . The relationship between the variable b_m used in [55] and our odd-parity $l = 2$ mode quadrupole moments are given by [58]

$$b_m = -\frac{8\pi G}{15} (\mathcal{Y}_{ij}^{2m})^* J_{ij}. \quad (4.23)$$

Therefore using (4.19), we have

$$\sum_{m=-2}^{m=+2} |\ddot{b}_m|^2 = \frac{8\pi G^2}{15} \ddot{J}_{ij} \ddot{J}_{ij}. \quad (4.24)$$

Note that by construction J_{ij} is symmetric and traceless. Hence the energy flux in terms of odd-parity quadrupole moments becomes

$$E^{(B)} = -\frac{G}{5} \int_{-\infty}^{+\infty} du \left(\ddot{J}_{ij} \ddot{J}_{ij} + 5H^2 \dot{J}_{ij} \dot{J}_{ij} + 4H^4 J_{ij} J_{ij} \right). \quad (4.25)$$

This can also be written as

$$E^{(B)} = -\frac{G}{8\pi} \int_{-\infty}^{+\infty} du \int_{S^2} d\Omega \left(\ddot{J}_{ij} \ddot{J}_{ij}^{tt} + 5H^2 \dot{J}_{ij} \dot{J}_{ij}^{tt} + 4H^4 J_{ij} J_{ij}^{tt} \right). \quad (4.26)$$

We have therefore demonstrated that the energy flux formulas (3.18), (3.26) exactly coincide with the ones derived in [55]. This is remarkable because these formulas have been obtained in different boundary gauges. In [55], the angular components g_{AB} of the boundary metric were kept fixed, while here we kept fixed the determinant of g_{AB} and the mixed components g_{uA} . This does not prove that the energy flux formulas are gauge invariant, but they are at least invariant under an interesting class of gauge transformations.

Given that the quadratic energy loss formula derived in [38,41] agree with the one derived in [55] as shown in [55] we also proved the equivalence of the energy flux formulas (3.18), (3.26) with [38,41].

V. CONCLUSION

We derived the full set of flux-balance laws for linear (even and odd) quadrupolar perturbations associated with background $SO(1,4)$ symmetries. These laws reduce to the standard Poincaré flux-balance laws in the flat limit. Even though the flux-balance laws have been determined, their split into the time derivative of the charge (left-hand side) and the flux (right-hand side) is not unique. We warn the reader that even if the symmetry generators form a $SO(1,4)$ algebra, we did not prove that the charges form a $SO(1,4)$ algebra under a suitable bracket. As proven in [39], the

charges Q_{ξ}^T defined from the holographic stress-energy tensor represent the Λ -BMS algebra under the adjusted bracket [69]. However, since the charges are here defined with a shift (2.25), one would need to study the impact of this shift on the representation theorem. This remains an open question.

After analysis, we obtained two distinct proposals for the energy loss. Both proposals admit the standard flat limit and are negative definite at quadratic level. They differ by terms proportional to the Hubble radius. The first proposal is given in Eq. (4.5). It corresponds to an upgrade of the formula first proposed in [7] after correcting the even-parity sector following [57], and after including the odd-parity contribution. The second proposal is given as Eq. (3.29). After integrating over \mathcal{I}^+ , it matches with the total energy loss formula derived in [55,58]. We further complemented the previous literature [55,58] by providing the corresponding definition of energy and energy flux on a fixed cut (i.e. fixed u) of \mathcal{I}^+ in terms of multipolar moments. The existence of two proposals for the energy loss indicates that fundamental requirements are missing to uniquely single out one expression. One such requirement is gauge invariance. The Bondi framework is based upon one gauge choice and even though it was successful in the asymptotically flat regime, we expect that a gauge-independent framework is required to settle this issue. The comparison of proposed definitions of energy loss based on Weyl

scalars and Killing spinors [9,13,23] with the current framework is left for future work.

ACKNOWLEDGMENTS

We gratefully thank A. Pérez and A. Virmani for their comments on the draft prior to submission. G.C. is Research Director of the F.R.S.-FNRS. The work of J.H. is supported in part by MSCA Fellowships CZ—UK2 (reg. n. CZ.02.01.01/00/22_010/0008115) from the Programme Johannes Amos Comenius co-funded by the European Union. J.H. also acknowledges the support from Czech Science Foundation Grant No. 22-14791S. E. S. K. is supported by Scuola Normale Superiore and partially by INFN (IS-GSS) and ERC (NOTIMEFORCOSMO, 101126304) grant funded by the European Union. E. S. K. has also been supported by TUBITAK-2219 Fellowship during the project.

Views and opinions expressed are, however, those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

DATA AVAILABILITY

No data were created or analyzed in this study.

-
- [1] A. G. Riess *et al.* (Supernova Search Team Collaboration), Observational evidence from supernovae for an accelerating universe and a cosmological constant, *Astron. J.* **116**, 1009 (1998).
 - [2] S. Perlmutter *et al.* (Supernova Cosmology Project Collaboration), Measurements of Ω and Λ from 42 high redshift supernovae, *Astrophys. J.* **517**, 565 (1999).
 - [3] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
 - [4] A. Ashtekar, B. Bonga, and A. Kesavan, Asymptotics with a positive cosmological constant: I. Basic framework, *Classical Quantum Gravity* **32**, 025004 (2015).
 - [5] C. G. Böhrer and M. Wright, Slowly rotating perfect fluids with a cosmological constant, *Gen. Relativ. Gravit.* **47**, 143 (2015).
 - [6] Y.-Z. Chu, Transverse traceless gravitational waves in a spatially flat FLRW universe: Causal structure from dimensional reduction, *Phys. Rev. D* **92**, 124038 (2015).
 - [7] A. Ashtekar, B. Bonga, and A. Kesavan, Asymptotics with a positive cosmological constant: III. The quadrupole formula, *Phys. Rev. D* **92**, 104032 (2015).
 - [8] A. Ashtekar, B. Bonga, and A. Kesavan, Gravitational waves from isolated systems: Surprising consequences of a positive cosmological constant, *Phys. Rev. Lett.* **116**, 051101 (2016).
 - [9] L. B. Szabados and P. Tod, A positive Bondi–type mass in asymptotically de Sitter spacetimes, *Classical Quantum Gravity* **32**, 205011 (2015).
 - [10] G. Date and S. J. Hoque, Gravitational waves from compact sources in a de Sitter background, *Phys. Rev. D* **94**, 064039 (2016).
 - [11] N. T. Bishop, Gravitational waves in a de Sitter universe, *Phys. Rev. D* **93**, 044025 (2016).
 - [12] P. T. Chruściel and L. Ifsits, The cosmological constant and the energy of gravitational radiation, *Phys. Rev. D* **93**, 124075 (2016).
 - [13] V.-L. Saw, Mass-loss of an isolated gravitating system due to energy carried away by gravitational waves with a cosmological constant, *Phys. Rev. D* **94**, 104004 (2016).
 - [14] A. Kesavan, Asymptotic structure of space-time with a positive cosmological constant, other thesis, Penn State U., Penn State U., 6, 2016.
 - [15] V.-L. Saw, Behavior of asymptotically electro- Λ spacetimes, *Phys. Rev. D* **95**, 084038 (2017).
 - [16] V. Schlue, Decay of the Weyl curvature in expanding black hole cosmologies, *Ann. PDE* **8**, 9 (2022).

- [17] G. Date and S. J. Hoque, Cosmological horizon and the quadrupole formula in de Sitter background, *Phys. Rev. D* **96**, 044026 (2017).
- [18] S. Viaggiu, Axial and polar gravitational wave equations in a de Sitter expanding universe by Laplace transform, *Classical Quantum Gravity* **34**, 035018 (2017).
- [19] Y. Hamada, M.-S. Seo, and G. Shiu, Memory in de Sitter space and Bondi-Metzner-Sachs-like supertranslations, *Phys. Rev. D* **96**, 023509 (2017).
- [20] S. J. Hoque, Physics of gravitational waves in presence of positive cosmological constant. Ph.D. thesis, HBNI, Mumbai, 2017.
- [21] V.-L. Saw, Mass loss due to gravitational waves with $\Lambda > 0$, *Mod. Phys. Lett. A* **32**, 1730020 (2017).
- [22] V.-L. Saw, Asymptotically simple spacetimes and mass loss due to gravitational waves, *Int. J. Mod. Phys. D* **27**, 1730027 (2017).
- [23] V.-L. Saw, Bondi mass with a cosmological constant, *Phys. Rev. D* **97**, 084017 (2018).
- [24] A. Ashtekar and B. Bonga, On the ambiguity in the notion of transverse traceless modes of gravitational waves, *Gen. Relativ. Gravit.* **49**, 122 (2017).
- [25] A. Ashtekar and B. Bonga, On a basic conceptual confusion in gravitational radiation theory, *Classical Quantum Gravity* **34**, 20LT01 (2017).
- [26] B. P. Bonga, Gravitational radiation with a positive cosmological constant. Ph.D. thesis, Penn State U., Penn State U., 6, 2017.
- [27] A. Ashtekar, Implications of a positive cosmological constant for general relativity, *Rep. Prog. Phys.* **80**, 102901 (2017).
- [28] S. Viaggiu, A possible new cosmological redshift effect due to Λ on traveling gravitational waves in Friedmann universes, *Int. J. Mod. Phys. D* **26**, 1750168 (2017).
- [29] B. Bonga and J. S. Hazboun, Power radiated by a binary system in a de Sitter Universe, *Phys. Rev. D* **96**, 064018 (2017).
- [30] S. J. Hoque and A. Virmani, On propagation of energy flux in de Sitter spacetime, *Gen. Relativ. Gravit.* **50**, 40 (2018).
- [31] L. B. Szabados and P. Tod, A review of total energy-momenta in GR with a positive cosmological constant, *Int. J. Mod. Phys. D* **28**, 1930003 (2018).
- [32] A. Poole, K. Skenderis, and M. Taylor, (A)dS₄ in Bondi gauge, *Classical Quantum Gravity* **36**, 095005 (2019).
- [33] P. Mao, Asymptotics with a cosmological constant: The solution space, *Phys. Rev. D* **99**, 104024 (2019).
- [34] A. Prema Balakrishnan, S. J. Hoque, and A. Virmani, Conserved charges in asymptotically de Sitter spacetimes, *Classical Quantum Gravity* **36**, 205008 (2019).
- [35] A. Ashtekar and S. Bahrami, Asymptotics with a positive cosmological constant. IV. The no-incoming radiation condition, *Phys. Rev. D* **100**, 024042 (2019).
- [36] G. Compère, A. Fiorucci, and R. Ruzziconi, The Λ -BMS₄ group of dS₄ and new boundary conditions for AdS₄, *Classical Quantum Gravity* **36**, 195017 (2019); **38**, 229501(E) (2021).
- [37] R. Ruzziconi, Asymptotic symmetries in the gauge fixing approach and the BMS Group, *Proc. Sci., Modave2019* (2020) 003 [arXiv:1910.08367].
- [38] P. T. Chruściel, S. J. Hoque, and T. Smořka, Energy of weak gravitational waves in spacetimes with a positive cosmological constant, *Phys. Rev. D* **103**, 064008 (2021).
- [39] G. Compère, A. Fiorucci, and R. Ruzziconi, The Λ -BMS₄ charge algebra, *J. High Energy Phys.* **10** (2020) 205.
- [40] F. Fernández-Álvarez and J. M. M. Senovilla, Gravitational radiation condition at infinity with a positive cosmological constant, *Phys. Rev. D* **102**, 101502 (2020).
- [41] M. Kolanowski and J. Lewandowski, Energy of gravitational radiation in the de Sitter universe at \mathcal{I}^+ and at a horizon, *Phys. Rev. D* **102**, 124052 (2020).
- [42] B. Bonga and K. Prabhu, BMS-like symmetries in cosmology, *Phys. Rev. D* **102**, 104043 (2020).
- [43] J. Fier, X. Fang, B. Li, S. Mukohyama, A. Wang, and T. Zhu, Gravitational wave cosmology: High frequency approximation, *Phys. Rev. D* **103**, 123021 (2021).
- [44] P. T. Chruściel, S. J. Hoque, M. Maliborski, and T. Smořka, On the canonical energy of weak gravitational fields with a cosmological constant $\Lambda \in \mathbb{R}$, *Eur. Phys. J. C* **81**, 696 (2021).
- [45] M. Kolanowski and J. Lewandowski, Hamiltonian charges in the asymptotically de Sitter spacetimes, *J. High Energy Phys.* **05** (2021) 063.
- [46] J. R. Fier, Gravitational waves and cosmology. Ph.D. thesis, Baylor U., Baylor U., 2021.
- [47] S. Chakraborty, S. J. Hoque, and R. Oliveri, Gravitational multipole moments for asymptotically de Sitter spacetimes, *Phys. Rev. D* **104**, 064019 (2021).
- [48] A. Fiorucci, Leaky covariant phase spaces: Theory and application to Λ -BMS symmetry. Ph.D. thesis, Brussels U., Intl. Solvay Inst., Brussels, 2021.
- [49] A. Poole, K. Skenderis, and M. Taylor, Charges, conserved quantities, and fluxes in de Sitter spacetime, *Phys. Rev. D* **106**, L061901 (2022).
- [50] D. Dobkowski-Ryłko and J. Lewandowski, Generalization of the quadrupole formula for the energy of gravitational radiation in de Sitter spacetime, *Phys. Rev. D* **106**, 044047 (2022).
- [51] M. Geiller and C. Zwikel, The partial Bondi gauge: Further enlarging the asymptotic structure of gravity, *SciPost Phys.* **13**, 108 (2022).
- [52] W. Kamiński, M. Kolanowski, and J. Lewandowski, Symmetries of the asymptotically de Sitter spacetimes, *Classical Quantum Gravity* **39**, 195009 (2022).
- [53] A. M. Grant, A. Saffer, L. C. Stein, and S. Tahura, Gravitational-wave energy and other fluxes in ghost-free bigravity, *Phys. Rev. D* **107**, 044041 (2023).
- [54] J. M. M. Senovilla, Gravitational radiation at infinity with non-negative cosmological constant, *Universe* **8**, 478 (2022).
- [55] B. Bonga, C. Bunster, and A. Pérez, Gravitational radiation with $\Lambda > 0$, *Phys. Rev. D* **108**, 064039 (2023).
- [56] F. Fernández-Álvarez, Degrees of freedom of gravitational radiation with positive cosmological constant, *Phys. Rev. D* **108**, 064028 (2023).
- [57] G. Compère, S. J. Hoque, and E. c. Kutluk, Quadrupolar radiation in de Sitter: Displacement memory and Bondi metric, *Classical Quantum Gravity* **41**, 155006 (2024).

- [58] Harsh, S. J. Hoque, S. P. Kashyap, and A. Virmani, de Sitter Teukolsky waves, *Classical Quantum Gravity* **41**, 225011 (2024).
- [59] F. Fernández-Álvarez, J. Podolský, and J. M. M. Senovilla, Analysis of gravitational radiation generated by type D black holes with positive cosmological constant, *Phys. Rev. D* **110**, 104029 (2024).
- [60] D. Dobkowski-Ryłko and J. Lewandowski, Quadrupole formulae with cosmological constant: Comparison, *J. High Energy Phys.* **01** (2025) 034.
- [61] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, Gravitational waves in general relativity. VII. Waves from axi-symmetric isolated systems, *Proc. R. Soc. A* **269**, 21 (1962).
- [62] R. Sachs, Asymptotic symmetries in gravitational theory, *Phys. Rev.* **128**, 2851 (1962).
- [63] D. Kastor and J. H. Traschen, A positive energy theorem for asymptotically de Sitter space-times, *Classical Quantum Gravity* **19**, 5901 (2002).
- [64] M.-x. Luo, N.-q. Xie, and X. Zhang, Positive mass theorems for asymptotically de Sitter spacetimes, *Nucl. Phys.* **B825**, 98 (2010).
- [65] W. R. Kelly and D. Marolf, Phase spaces for asymptotically de Sitter cosmologies, *Classical Quantum Gravity* **29**, 205013 (2012).
- [66] X. Zhang, Recent progress on the positive energy theorem, *Int. J. Mod. Phys. A* **30**, 1545018 (2015).
- [67] A. A. Starobinsky, Isotropization of arbitrary cosmological expansion given an effective cosmological constant, *JETP Lett.* **37**, 66 (1983).
- [68] G. Compère, S. E. Gralla, and H. Wei, An asymptotic framework for gravitational scattering, *Classical Quantum Gravity* **40**, 205018 (2023).
- [69] G. Barnich and C. Troessaert, BMS charge algebra, *J. High Energy Phys.* **12** (2011) 105.
- [70] G. Compère, R. Oliveri, and A. Seraj, The Poincaré and BMS flux-balance laws with application to binary systems, *J. High Energy Phys.* **10** (2020) 116; **06** (2024) 45.
- [71] R. Epstein and R. V. Wagoner, Post-Newtonian generation of gravitational waves., *Astron. J.* **197**, 717 (1975).
- [72] A. Peres, Classical radiation recoil, *Phys. Rev.* **128**, 2471 (1962).
- [73] J. D. Bekenstein, Gravitational-radiation recoil and runaway black holes, *Astron. J.* **183**, 657 (1973).
- [74] K. S. Thorne, Multipole expansions of gravitational radiation, *Rev. Mod. Phys.* **52**, 299 (1980).
- [75] C. N. Kozameh, J. I. Nieva, and G. D. Quiroga, Spin and center of mass comparison between the post-Newtonian approach and the asymptotic formulation, *Phys. Rev. D* **98**, 064005 (2018).
- [76] S. J. Hoque and A. Aggarwal, Quadrupolar power radiation by a binary system in de Sitter background, *Int. J. Mod. Phys. D* **28**, 1950025 (2018).
- [77] V. Balasubramanian and P. Kraus, A stress tensor for Anti-de Sitter gravity, *Commun. Math. Phys.* **208**, 413 (1999).
- [78] M. Maggiore, *Gravitational Waves. Vol. 1: Theory and Experiments* (Oxford University Press, New York, 2007).