

Measurement of the Ω_c^0 and Ξ_c^0 baryon lifetimes using hadronic b -baryon decays



The LHCb collaboration

E-mail: Lars.Eklund@cern.ch

ABSTRACT: The lifetimes of the Ω_c^0 and Ξ_c^0 baryons are measured using a pp collision dataset collected by the LHCb experiment, corresponding to an integrated luminosity of 9 fb^{-1} . The charm baryons are produced in the fully reconstructed decay chains $\Omega_b^- \rightarrow \Omega_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$ and $\Xi_b^- \rightarrow \Xi_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$. The measurement uses topologically and kinematically similar $B^- \rightarrow D^0(\rightarrow K^-K^+\pi^-\pi^+)\pi^-$ decays for normalisation. The measured lifetimes are

$$\begin{aligned}\tau_{\Omega_c^0} &= 276.3 \pm 19.4 (\text{stat}) \pm 1.8 (\text{syst}) \pm 0.7 (\tau_{D^0}) \text{ fs}, \\ \tau_{\Xi_c^0} &= 149.2 \pm 2.5 (\text{stat}) \pm 0.9 (\text{syst}) \pm 0.4 (\tau_{D^0}) \text{ fs},\end{aligned}$$

where the first uncertainty is statistical, the second systematic and the third due to the uncertainty of the D^0 lifetime. These results are consistent with previous measurements performed by the LHCb experiment.

KEYWORDS: Charm Physics, Flavour Physics, Hadron-Hadron Scattering, Lifetime

ARXIV EPRINT: [2506.13334](https://arxiv.org/abs/2506.13334)

Contents

1	Introduction	1
2	Detector and simulation	2
3	Selection of signal candidates	3
4	Fits to the invariant-mass spectra	5
5	Lifetime measurements	8
6	Systematic uncertainties	10
7	Results and conclusion	12
	The LHCb collaboration	17

1 Introduction

Charm and beauty hadron lifetime measurements are important means to explore and document the properties of hadrons. Experimentally, accurate simulation of particle decays requires good knowledge of the lifetimes. From a theoretical viewpoint, charm hadron lifetimes are an excellent testing ground to evaluate different techniques for calculating hadron properties. Recent measurements of the Ω_c^0 and Ξ_c^0 lifetimes are in tension with earlier results, and this paper presents new results that confirm the revised lifetimes.

The lifetime of the Ω_c^0 baryon, with a valence quark content of css , was first measured by the E687 and WA89 collaborations [1, 2] and the FOCUS experiment [3], resulting in an average of 69 ± 12 fs [4]. In a more recent measurement, the LHCb experiment measured the Ω_c^0 lifetime to be almost four times larger than, and inconsistent with, the previous average. This measurement was based on a significantly larger sample of Ω_c^0 baryons originating from semileptonic b -baryon decays [5]. This result was subsequently confirmed by a measurement from the LHCb experiment using an independent sample of Ω_c^0 baryons produced promptly in pp collisions [6], yielding an average for the LHCb measurements of 275 ± 12 fs. The Belle II experiment recently measured the Ω_c^0 lifetime to be 243 ± 49 fs [7], consistent with the longer lifetime reported by LHCb.

Similarly, the lifetime of the Ξ_c^0 baryon, with a valence quark content of csd , was first measured by the ACCMOR, WA687 and FOCUS collaborations [8–10], resulting in an average of 112_{-10}^{+13} fs [4]. The more recent measurements by the LHCb experiment, using Ξ_c^0 baryons produced in semileptonic decays [11] and promptly in pp collisions [6], average to a lifetime of 152.0 ± 2.0 fs, in tension with the previous average. These discrepancies in measured lifetimes for the Ω_c^0 and Ξ_c^0 baryons highlight the need for additional measurements.

The lifetimes of charm baryons can be calculated using heavy-quark-expansion theory [12–19], which is an effective theory based on an expansion in inverse powers of the mass of the

heaviest quark. At leading order, the lifetime only depends on the mass of the charm quark, resulting in the same lifetime for all four weakly decaying single-charm baryons. Differences in the lifetimes of the single-charm baryons are expected to arise from higher-order effects, such as weak W -exchange and Pauli interference, caused by the spectator quarks. Charm hadron lifetimes are more sensitive to these contributions than beauty hadrons since the size of the contributions increases as the mass of the heaviest quark decreases [20–25].

The higher-order effects on the lifetimes of single-charm baryons have been evaluated to give a hierarchy of $\tau_{\Xi_c^+} > \tau_{\Lambda_c^+} > \tau_{\Xi_c^0} > \tau_{\Omega_c^0}$ [22–26], consistent with the earlier measurements of the lifetimes. Different treatments of the higher-order effects predict the hierarchy $\tau_{\Xi_c^+} > \tau_{\Omega_c^0} > \tau_{\Lambda_c^+} > \tau_{\Xi_c^0}$ [22, 25, 27], which is consistent with the more recent measurements. Hence, experimental input is crucial to arbitrate between these different calculations.

The measurements presented in this paper exploit charm baryons produced in hadronic b -baryon decays recorded by the LHCb experiment, specifically $\Omega_b^- \rightarrow \Omega_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$ and $\Xi_b^- \rightarrow \Xi_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$ decays.¹ These fully reconstructed decays have well defined production and decay vertices and are statistically independent from previous measurements by the LHCb experiment. The lifetimes are measured using the topologically and kinematically similar $B^- \rightarrow D^0(\rightarrow K^-K^+\pi^-\pi^+)\pi^-$ decay for normalisation to minimise the uncertainties associated with the decay time acceptance. Two different methods for the background subtraction are used to cross-check the results, giving consistent values. In order to avoid experimenter’s bias, the results of the analysis were not examined until the full procedure had been finalised.

The data sample was collected by the LHCb experiment in the years 2011, 2012 and 2015 – 2018 at centre-of-mass energies of $\sqrt{s} = 7, 8$ and 13 TeV, respectively, corresponding to an integrated luminosity of 9 fb^{-1} .

2 Detector and simulation

The LHCb detector [28, 29] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [30], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 T m, and three stations of silicon-strip detectors and straw drift tubes [31, 32] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [33]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [34].

¹The inclusion of charge-conjugate processes is implied throughout.

The online event selection is performed by a trigger [35], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. At the hardware trigger stage, events are required to have a muon with high p_T or a hadron, photon or electron with high transverse energy in the calorimeters. The signal candidates selected for the measurement are not required to have passed a specific set of criteria at the hardware stage; all candidates accepted by the trigger are included. The software trigger requires a two-, three- or four-track secondary vertex with a significant displacement from any PV. At least one charged particle must have a transverse momentum $p_T > 1.6 \text{ GeV}/c$ and be inconsistent with originating from a PV. A multivariate algorithm [36, 37] is used for the identification of secondary vertices consistent with the decay of a beauty hadron.

Simulation is required to determine how the detector acceptance and selection criteria affect the decay time distributions, and to determine the decay time resolution. In addition, simulation is used to develop the candidate selections. In the simulation, pp collisions are generated using PYTHIA [38, 39] with a specific LHCb configuration [40]. Decays of unstable particles are described by EVTGEN [41], in which final-state radiation is generated using PHOTOS [42]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [43, 44] as described in ref. [45]. To make more efficient use of computing resources, the underlying pp interaction is reused multiple times and combined with independently generated signal decays each time [46]. The simulated data are produced with charm and beauty hadron lifetimes consistent with the known values [47], or weighted to match those values if different.

The simulation provides a good description of the data, with the agreement further improved by data-driven corrections. The b -baryon p_T spectrum, track multiplicity and hardware-trigger efficiency are corrected with weights determined from the observed differences between data and simulation. The tracking efficiency corrections are applied using tables binned in pseudorapidity and p_T , derived from calibration samples. The simulation is corrected so that the fractions of candidates triggered by the signal or by other particles in the event match those in data. The particle identification performance in simulation is corrected using information from calibration data samples [48, 49].

The lifetime is measured from the distribution of decay times t for the signal candidates, evaluated in the particles' rest frames. It is determined from the reconstructed decay length l , invariant mass m and momentum p of the signal candidate, where $t = \frac{m \cdot l}{p}$ [30]. This relation is evaluated in a fit where the candidate is constrained to originate from its associated PV. The associated PV is that to which the candidates has the smallest χ_{IP}^2 , defined as the difference in the vertex-fit χ^2 of a given PV reconstructed with and without the track under consideration.

3 Selection of signal candidates

The selection of signal candidates exploits the long lifetime of the b -hadrons, avoiding requirements on quantities that directly affect the decay time distribution of the charm hadrons. Hence, the selection efficiencies only depend weakly on the charm hadron decay time, which simplifies the corrections for the decay time acceptance. The selection of the normalisation mode closely matches that of the signal modes, to ensure good cancellation

of the acceptance functions in the ratio of the signal and normalisation modes. Differences are limited to particle identification requirements and mass-range selections.

Four charged tracks, not originating from a PV and with requirements on their p and p_T , are selected to form a good-quality displaced vertex. The tracks must be consistent with the pion, kaon or proton hypotheses, as required to form the $\Omega_c^0 \rightarrow pK^-K^-\pi^+$, $\Xi_c^0 \rightarrow pK^-K^-\pi^+$ and $D^0 \rightarrow K^-K^+\pi^-\pi^+$ candidate decays.

Each charm hadron candidate is combined with a charged track that does not originate from a PV, has a high probability of being a pion and satisfies requirements on its p and p_T , to form a beauty-hadron candidate. The charm hadron and the additional track are required to originate from a common vertex, which must be displaced from all PVs and be of good quality. In addition, the reconstructed momentum vector of the b -hadron candidate must point back to the associated PV. A kinematic fit to the full decay tree [50] is performed on these $\Omega_b^- \rightarrow \Omega_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$, $\Xi_b^- \rightarrow \Xi_c^0(\rightarrow pK^-K^-\pi^+)\pi^-$ and $B^- \rightarrow D^0(\rightarrow K^-K^+\pi^-\pi^+)\pi^-$ candidates, in which they are constrained to originate from their associated PV. A loose requirement on the quality of that fit is applied, estimated by its χ^2 per degree of freedom ($\chi_{\text{DTF}}^2/\text{ndf}$).

A boosted decision tree (BDT) classifier [51, 52], implemented in the TMVA toolkit [53], is employed to further reject background. The classifier is trained using proxy datasets to represent signal and background decays. The signal is represented by simulated signal decays. The background is represented by candidates from data with $pK^-K^-\pi^+$ ($K^+K^-\pi^+\pi^-$) invariant masses outside the charm baryon (D^0 meson) signal regions, or with $\Omega_c^0\pi^-$, $\Xi_c^0\pi^-$ or $D^0\pi^-$ invariant masses significantly displaced above the b -hadron signal peaks. These training datasets are subject to the selection criteria described above.

Fifteen variables are used in the BDT to differentiate between signal and background. Among these, the variables associated with the pions from the b -hadron decays are their p_T , their χ_{IP}^2 and their probability of being a pion. Variables associated with the charm hadrons are their total momentum, the sum of p_T of their decay products and the probability of each decay product being assigned the correct particle identity. Variables associated with the b -hadron candidates are their p_T , χ_{IP}^2 and $\chi_{\text{DTF}}^2/\text{ndf}$, the χ^2/ndf of their decay vertex fits, the angle between their reconstructed flight path and momentum vector, and the flight distance significance, defined as the distance divided by its uncertainty.

The requirements on the BDT output variable for accepting charm baryon candidates are optimised by maximising the signal significance, which is defined as $S/\sqrt{S+B} = S_0\epsilon_s/\sqrt{S_0\epsilon_s+B}$. The background yield in the signal region, B , is determined from a fit to data. The signal yield S_0 is determined from a fit to data when applying a loose BDT requirement, which is then scaled with the change in selection efficiency (ϵ_s) determined from simulation, as the BDT requirement is tightened. This ensures that the measurement is not biased by the optimisation procedure. For the normalisation channel, a requirement that is tighter than the optimum given by the described procedure is applied. The number of selected candidates in the normalisation mode is significantly higher than for the signal modes; hence, the tighter selection does not affect the statistical precision, and the very clean selected sample simplifies the modelling of the invariant mass spectrum.

The training of the BDT classifier and the optimisation of the output variable requirement are performed separately for the data-taking periods 2011–2012 and 2015–2018, to account for the differences in running conditions and centre-of-mass energies.

Candidates are required to have $pK^-K^-\pi^+$ ($K^+K^-\pi^+\pi^-$) invariant masses within three times the mass resolution around the known charm baryon (D^0) masses [47], and $\Omega_c^0\pi^-$, $\Xi_c^0\pi^-$ and $D^0\pi^-$ invariant masses in ranges from 300, 200 and 150 MeV/ c^2 below to 550, 550 and 450 MeV/ c^2 above the known Ω_b^- , Ξ_b^- and B^- masses, respectively.

The selected decay time ranges for the two charm baryons are chosen to extend up to six times their lifetimes, which gives ranges of -0.15 to 1.50 ps for the Ω_c^0 candidates and -0.15 to 0.90 ps for the Ξ_c^0 candidates. Some candidates are reconstructed with negative decay times due to the finite decay time resolution; hence, the measurements include the negative decay time region. The normalisation mode candidates are analysed separately in two subsamples based on their lifetime, that match the ranges of the Ω_c^0 and Ξ_c^0 lifetime measurements.

After the full selection sequence is applied, approximately one percent of the events contain more than one signal candidate. These multiple candidates are retained for the measurement [54], to avoid the risk of biasing the measurement in the choice of which candidates to keep.

4 Fits to the invariant-mass spectra

The signal and background components are separated through unbinned maximum-likelihood fits to the invariant-mass spectra of the selected candidates. The mass fits are performed to the $\Omega_b^- \rightarrow \Omega_c^0\pi^-$, $\Xi_b^- \rightarrow \Xi_c^0\pi^-$ and $B^- \rightarrow D^0\pi^-$ invariant-mass spectra, since this gives a better discrimination between the signal and backgrounds with charm hadrons in the decay chain than fits to the $pK^-K^-\pi^+$ and $K^+K^-\pi^+\pi^-$ spectra. The invariant-mass spectra and fits are shown in figure 1 for the signal modes and in figure 2 for the normalisation mode.

The b -hadron signals are modelled with a modified Crystal Ball function [55] with power law tails on both sides of the peak (double-sided Crystal Ball, DCB), plus an additional Gaussian function. The tail parameters, as well as the relative widths and fractions of the two functions, are determined from simulation, whereas the peak position and overall width are determined from data.

The three spectra contain small fractions, less than 1%, of the Cabibbo suppressed $\Omega_b^- \rightarrow \Omega_c^0 K^-$, $\Xi_b^- \rightarrow \Xi_c^0 K^-$ and $B^- \rightarrow D^0 K^-$ decays where the kaons have been misidentified as pions. These components are modelled with DCB functions where all parameters are determined from simulation and their yields are constrained relative to the yields of the signal decays. The branching fractions of the b -baryon decays are not known; hence, the ratios of branching fractions between the Cabibbo favoured and suppressed decays are assumed to be the nominal Cabibbo suppression of approximately twenty, with an uncertainty of 30% to account for SU(3) breaking effects. The branching fractions of the $B^- \rightarrow D^0\pi^-$ and $B^- \rightarrow D^0 K^-$ decays are known with good accuracy and their ratio is calculated to be $(7.9 \pm 0.4)\%$ [47]. The relative efficiencies of the Cabibbo favoured and suppressed decays are determined from simulation. These ratios, including uncertainties, are combined to form Gaussian constraints in the fit.




Figure 1. Top row: invariant-mass distributions of reconstructed (left) $\Omega_b^- \rightarrow \Omega_c^0 \pi^-$ and (right) $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ candidates, with $pK^-K^-\pi^+$ invariant masses falling in the shaded regions of the bottom row plots. Bottom row: distributions of the reconstructed $pK^-K^-\pi^+$ invariant mass, for (left) Ω_c^0 and (right) Ξ_c^0 candidates with invariant masses within three times the mass resolution of the Ω_b^- and Ξ_b^- baryons, respectively.

Partially reconstructed decays, where one of the decay products is not included in the decay hypothesis, may enter the invariant-mass spectra. The decays $\Omega_b^- \rightarrow \Omega_c(2770)^0(\rightarrow \Omega_c^0 \gamma) \pi^-$, where the photons are not included, are modelled with a template determined from an analytic calculation. The shape depends on the helicity of the excited charm baryon, which is either $\frac{1}{2}$ or $\frac{3}{2}$, whose relative couplings are not known. Hence, both states are included as independent components of the fit, where their yields are determined to be approximately equal.




Figure 2. Top row: invariant-mass distributions of reconstructed $B^- \rightarrow D^0 \pi^-$ candidates, with $K^+ K^- \pi^+ \pi^-$ invariant masses falling in the shaded regions of the bottom-row plots. Bottom row: distributions of the reconstructed $K^+ K^- \pi^+ \pi^-$ invariant mass for B^- candidates with invariant masses within three times the mass resolution of the B^- meson. Candidates with D^0 decay times in the ranges used in the (left column) Ω_c^0 and (right column) Ξ_c^0 lifetime measurements are included.

The corresponding partially reconstructed $\Xi_b^- \rightarrow \Xi_c^{*0} (\rightarrow \Xi_c^0 \gamma) \pi^-$ decays are modelled with a sum of a generalised Gaussian function and a Johnsons’s S_U distribution [56], where the parameters are determined from fast simulation [57]. The dependence on the helicity of these decays is neglected since the yield is modest.

The $\Omega_c^0 \pi^-$ and $\Xi_c^0 \pi^-$ invariant-mass spectra contain partially reconstructed $\Omega_b^- \rightarrow \Omega_c^0 \pi^- \pi^0$ and $\Xi_b^- \rightarrow \Xi_c^0 \pi^- \pi^0$ decays, where the $\pi^- \pi^0$ either originate from $\rho(770)^-$ decays or are nonresonant. These decays are modelled with an Johnsons’s S_U distribution, with parameters fixed from fast simulation. The shapes of the invariant-mass distributions are not significantly affected by the helicity of the $\pi^- \pi^0$ system [58]. It is not possible to

determine the relative fractions of the resonant and nonresonant decays from data; hence, the model giving smallest fit χ^2 is selected. The nonresonant model is used for the fit to the $\Omega_c^0\pi^-$ invariant-mass spectrum and the resonant model is used for the $\Xi_c^0\pi^-$ invariant-mass spectrum. The yields of these decays are determined from data.

Partially reconstructed $B^- \rightarrow D^{*0}(\rightarrow D^0\gamma)\pi^-$ decays enter the $B^- \rightarrow D^0\pi^-$ invariant-mass spectrum and are modelled with a Johnsons's S_U distribution with parameters determined from fast simulation [57]. The ratio of yields relative to the $B^- \rightarrow D^0\pi^-$ decays, which is estimated from the relative branching fractions [47] and the relative reconstruction and selection efficiencies, is used as a Gaussian constraint in the fit.

Other peaking backgrounds are found to be negligible, for instance those from charmless Ξ_b^- and Ω_b^- baryon decays.

The combinatorial background in these spectra originate from two sources: either from correctly reconstructed Ω_c^0 , Ξ_c^0 or D^0 hadrons combined with an unassociated charged pion or from candidates built from five charged hadrons that originate from different processes in the event. The combinatorial background in the $\Omega_c^0\pi^-$ and $\Xi_c^0\pi^-$ invariant-mass spectra are dominated by the second category and the shapes are modelled with exponential functions, where the slope parameters are determined from a sample of candidates where the $pK^-K^-\pi^+$ invariant mass is not consistent with originating from Ω_c^0 or Ξ_c^0 decays. These parameters, including uncertainties, are used as Gaussian constraints in the fit to data.

The combinatorial background in the $B^- \rightarrow D^0\pi^-$ spectrum is very small and modelled by an exponential function with an unconstrained slope.

The fits give signal yields of 355 ± 26 for the $\Omega_b^- \rightarrow \Omega_c^0\pi^-$ decays and $8\,260 \pm 100$ for the $\Xi_b^- \rightarrow \Xi_c^0\pi^-$ decays. The fitted yields for the $B^- \rightarrow D^0\pi^-$ decays are $53\,420 \pm 260$ ($48\,720 \pm 250$) in the decay time range used for the Ω_c^0 (Ξ_c^0) lifetime measurement.

5 Lifetime measurements

The Ω_c^0 and Ξ_c^0 lifetimes are measured with a least-squares fit to the ratio of yields between the signal and normalisation modes, in bins of decay time. The bin boundaries are chosen to give approximately the same number of signal candidates in each bin, where six bins are used for the Ω_c^0 lifetime measurement and fifteen are used for the Ξ_c^0 lifetime measurement. The ratio of yields in decay time bin i with boundaries t_i and t_{i+1} is modelled to be

$$r_i \equiv \frac{\int_{t_i}^{t_{i+1}} A_{\Omega_c^0, \Xi_c^0}(t) \cdot \left(e^{-t/\tau_{\Omega_c^0, \Xi_c^0}} * R_{\Omega_c^0, \Xi_c^0}(t) \right) dt}{\int_{t_i}^{t_{i+1}} A_{D^0}(t) \cdot \left(e^{-t/\tau_{D^0}} * R_{D^0}(t) \right) dt}, \quad (5.1)$$

where $\tau_{\Omega_c^0, \Xi_c^0, D^0}$ are the hadron lifetimes, $A_{\Omega_c^0, \Xi_c^0, D^0}$ the decay time acceptance functions, $R_{\Omega_c^0, \Xi_c^0, D^0}$ the decay time resolution functions and the symbol $*$ represents a convolution. The D^0 lifetime is fixed to its known value of $\tau_{D^0} = 410.3 \pm 1.0$ fs [47], while the charm baryon lifetimes are determined from the fit.

The decay time resolution function is determined from simulation and modelled with a sum of three Gaussian functions. The effective resolutions, defined as the square root of the sum of the Gaussian variances weighted by their fractions, are 83 fs for the Ω_c^0 , 101 fs for the

Ξ_c^0 and 75 fs for the D^0 decay times. A multiplicative scale factor to account for differences in resolution between data and simulation is determined from a fit to the D^0 decay time distribution. The distribution is modelled with an exponential function, with the D^0 lifetime fixed to its known value, convolved with the resolution function including a scale factor for its width, and multiplied by the acceptance function. The scale factor is determined to be 1.08 ± 0.02 , which is applied to all three decay modes.

The decay time acceptance functions are determined from simulation, from the ratio between the reconstructed decay time distributions and a model consisting of an exponential function convolved with the resolution function. The lifetimes in the exponential functions are set to the values used in the simulation. The acceptance is almost independent of the charm hadron decay time, which is expected since the selection is designed to exploit the long b -hadron lifetimes. They are modelled with the sum of second-order polynomials and a Gaussian function with a mean of approximately -0.15 ps to account for a small deviation from a polynomial shape in that region.

Two different methods are employed to determine the decay time distribution of the signal. In the first method, the data are divided into bins of decay time and a mass fit is performed on each subset. The same fit functions as those described in section 4 are used, with only the yields allowed to vary and all other parameters fixed to the values given by the time-integrated mass fits. The ratios of signal and normalisation mode yields in each bin are fitted with the function given in eq. 5.1.

The second method uses the *sPlot* technique [59], a statistical tool that determines the decay time distributions of the components included in the mass fits, using the $\Omega_b^- \rightarrow \Omega_c^0 \pi^-$, $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ and $B^- \rightarrow D^0 \pi^-$ invariant masses as the discriminating variables. This results in binned decay time distributions for the signal and normalisation modes. The ratios of the distributions for the signal and normalisation modes are fit with the function given in eq. 5.1.

Both methods return the generated lifetimes when applied to signal simulation, and give unbiased results when evaluated on a large number of pseudoexperiments that include backgrounds. The statistical correlations of the measurements with the two methods, determined from pseudoexperiments, are 92% for the Ω_c^0 lifetime and 99% for the Ξ_c^0 lifetime. The statistical and systematic uncertainties of the two methods are nearly identical. The results of the two methods were compared before unblinding and were found to be consistent when accounting for their total uncertainties and correlations.

The first method for background subtraction, fitting the signal yields in bins of decay time, is used for the nominal result of this measurement. This method was selected since a similar method was used in a previous LHCb measurement [6] and since pseudoexperiments indicate that it is slightly more precise on average. This choice was made before the results were unblinded.

The fits to the lifetime ratios are shown in figure 3 and the measured lifetimes are

$$\begin{aligned}\tau_{\Omega_c^0} &= 276.3 \pm 19.4 \text{ (stat) fs,} \\ \tau_{\Xi_c^0} &= 149.2 \pm 2.5 \text{ (stat) fs.}\end{aligned}$$

For comparison, the measured lifetimes for the second method are $\tau_{\Omega_c^0} = 265.5 \pm 19.8$ (stat) and $\tau_{\Xi_c^0} = 148.2 \pm 2.6$ (stat) fs.




Figure 3. Ratio of yields (r_i) between the (left) $\Omega_b^- \rightarrow \Omega_c^0 \pi^-$ and (right) $\Xi_b^- \rightarrow \Xi_c^0 \pi^-$ decays, and the $B^- \rightarrow D^0 \pi^-$ decays, in bins of decay time. The results from the fit with the model given in eq. 5.1 are shown.

Sources	$\sigma_{\tau_{\Omega_c^0}}$ [fs]	$\sigma_{\tau_{\Xi_c^0}}$ [fs]
Signal mass modelling	0.1	< 0.1
Partially reconstructed background modelling	< 0.1	< 0.1
Cabibbo suppressed $\Xi_b^- \rightarrow \Omega_c^0 \pi^-$ decay	< 0.1	—
Helicity of the $\Omega_b^- \rightarrow \Omega_c(2770)^0 (\rightarrow \Omega_c^0 \gamma) \pi^-$ decay	0.3	—
Combinatorial background modelling	0.3	< 0.1
Decay time acceptance modelling	1.7	0.8
Decay time resolution modelling	0.1	0.5
Decay time resolution correction	< 0.1	< 0.1
Uncertainty of the Ω_b^- and Ξ_b^- lifetimes	0.5	0.1
Total systematic uncertainty	1.8	0.9
Uncertainty of the D^0 lifetime	0.7	0.4

Table 1. Systematic uncertainties for the Ω_c^0 and Ξ_c^0 lifetime measurements.

6 Systematic uncertainties

Several sources of systematic uncertainty are considered and they are summarised in table 1. In addition, the quoted statistical uncertainty includes a small component of systematic uncertainty due to the Gaussian constraints used in the mass fits.

The systematic uncertainties associated with the shape of the signal mass models are evaluated by generating pseudoexperiments with the nominal model described in section 4, and fitting them with an alternative model consisting of a DCB function without the additional

Gaussian function. The difference in average lifetime between the nominal and alternative fits, across all pseudoexperiments, is taken as the systematic uncertainty.

The uncertainty related to the modelling of the partially reconstructed background is evaluated by generating pseudoexperiments using the nominal values for the parameters governing the shapes, and fitting them with parameter values that are varied within their uncertainties. The difference in fitted lifetime with the two sets of parameter values is determined for each pseudoexperiment, and the spread of this difference plus its average value is assigned as the systematic uncertainty. The decay $\Xi_b^- \rightarrow \Xi_c(2645)^0 (\rightarrow \Xi_c^0 \pi^0) \pi^-$ could contribute to the selected data, and an associated uncertainty is evaluated from pseudoexperiments that include this background, which are fit with the nominal model. This uncertainty is found to be negligible, since the background is absorbed in the modelling of the other partially reconstructed backgrounds.

Similarly, background from the Cabibbo suppressed decay $\Xi_b^- \rightarrow \Omega_c^0 \pi^-$ could be present in the $\Omega_b^- \rightarrow \Omega_c^0 \pi^-$ mass spectrum, but its influence on the measurement is found to be negligible.

The uncertainty associated with the unknown fractions of the two helicity states for the $\Omega_b^- \rightarrow \Omega_c(2770)^0 (\rightarrow \Omega_c^0 \gamma) \pi^-$ decay is evaluated by generating pseudoexperiments with the helicity fractions taken from the nominal fit result. The pseudoexperiments are fit with two alternative models, either with helicity $\frac{1}{2}$ or $\frac{3}{2}$. The average of the difference between the nominal and alternative models is computed, and the largest of the two differences is taken as the systematic uncertainty.

The uncertainty related to the combinatorial background modelling is partially included in the statistical uncertainty, through the Gaussian constraint on the slope parameter. An additional uncertainty is assigned by fitting the pseudoexperiments with the slope parameter allowed to vary freely in the fit. The results from these fits are compared with those from the nominal fit, and the average difference is assigned as the systematic uncertainty.

The systematic uncertainties associated with the decay time acceptance and decay time resolution modelling are evaluated by generating pseudoexperiments with the nominal models described in section 5 and fitting them with alternative models. For the acceptance functions, a first order polynomial is used as the alternative model, without the additional Gaussian function. This is the dominant systematic uncertainty for both measurements.

The uncertainty related to the decay time resolution modelling is evaluated with an alternative model consisting of a sum of two Gaussian functions. The average differences between the nominal and alternative models are taken as the systematic uncertainties.

The uncertainty on the decay time resolution scale factor extracted from data is propagated to the lifetime measurements. Pseudoexperiments are generated using the nominal value of the scale factor and fitted with the value varied within its uncertainty, independently for the signal and normalisation modes. The spread of this difference plus its average value is assigned as the systematic uncertainty.

The simulated samples used to determine the acceptance functions are generated with the known lifetimes for the Ω_b^- and Ξ_b^- baryons [47]. Since these lifetimes are known with finite precision, the associated systematic uncertainty is evaluated by weighting the simulated samples to vary the b -baryon lifetimes up and down by one unit of their uncertainty. The acceptance functions determined from these weighted samples are used as alternative models

to generate pseudoexperiments, which are fit with the nominal model. The average shift in fitted lifetime is assigned as the systematic uncertainty.

The charm baryon lifetimes are measured relative to the D^0 lifetime; hence, the fractional uncertainty on its lifetime is assigned as a separate systematic uncertainty.

The consistency of the results is validated by performing the measurement on subsets of the data, divided into years of data taking, polarity of the LHCb magnet, centre-of-mass energy (7 and 8 TeV versus 13 TeV) and divided into two regions of $m(K^-\pi^+)$ invariant mass. The latter is to verify that the resonant structures present in the charm hadron decays do not affect the measured lifetimes. The data are also divided into two trigger categories, depending on if the hardware trigger decision was based on properties of the signal candidate or based on other particles produced in the pp collisions. The results from all these subsets are consistent for both methods of background subtraction.

7 Results and conclusion

The lifetimes of the Ω_c^0 and Ξ_c^0 baryons are measured in fully reconstructed decay chains using data collected by the LHCb experiment in the years 2011–2012 and 2015–2018 corresponding to an integrated luminosity of 9 fb^{-1} . The measured lifetimes are

$$\begin{aligned}\tau_{\Omega_c^0} &= 276.3 \pm 19.4 (\text{stat}) \pm 1.8 (\text{syst}) \pm 0.7 (\tau_{D^0}) \text{ fs}, \\ \tau_{\Xi_c^0} &= 149.2 \pm 2.5 (\text{stat}) \pm 0.9 (\text{syst}) \pm 0.4 (\tau_{D^0}) \text{ fs},\end{aligned}$$

where the first uncertainty is statistical, the second is systematic and the third is due to the uncertainty of the D^0 lifetime. The statistical uncertainties are dominant, in particular for the Ω_c^0 lifetime. These results are consistent with the previous measurements by the LHCb experiment [5, 6, 11], confirming the revised hierarchy of charm baryon lifetimes, and provide guidance on the accuracy of the different techniques used to calculate charm baryon lifetimes.

The measured lifetimes are combined with the previous LHCb measurements [5, 6, 11], using a linear unbiased estimator [60, 61], accounting for correlations. The measurements are statistically independent and all non-negligible systematic uncertainties are uncorrelated, apart from those related to the decay time resolution, which conservatively are considered to be fully correlated.

The studies of semileptonic decays [5, 11] measured the ratio between the charm baryons and the D^+ lifetimes, with the known value of the D^+ lifetime [4] as input. The knowledge of the D^+ lifetime has since improved [62], resulting in a new average [47] which is used to re-evaluate the Ω_c^0 and Ξ_c^0 lifetimes. The LHCb combinations use these updated lifetimes. The uncertainties associated with the D^+ and D^0 lifetime uncertainties are uncorrelated.

The measurements that used promptly produced charm baryons [6] were performed relative to a D^0 decay, using a previous average for the D^0 lifetime [63] as input. The measurements with hadronic decays use an updated average D^0 lifetime [47], which includes a new measurement [62]. The correlation between the systematic uncertainties associated with the D^0 lifetime between the two measurements is estimated to be 68%.

The combinations give χ^2/ndf values of 0.13/2 for the Ω_c^0 lifetime and 2.1/2 for the Ξ_c^0 lifetime and the resulting averages of the LHCb measurements are

$$\begin{aligned}\tau_{\Omega_c^0} &= 274.8 \pm 10.5 \text{ fs}, \\ \tau_{\Xi_c^0} &= 150.7 \pm 1.6 \text{ fs},\end{aligned}$$

where the uncertainties include both statistical and systematic effects.

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: ARC (Australia); CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICIU and AEI (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (U.S.A.). We acknowledge the computing resources that are provided by ARDC (Australia), CBPF (Brazil), CERN, IHEP and LZU (China), IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), Polish WLCG (Poland), IFIN-HH (Romania), PIC (Spain), CSCS (Switzerland), and GridPP (United Kingdom). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Alexander-von-Humboldt Foundation (Germany); ICSC (Italy); Severo Ochoa and María de Maeztu Units of Excellence, GVA, XuntaGal, GENCAT, InTalent-Inditex and Prog. Atracción Talento CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

Data Availability Statement. This article has no associated data or the data will not be deposited.

Code Availability Statement. This article has no associated code or the code will not be deposited.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E687 collaboration, *First measurement of the lifetime of the Ω_c^0* , *Phys. Lett. B* **357** (1995) 678 [[INSPIRE](#)].
- [2] WA89 collaboration, *Measurement of the Ω_c^0 lifetime*, *Phys. Lett. B* **358** (1995) 151 [[hep-ex/9507004](#)] [[INSPIRE](#)].

- [3] FOCUS collaboration, *Measurement of the Ω_c^0 lifetime*, *Phys. Lett. B* **561** (2003) 41 [[hep-ex/0302033](#)] [[INSPIRE](#)].
- [4] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **98** (2018) 030001 [[INSPIRE](#)].
- [5] LHCb collaboration, *Measurement of the Ω_c^0 baryon lifetime*, *Phys. Rev. Lett.* **121** (2018) 092003 [[arXiv:1807.02024](#)] [[INSPIRE](#)].
- [6] LHCb collaboration, *Measurement of the lifetimes of promptly produced Ω_c^0 and Ξ_c^0 baryons*, *Sci. Bull.* **67** (2022) 479 [[arXiv:2109.01334](#)] [[INSPIRE](#)].
- [7] BELLE-II collaboration, *Measurement of the Ω_c^0 lifetime at Belle II*, *Phys. Rev. D* **107** (2023) L031103 [[arXiv:2208.08573](#)] [[INSPIRE](#)].
- [8] ACCMOR collaboration, *First measurement of the lifetime of the charmed strange baryon Ξ_c^0* , *Phys. Lett. B* **236** (1990) 495 [[INSPIRE](#)].
- [9] E687 collaboration, *Measurement of the lifetime of the Ξ_c^0* , *Phys. Rev. Lett.* **70** (1993) 2058 [[INSPIRE](#)].
- [10] FOCUS collaboration, *A new measurement of the Ξ_c^0 lifetime*, *Phys. Lett. B* **541** (2002) 211 [[hep-ex/0206069](#)] [[INSPIRE](#)].
- [11] LHCb collaboration, *Precision measurement of the Λ_c^+ , Ξ_c^+ and Ξ_c^0 baryon lifetimes*, *Phys. Rev. D* **100** (2019) 032001 [[arXiv:1906.08350](#)] [[INSPIRE](#)].
- [12] V.A. Khoze and M.A. Shifman, *Heavy quarks*, *Sov. Phys. Usp.* **26** (1983) 387 [[INSPIRE](#)].
- [13] I.I.Y. Bigi and N.G. Uraltsev, *Gluonic enhancements in non-spectator beauty decays: an inclusive mirage though an exclusive possibility*, *Phys. Lett. B* **280** (1992) 271 [[INSPIRE](#)].
- [14] I.I.Y. Bigi, N.G. Uraltsev and A.I. Vainshtein, *Nonperturbative corrections to inclusive beauty and charm decays: QCD versus phenomenological models*, *Phys. Lett. B* **293** (1992) 430 [Erratum *ibid.* **297** (1992) 477] [[hep-ph/9207214](#)] [[INSPIRE](#)].
- [15] B. Blok and M.A. Shifman, *The rule of discarding $1/N_c$ in inclusive weak decays. 1*, *Nucl. Phys. B* **399** (1993) 441 [[hep-ph/9207236](#)] [[INSPIRE](#)].
- [16] B. Blok and M.A. Shifman, *The rule of discarding $1/N_c$ in inclusive weak decays. 2*, *Nucl. Phys. B* **399** (1993) 459 [[hep-ph/9209289](#)] [[INSPIRE](#)].
- [17] M. Neubert, *B decays and the heavy quark expansion*, *Adv. Ser. Direct. High Energy Phys.* **15** (1998) 239 [[hep-ph/9702375](#)] [[INSPIRE](#)].
- [18] N. Uraltsev, *Heavy quark expansion in beauty and its decays*, *Proc. Int. Sch. Phys. Fermi* **137** (1998) 329 [[hep-ph/9804275](#)] [[INSPIRE](#)].
- [19] I.I.Y. Bigi, *The QCD perspective on lifetimes of heavy flavor hadrons*, [hep-ph/9508408](#) [[INSPIRE](#)].
- [20] M. Kirk, A. Lenz and T. Rauh, *Dimension-six matrix elements for meson mixing and lifetimes from sum rules*, *JHEP* **12** (2017) 068 [Erratum *ibid.* **06** (2020) 162] [[arXiv:1711.02100](#)] [[INSPIRE](#)].
- [21] A. Lenz and T. Rauh, *D-meson lifetimes within the heavy quark expansion*, *Phys. Rev. D* **88** (2013) 034004 [[arXiv:1305.3588](#)] [[INSPIRE](#)].
- [22] H.-Y. Cheng, *Charmed baryon physics circa 2021*, *Chin. J. Phys.* **78** (2022) 324 [[arXiv:2109.01216](#)] [[INSPIRE](#)].
- [23] S. Bianco, F.L. Fabbri, D. Benson and I. Bigi, *A Cicerone for the physics of charm*, *Riv. Nuovo Cim.* **26** (2003) 1 [[hep-ex/0309021](#)] [[INSPIRE](#)].

- [24] G. Bellini, I.I.Y. Bigi and P.J. Dornan, *Lifetimes of charm and beauty hadrons*, *Phys. Rept.* **289** (1997) 1 [INSPIRE].
- [25] B. Blok and M.A. Shifman, *Lifetimes of charmed hadrons revisited. Facts and fancy*, in the proceedings of the *3rd workshop on the tau-charm factory*, (1991) [hep-ph/9311331] [INSPIRE].
- [26] H.-Y. Cheng, *A phenomenological analysis of heavy hadron lifetimes*, *Phys. Rev. D* **56** (1997) 2783 [hep-ph/9704260] [INSPIRE].
- [27] H.-Y. Cheng, *Phenomenological study of heavy hadron lifetimes*, *JHEP* **11** (2018) 014 [arXiv:1807.00916] [INSPIRE].
- [28] LHCb collaboration, *The LHCb detector at the LHC*, 2008 *JINST* **3** S08005 [INSPIRE].
- [29] LHCb collaboration, *LHCb detector performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [arXiv:1412.6352] [INSPIRE].
- [30] R. Aaij et al., *Performance of the LHCb vertex locator*, 2014 *JINST* **9** P09007 [arXiv:1405.7808] [INSPIRE].
- [31] LHCb OUTER TRACKER GROUP collaboration, *Performance of the LHCb outer tracker*, 2014 *JINST* **9** P01002 [arXiv:1311.3893] [INSPIRE].
- [32] LHCb OUTER TRACKER GROUP collaboration, *Improved performance of the LHCb outer tracker in LHC run 2*, 2017 *JINST* **12** P11016 [arXiv:1708.00819] [INSPIRE].
- [33] LHCb RICH GROUP collaboration, *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [arXiv:1211.6759] [INSPIRE].
- [34] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, 2013 *JINST* **8** P02022 [arXiv:1211.1346] [INSPIRE].
- [35] R. Aaij et al., *The LHCb trigger and its performance in 2011*, 2013 *JINST* **8** P04022 [arXiv:1211.3055] [INSPIRE].
- [36] V.V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, 2013 *JINST* **8** P02013 [arXiv:1210.6861] [INSPIRE].
- [37] T. Likhomanenko et al., *LHCb topological trigger reoptimization*, *J. Phys. Conf. Ser.* **664** (2015) 082025 [arXiv:1510.00572] [INSPIRE].
- [38] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [arXiv:0710.3820] [INSPIRE].
- [39] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [hep-ph/0603175] [INSPIRE].
- [40] LHCb collaboration, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [INSPIRE].
- [41] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [INSPIRE].
- [42] N. Davidson, T. Przydzinski and Z. Was, *PHOTOS interface in C++: technical and physics documentation*, *Comput. Phys. Commun.* **199** (2016) 86 [arXiv:1011.0937] [INSPIRE].
- [43] J. Allison et al., *GEANT4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [INSPIRE].
- [44] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [INSPIRE].
- [45] LHCb collaboration, *The LHCb simulation application, Gauss: design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [INSPIRE].

- [46] D. Müller, M. Clemencic, G. Corti and M. Gersabeck, *ReDecay: a novel approach to speed up the simulation at LHCb*, *Eur. Phys. J. C* **78** (2018) 1009 [[arXiv:1810.10362](#)] [[INSPIRE](#)].
- [47] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **110** (2024) 030001 [[INSPIRE](#)].
- [48] ATLAS collaboration, *Prospect for a search for direct stau production in events with at least two hadronic taus and missing transverse momentum at the High Luminosity LHC with the ATLAS Detector*, [ATL-PHYS-PUB-2016-021](#), CERN, Geneva, Switzerland (2016).
- [49] R. Aaij et al., *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in run 2*, *EPJ Tech. Instrum.* **6** (2019) 1 [[arXiv:1803.00824](#)] [[INSPIRE](#)].
- [50] W.D. Hulsbergen, *Decay chain fitting with a Kalman filter*, *Nucl. Instrum. Meth. A* **552** (2005) 566 [[physics/0503191](#)] [[INSPIRE](#)].
- [51] L. Breiman, J. Friedman, R.A. Olshen and C.J. Stone, *Classification and regression trees*, Chapman and Hall/CRC (2017) [[DOI:10.1201/9781315139470](#)] [[INSPIRE](#)].
- [52] Y. Freund and R.E. Schapire, *A decision-theoretic generalization of on-line learning and an application to boosting*, *J. Comput. Syst. Sci.* **55** (1997) 119 [[INSPIRE](#)].
- [53] H. Voss, A. Hocker, J. Stelzer and F. Tegenfeldt, *TMVA, the Toolkit for Multivariate Data Analysis with ROOT*, *PoS ACAT* (2007) 040 [[INSPIRE](#)].
- [54] P. Koppenburg, *Statistical biases in measurements with multiple candidates*, [arXiv:1703.01128](#) [[INSPIRE](#)].
- [55] T. Skwarnicki, *A study of the radiative CASCADE transitions between the Upsilon-Prime and Upsilon resonances*, Ph.D. thesis, INP, Cracow, Poland (1986) [[INSPIRE](#)].
- [56] N.L. Johnson, *Systems of frequency curves generated by methods of translation*, *Biometrika* **36** (1949) 149 [[INSPIRE](#)].
- [57] G.A. Cowan, D.C. Craik and M.D. Needham, *RapidSim: an application for the fast simulation of heavy-quark hadron decays*, *Comput. Phys. Commun.* **214** (2017) 239 [[arXiv:1612.07489](#)] [[INSPIRE](#)].
- [58] LHCb collaboration, *Observation of excited Ω_c^0 baryons in $\Omega_b^- \rightarrow \Xi_c^+ K^- \pi^-$ decays*, *Phys. Rev. D* **104** (2021) L091102 [[arXiv:2107.03419](#)] [[INSPIRE](#)].
- [59] M. Pivk and F.R. Le Diberder, *SPlot: a statistical tool to unfold data distributions*, *Nucl. Instrum. Meth. A* **555** (2005) 356 [[physics/0402083](#)] [[INSPIRE](#)].
- [60] R. Nisius, *BLUE: combining correlated estimates of physics observables within ROOT using the Best Linear Unbiased Estimate method*, *SoftwareX* **11** (2020) 100468 [[arXiv:2001.10310](#)] [[INSPIRE](#)].
- [61] R. Nisius, *On the combination of correlated estimates of a physics observable*, *Eur. Phys. J. C* **74** (2014) 3004 [[arXiv:1402.4016](#)] [[INSPIRE](#)].
- [62] BELLE-II collaboration, *Precise measurement of the D^0 and D^+ lifetimes at Belle II*, *Phys. Rev. Lett.* **127** (2021) 211801 [[arXiv:2108.03216](#)] [[INSPIRE](#)].
- [63] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2020** (2020) 083C01 [[INSPIRE](#)].

The LHCb collaboration

R. Aaij³⁸, A.S.W. Abdelmotteleb⁵⁷, C. Abellan Beteta⁵¹, F. Abudinén⁵⁷, T. Ackernley⁶¹, A. A. Adefisoye⁶⁹, B. Adeva⁴⁷, M. Adinolfi⁵⁵, P. Adlarson⁸⁴, C. Agapopoulou¹⁴, C.A. Aidala⁸⁶, Z. Ajaltouni¹¹, S. Akar¹¹, K. Akiba³⁸, P. Albicocco²⁸, J. Albrecht^{19,f}, F. Alessio⁴⁹, M. Alexander⁶⁰, Z. Aliouche⁶³, P. Alvarez Cartelle⁵⁶, R. Amalric¹⁶, S. Amato³, J.L. Amey⁵⁵, Y. Amhis¹⁴, L. An⁶, L. Anderlini²⁷, M. Andersson⁵¹, P. Andreola⁵¹, M. Andreotti²⁶, S. Andres Estrada⁸³, A. Anelli^{31,o,49}, D. Ao⁷, F. Archilli^{37,v}, Z. Areg⁶⁹, M. Argenton²⁶, S. Arguedas Cuendis^{9,49}, A. Artamonov⁴⁴, M. Artuso⁶⁹, E. Aslanides¹³, R. Ataíde Da Silva⁵⁰, M. Atzeni⁶⁵, B. Audurier¹², J. A. Authier¹⁵, D. Bacher⁶⁴, I. Bachiller Perea⁵⁰, S. Bachmann²², M. Bachmayer⁵⁰, J.J. Back⁵⁷, P. Baladron Rodriguez⁴⁷, V. Balagura¹⁵, A. Balboni²⁶, W. Baldini²⁶, L. Balzani¹⁹, H. Bao⁷, J. Baptista de Souza Leite⁶¹, C. Barbero Pretel^{47,12}, M. Barbetti²⁷, I. R. Barbosa⁷⁰, R.J. Barlow⁶³, M. Barnyakov²⁵, S. Barsuk¹⁴, W. Barter⁵⁹, J. Bartz⁶⁹, S. Bashir⁴⁰, B. Batsukh⁵, P. B. Battista¹⁴, A. Bay⁵⁰, A. Beck⁶⁵, M. Becker¹⁹, F. Bedeschi³⁵, I.B. Bediaga², N. A. Behling¹⁹, S. Belin⁴⁷, K. Belous⁴⁴, I. Belov²⁹, I. Belyaev³⁶, G. Benane¹³, G. Bencivenni²⁸, E. Ben-Haim¹⁶, A. Berezhnoy⁴⁴, R. Bernet⁵¹, S. Bernet Andres⁴⁶, A. Bertolin³³, C. Betancourt⁵¹, F. Betti⁵⁹, J. Bex⁵⁶, Ia. Bezshyiko⁵¹, O. Bezshyiko³⁵, J. Bhom⁴¹, M.S. Bieker¹⁸, N.V. Biesuz²⁶, P. Billoir¹⁶, A. Biolchini³⁸, M. Birch⁶², F.C.R. Bishop¹⁰, A. Bitadze⁶³, A. Bizzeti^{27,p}, T. Blake^{57,b}, F. Blanc⁵⁰, J.E. Blank¹⁹, S. Blusk⁶⁹, V. Bocharnikov⁴⁴, J.A. Boelhauve¹⁹, O. Boente Garcia¹⁵, T. Boettcher⁶⁸, A. Bohare⁵⁹, A. Boldyrev⁴⁴, C.S. Bolognani⁸¹, R. Bolzonella^{26,l}, R. B. Bonacci¹, N. Bondar^{44,49}, A. Bordelius⁴⁹, F. Borgato^{33,49}, S. Borghi⁶³, M. Borsato^{31,o}, J.T. Borsuk⁸², E. Bottalico⁶¹, S.A. Bouchiba⁵⁰, M. Bovill⁶⁴, T.J.V. Bowcock⁶¹, A. Boyer⁴⁹, C. Bozzi²⁶, J. D. Brandenburg⁸⁷, A. Brea Rodriguez⁵⁰, N. Breer¹⁹, J. Brodzicka⁴¹, A. Brossa Gonzalo^{47,†}, J. Brown⁶¹, D. Brundu³², E. Buchanan⁵⁹, L. Buonincontri^{33,q}, M. Burgos Marcos⁸¹, A.T. Burke⁶³, C. Burr⁴⁹, J.S. Butter⁵⁶, J. Buytaert⁴⁹, W. Byczynski⁴⁹, S. Cadeddu³², H. Cai⁷⁴, Y. Cai⁵, A. Caillet¹⁶, R. Calabrese^{26,l}, S. Calderon Ramirez⁹, L. Calefice⁴⁵, S. Cali²⁸, M. Calvi^{31,o}, M. Calvo Gomez⁴⁶, P. Camargo Magalhaes^{2,aa}, J. I. Cambon Bouzas⁴⁷, P. Campana²⁸, D.H. Campora Perez⁸¹, A.F. Campoverde Quezada⁷, S. Capelli³¹, L. Capriotti²⁶, R. Caravaca-Mora⁹, A. Carbone^{25,j}, L. Carcedo Salgado⁴⁷, R. Cardinale^{29,m}, A. Cardini³², P. Carniti³¹, L. Carus²², A. Casais Vidal⁶⁵, R. Caspary²², G. Casse⁶¹, M. Cattaneo⁴⁹, G. Cavallero²⁶, V. Cavallini^{26,l}, S. Celani²², S. Cesare^{30,n}, A.J. Chadwick⁶¹, I. Chahrouh⁸⁶, H. Chang^{4,c}, M. Charles¹⁶, Ph. Charpentier⁴⁹, E. Chatzianagnostou³⁸, R. Cheaib⁷⁸, M. Chefdeville¹⁰, C. Chen⁵⁶, J. Chen⁵⁰, S. Chen⁵, Z. Chen⁷, A. Chernov⁴¹, S. Chernyshenko⁵³, X. Chiotopoulos⁸¹, V. Chobanova⁸³, M. Chrzaszcz⁴¹, A. Chubykin⁴⁴, V. Chulikov^{28,36}, P. Ciambone²⁸, X. Cid Vidal⁴⁷, G. Ciezarek⁴⁹, P. Cifra³⁸, P.E.L. Clarke⁵⁹, M. Clemencic⁴⁹, H.V. Cliff⁵⁶, J. Closier⁴⁹, C. Cocha Toapaxi²², V. Coco⁴⁹, J. Cogan¹³, E. Cogneras¹¹, L. Cojocariu⁴³, S. Collaviti⁵⁰, P. Collins⁴⁹, T. Colombo⁴⁹, M. Colonna¹⁹, A. Comerma-Montells⁴⁵, L. Congedo²⁴, A. Contu³², N. Cooke⁶⁰, C. Coronel⁶⁶, I. Corredoira¹², A. Correia¹⁶, G. Corti⁴⁹, J. Cottee Meldrum⁵⁵, B. Couturier⁴⁹, D.C. Craik⁵¹, M. Cruz Torres^{2,g}, E. Curras Rivera⁵⁰, R. Currie⁵⁹, C.L. Da Silva⁶⁸, S. Dadabaev⁴⁴, L. Dai⁷¹, X. Dai⁴,

E. Dall’Occo ⁴⁹, J. Dalseno ⁸³, C. D’Ambrosio ⁶², J. Daniel ¹¹, P. d’Argent ²⁴, G. Darze ³, A. Davidson ⁵⁷, J.E. Davies ⁶³, O. De Aguiar Francisco ⁶³, C. De Angelis ^{32,k}, F. De Benedetti ⁴⁹, J. de Boer ³⁸, K. De Bruyn ⁸⁰, S. De Capua ⁶³, M. De Cian ⁶³,
 U. De Freitas Carneiro Da Graca ^{2,a}, E. De Lucia ²⁸, J.M. De Miranda ², L. De Paula ³,
 M. De Serio ^{24,h}, P. De Simone ²⁸, F. De Vellis ¹⁹, J.A. de Vries ⁸¹, F. Debernardis ²⁴,
 D. Decamp ¹⁰, S. Dekkers ¹, L. Del Buono ¹⁶, B. Delaney ⁶⁵, H.-P. Dembinski ¹⁹, J. Deng ⁸,
 V. Denysenko ⁵¹, O. Deschamps ¹¹, F. Dettori ^{32,k}, B. Dey ⁷⁸, P. Di Nezza ²⁸, I. Diachkov ⁴⁴,
 S. Didenko ⁴⁴, S. Ding ⁶⁹, Y. Ding ⁵⁰, L. Dittmann ²², V. Dobishuk ⁵³, A. D. Docheva ⁶⁰,
 C. Dong ^{4,c}, A.M. Donohoe ²³, F. Dordei ³², A.C. dos Reis ², A. D. Dowling ⁶⁹,
 W. Duan ⁷², P. Duda ⁸², M.W. Dudek ⁴¹, L. Dufour ⁴⁹, V. Duk ³⁴, P. Durante ⁴⁹, M.
 M. Duras ⁸², J.M. Durham ⁶⁸, O. D. Durmus ⁷⁸, A. Dziurda ⁴¹, A. Dzyuba ⁴⁴, S. Easo ⁵⁸,
 E. Eckstein ¹⁸, U. Egede ¹, A. Egorychev ⁴⁴, V. Egorychev ⁴⁴, S. Eisenhardt ⁵⁹, E. Ejopu ⁶³,
 L. Eklund ⁸⁴, M. Elashri ⁶⁶, J. Ellbracht ¹⁹, S. Ely ⁶², A. Ene ⁴³, J. Eschle ⁶⁹, S. Esen ²²,
 T. Evans ³⁸, F. Fabiano ³², S. Faghih ⁶⁶, L.N. Falcao ², B. Fang ⁷, R. Fantechi ³⁵,
 L. Fantini ^{34,r}, M. Faria ⁵⁰, K. Farmer ⁵⁹, D. Fazzini ^{31,o}, L. Felkowski ⁸², M. Feng ^{5,7},
 M. Feo ¹⁹, A. Fernandez Casani ⁴⁸, M. Fernandez Gomez ⁴⁷, A.D. Fernez ⁶⁷, F. Ferrari ^{25,j},
 F. Ferreira Rodrigues ³, M. Ferrillo ⁵¹, M. Ferro-Luzzi ⁴⁹, S. Filippov ⁴⁴, R.A. Fini ²⁴,
 M. Fiorini ^{26,l}, M. Firlej ⁴⁰, K.L. Fischer ⁶⁴, D.S. Fitzgerald ⁸⁶, C. Fitzpatrick ⁶³,
 T. Fiutowski ⁴⁰, F. Fleuret ¹⁵, A. Fomin ⁵², M. Fontana ²⁵, L. F. Foreman ⁶³, R. Forty ⁴⁹,
 D. Foulds-Holt ⁵⁹, V. Franco Lima ³, M. Franco Sevilla ⁶⁷, M. Frank ⁴⁹, E. Franzoso ^{26,l},
 G. Frau ⁶³, C. Frei ⁴⁹, D.A. Friday ⁶³, J. Fu ⁷, Q. Führung ^{19,f,56}, T. Fulghesu ¹³,
 G. Galati ²⁴, M.D. Galati ³⁸, A. Gallas Torreira ⁴⁷, D. Galli ^{25,j}, S. Gambetta ⁵⁹,
 M. Gandelman ³, P. Gandini ³⁰, B. Ganie ⁶³, H. Gao ⁷, R. Gao ⁶⁴, T.Q. Gao ⁵⁶, Y. Gao ⁸,
 Y. Gao ⁶, Y. Gao ⁸, L.M. Garcia Martin ⁵⁰, P. Garcia Moreno ⁴⁵, J. García Pardiñas ⁶⁵, P.
 Gardner ⁶⁷, K. G. Garg ⁸, L. Garrido ⁴⁵, C. Gaspar ⁴⁹, A. Gavrikov ³³, L.L. Gerken ¹⁹,
 E. Gersabeck ²⁰, M. Gersabeck ²⁰, T. Gershon ⁵⁷, S. Ghizzo ^{29,m}, Z. Ghorbanimoghaddam ⁵⁵,
 L. Giambastiani ^{33,q}, F. I. Giasemis ^{16,e}, V. Gibson ⁵⁶, H.K. Gienza ⁴², A.L. Gilman ⁶⁴,
 M. Giovannetti ²⁸, A. Gioventù ⁴⁵, L. Girardey ^{63,58}, M.A. Giza ⁴¹, F.C. Glaser ^{14,22},
 V.V. Gligorov ¹⁶, C. Göbel ⁷⁰, L. Golinka-Bezshyyko ⁸⁵, E. Golobardes ⁴⁶, D. Golubkov ⁴⁴,
 A. Golutvin ^{62,49}, S. Gomez Fernandez ⁴⁵, W. Gomulka ⁴⁰, I. Gonçalves Vaz ⁴⁹,
 F. Goncalves Abrantes ⁶⁴, M. Goncerz ⁴¹, G. Gong ^{4,c}, J. A. Gooding ¹⁹, I.V. Gorelov ⁴⁴,
 C. Gotti ³¹, E. Govorkova ⁶⁵, J.P. Grabowski ¹⁸, L.A. Granado Cardoso ⁴⁹, E. Graugés ⁴⁵,
 E. Graverini ^{50,t}, L. Grazette ⁵⁷, G. Graziani ²⁷, A. T. Grecu ⁴³, L.M. Greeven ³⁸,
 N.A. Grieser ⁶⁶, L. Grillo ⁶⁰, S. Gromov ⁴⁴, C. Gu ¹⁵, M. Guarise ²⁶, L. Guerry ¹¹,
 V. Guliaeva ⁴⁴, P. A. Günther ²², A.-K. Guseinov ⁵⁰, E. Gushchin ⁴⁴, Y. Guz ^{6,49}, T. Gys ⁴⁹,
 K. Habermann ¹⁸, T. Hadavizadeh ¹, C. Hadjivasiliou ⁶⁷, G. Haefeli ⁵⁰, C. Haen ⁴⁹, G.
 Hallett ⁵⁷, P.M. Hamilton ⁶⁷, J. Hammerich ⁶¹, Q. Han ³³, X. Han ^{22,49},
 S. Hansmann-Menzemer ²², L. Hao ⁷, N. Harnew ⁶⁴, T. H. Harris ¹, M. Hartmann ¹⁴,
 S. Hashmi ⁴⁰, J. He ^{7,d}, F. Hemmer ⁴⁹, C. Henderson ⁶⁶, R. Henderson ¹⁴,
 R.D.L. Henderson ¹, A.M. Hennequin ⁴⁹, K. Hennessy ⁶¹, L. Henry ⁵⁰, J. Herd ⁶²,
 P. Herrero Gascon ²², J. Heuel ¹⁷, A. Hicheur ³, G. Hijano Mendizabal ⁵¹, J. Horswill ⁶³,
 R. Hou ⁸, Y. Hou ¹¹, N. Howarth ⁶¹, J. Hu ⁷², W. Hu ⁷, X. Hu ^{4,c}, W. Hulsbergen ³⁸,
 R.J. Hunter ⁵⁷, M. Hushchyn ⁴⁴, D. Hutchcroft ⁶¹, M. Idzik ⁴⁰, D. Ilin ⁴⁴, P. Ilten ⁶⁶,

A. Iniukhin [ID](#)⁴⁴, A. Ishteev [ID](#)⁴⁴, K. Ivshin [ID](#)⁴⁴, H. Jage [ID](#)¹⁷, S.J. Jaimes Elles [ID](#)^{76,49,48},
 S. Jakobsen [ID](#)⁴⁹, E. Jans [ID](#)³⁸, B.K. Jashal [ID](#)⁴⁸, A. Jawahery [ID](#)⁶⁷, C. Jayaweera [ID](#)⁵⁴, V. Jevtic [ID](#)¹⁹,
 E. Jiang [ID](#)⁶⁷, X. Jiang [ID](#)^{5,7}, Y. Jiang [ID](#)⁷, Y. J. Jiang [ID](#)⁶, N. Jindal [ID](#)⁸⁷, M. John [ID](#)⁶⁴, A.
 John Rubesh Rajan [ID](#)²³, D. Johnson [ID](#)⁵⁴, C.R. Jones [ID](#)⁵⁶, T.P. Jones [ID](#)⁵⁷, S. Joshi [ID](#)⁴², B. Jost [ID](#)⁴⁹, J.
 Juan Castella [ID](#)⁵⁶, N. Jurik [ID](#)⁴⁹, I. Juszcak [ID](#)⁴¹, D. Kaminaris [ID](#)⁵⁰, S. Kandybei [ID](#)⁵², M. Kane [ID](#)⁵⁹,
 Y. Kang [ID](#)^{4,c}, C. Kar [ID](#)¹¹, M. Karacson [ID](#)⁴⁹, D. Karpenkov [ID](#)⁴⁴, A. Kauniskangas [ID](#)⁵⁰,
 J.W. Kautz [ID](#)⁶⁶, M.K. Kazanecki [ID](#)⁴¹, F. Keizer [ID](#)⁴⁹, M. Kenzie [ID](#)⁵⁶, T. Ketel [ID](#)³⁸, B. Khanji [ID](#)⁶⁹,
 A. Kharisova [ID](#)⁴⁴, S. Kholodenko [ID](#)^{35,49}, G. Khreich [ID](#)¹⁴, T. Kirn [ID](#)¹⁷, V.S. Kirsebom [ID](#)^{31,o},
 O. Kitouni [ID](#)⁶⁵, S. Klaver [ID](#)³⁹, N. Kleijne [ID](#)^{35,s}, K. Klimaszewski [ID](#)⁴², M.R. Kmiec [ID](#)⁴², S. Koliiev [ID](#)⁵³,
 L. Kolk [ID](#)¹⁹, A. Konoplyannikov [ID](#)⁶, P. Kopciewicz [ID](#)⁴⁹, P. Koppenburg [ID](#)³⁸, A. Korchin [ID](#)⁵²,
 M. Korolev [ID](#)⁴⁴, I. Kostiuk [ID](#)³⁸, O. Kot [ID](#)⁵³, S. Kotriakhova [ID](#), E. Kowalczyk [ID](#)⁶⁷, A. Kozachuk [ID](#)⁴⁴,
 P. Kravchenko [ID](#)⁴⁴, L. Kravchuk [ID](#)⁴⁴, M. Kreps [ID](#)⁵⁷, P. Krokovny [ID](#)⁴⁴, W. Krupa [ID](#)⁶⁹,
 W. Krzemien [ID](#)⁴², O. Kshyvanskyi [ID](#)⁵³, S. Kubis [ID](#)⁸², M. Kucharczyk [ID](#)⁴¹, V. Kudryavtsev [ID](#)⁴⁴,
 E. Kulikova [ID](#)⁴⁴, A. Kupsc [ID](#)⁸⁴, V. Kushnir [ID](#)⁵², B. Kutsenko [ID](#)¹³, I. Kyryllin [ID](#)⁵², D. Lacarrere [ID](#)⁴⁹,
 P. Laguarda Gonzalez [ID](#)⁴⁵, A. Lai [ID](#)³², A. Lampis [ID](#)³², D. Lancierini [ID](#)⁶², C. Landesa Gomez [ID](#)⁴⁷,
 J.J. Lane [ID](#)¹, G. Lanfranchi [ID](#)²⁸, C. Langenbruch [ID](#)²², J. Langer [ID](#)¹⁹, O. Lantwin [ID](#)⁴⁴, T. Latham [ID](#)⁵⁷,
 F. Lazzari [ID](#)^{35,t,49}, C. Lazzeroni [ID](#)⁵⁴, R. Le Gac [ID](#)¹³, H. Lee [ID](#)⁶¹, R. Lefèvre [ID](#)¹¹, A. Leflat [ID](#)⁴⁴,
 S. Legotin [ID](#)⁴⁴, M. Lehuraux [ID](#)⁵⁷, E. Lemos Cid [ID](#)⁴⁹, O. Leroy [ID](#)¹³, T. Lesiak [ID](#)⁴¹, E. D. Lesser [ID](#)⁴⁹,
 B. Leverington [ID](#)²², A. Li [ID](#)^{4,c}, C. Li [ID](#)⁴, C. Li [ID](#)¹³, H. Li [ID](#)⁷², J. Li [ID](#)⁸, K. Li [ID](#)⁷⁵, L. Li [ID](#)⁶³,
 M. Li [ID](#)⁸, P. Li [ID](#)⁷, P.-R. Li [ID](#)⁷³, Q. Li [ID](#)^{5,7}, S. Li [ID](#)⁸, T. Li [ID](#)⁷¹, T. Li [ID](#)⁷², Y. Li [ID](#)⁸, Y. Li [ID](#)⁵, Y.
 Li [ID](#)⁴, Z. Lian [ID](#)^{4,c}, Q. Liang [ID](#)⁸, X. Liang [ID](#)⁶⁹, S. Libralon [ID](#)⁴⁸, C. Lin [ID](#)⁷, T. Lin [ID](#)⁵⁸, R. Lindner [ID](#)⁴⁹,
 H. Linton [ID](#)⁶², R. Litvinov [ID](#)³², D. Liu [ID](#)⁸, F. L. Liu [ID](#)¹, G. Liu [ID](#)⁷², K. Liu [ID](#)⁷³, S. Liu [ID](#)^{5,7}, W.
 Liu [ID](#)⁸, Y. Liu [ID](#)⁵⁹, Y. Liu [ID](#)⁷³, Y. L. Liu [ID](#)⁶², G. Loachamin Ordonez [ID](#)⁷⁰, A. Lobo Salvia [ID](#)⁴⁵,
 A. Loi [ID](#)³², T. Long [ID](#)⁵⁶, J.H. Lopes [ID](#)³, A. Lopez Huertas [ID](#)⁴⁵, S. López Soliño [ID](#)⁴⁷, Q. Lu [ID](#)¹⁵,
 C. Lucarelli [ID](#)⁴⁹, D. Lucchesi [ID](#)^{33,q}, M. Lucio Martinez [ID](#)⁴⁸, Y. Luo [ID](#)⁶, A. Lupato [ID](#)^{33,i},
 E. Luppi [ID](#)^{26,l}, K. Lynch [ID](#)²³, X.-R. Lyu [ID](#)⁷, G. M. Ma [ID](#)^{4,c}, S. Maccolini [ID](#)¹⁹, F. Machefer [ID](#)¹⁴,
 F. Maciuc [ID](#)⁴³, B. Mack [ID](#)⁶⁹, I. Mackay [ID](#)⁶⁴, L. M. Mackey [ID](#)⁶⁹, L.R. Madhan Mohan [ID](#)⁵⁶, M. J.
 Madurai [ID](#)⁵⁴, D. Magdalinski [ID](#)³⁸, D. Maisuzenko [ID](#)⁴⁴, J.J. Malczewski [ID](#)⁴¹, S. Malde [ID](#)⁶⁴,
 L. Malentacca [ID](#)⁴⁹, A. Malinin [ID](#)⁴⁴, T. Maltsev [ID](#)⁴⁴, G. Manca [ID](#)^{32,k}, G. Mancinelli [ID](#)¹³,
 C. Mancuso [ID](#)¹⁴, R. Manera Escalero [ID](#)⁴⁵, F. M. Manganella [ID](#)³⁷, D. Manuzzi [ID](#)²⁵,
 D. Marangotto [ID](#)^{30,n}, J.F. Marchand [ID](#)¹⁰, R. Marchevski [ID](#)⁵⁰, U. Marconi [ID](#)²⁵, E. Mariani [ID](#)¹⁶,
 S. Mariani [ID](#)⁴⁹, C. Marin Benito [ID](#)⁴⁵, J. Marks [ID](#)²², A.M. Marshall [ID](#)⁵⁵, L. Martel [ID](#)⁶⁴,
 G. Martelli [ID](#)³⁴, G. Martellotti [ID](#)³⁶, L. Martinazzoli [ID](#)⁴⁹, M. Martinelli [ID](#)^{31,o}, D. Martinez Gomez [ID](#)⁸⁰,
 D. Martinez Santos [ID](#)⁸³, F. Martinez Vidal [ID](#)⁴⁸, A. Martorell i Granollers [ID](#)⁴⁶, A. Massafferri [ID](#)²,
 R. Matev [ID](#)⁴⁹, A. Mathad [ID](#)⁴⁹, V. Matiunin [ID](#)⁴⁴, C. Matteuzzi [ID](#)⁶⁹, K.R. Mattioli [ID](#)¹⁵, A. Mauri [ID](#)⁶²,
 E. Maurice [ID](#)¹⁵, J. Mauricio [ID](#)⁴⁵, P. Mayencourt [ID](#)⁵⁰, J. Mazorra de Cos [ID](#)⁴⁸, M. Mazurek [ID](#)⁴²,
 M. McCann [ID](#)⁶², T.H. McGrath [ID](#)⁶³, N.T. McHugh [ID](#)⁶⁰, A. McNab [ID](#)⁶³, R. McNulty [ID](#)²³,
 B. Meadows [ID](#)⁶⁶, G. Meier [ID](#)¹⁹, D. Melnychuk [ID](#)⁴², F. M. Meng [ID](#)^{4,c}, M. Merk [ID](#)^{38,81}, A. Merli [ID](#)^{50,30},
 L. Meyer Garcia [ID](#)⁶⁷, D. Miao [ID](#)^{5,7}, H. Miao [ID](#)⁷, M. Mikhasenko [ID](#)⁷⁷, D.A. Milanes [ID](#)^{76,y},
 A. Minotti [ID](#)^{31,o}, E. Minucci [ID](#)²⁸, T. Miralles [ID](#)¹¹, B. Mitreska [ID](#)¹⁹, D.S. Mitzel [ID](#)¹⁹, A. Modak [ID](#)⁵⁸,
 L. Moeser [ID](#)¹⁹, R.A. Mohammed [ID](#)⁶⁴, R.D. Moise [ID](#)¹⁷, E. F. Molina Cardenas [ID](#)⁸⁶,
 T. Mombächer [ID](#)⁴⁹, M. Monk [ID](#)^{57,1}, S. Monteil [ID](#)¹¹, A. Morcillo Gomez [ID](#)⁴⁷, G. Morello [ID](#)²⁸,
 M.J. Morello [ID](#)^{35,s}, M.P. Morgenthaler [ID](#)²², J. Moron [ID](#)⁴⁰, W. Morren [ID](#)³⁸, A.B. Morris [ID](#)⁴⁹,

A.G. Morris [ID](#)¹³, R. Mountain [ID](#)⁶⁹, H. Mu [ID](#)^{4,c}, Z. M. Mu [ID](#)⁶, E. Muhammad [ID](#)⁵⁷, F. Muheim [ID](#)⁵⁹, M. Mulder [ID](#)⁸⁰, K. Müller [ID](#)⁵¹, F. Muñoz-Rojas [ID](#)⁹, R. Murta [ID](#)⁶², V. Mytrochenko [ID](#)⁵², P. Naik [ID](#)⁶¹, T. Nakada [ID](#)⁵⁰, R. Nandakumar [ID](#)⁵⁸, T. Nanut [ID](#)⁴⁹, I. Nasteva [ID](#)³, M. Needham [ID](#)⁵⁹, E. Nekrasova [ID](#)⁴⁴, N. Neri [ID](#)^{30,n}, S. Neubert [ID](#)¹⁸, N. Neufeld [ID](#)⁴⁹, P. Neustroev [ID](#)⁴⁴, J. Nicolini [ID](#)⁴⁹, D. Nicotra [ID](#)⁸¹, E.M. Niel [ID](#)¹⁵, N. Nikitin [ID](#)⁴⁴, Q. Niu [ID](#)⁷³, P. Nogarolli [ID](#)³, P. Nogga [ID](#)¹⁸, C. Normand [ID](#)⁵⁵, J. Novoa Fernandez [ID](#)⁴⁷, G. Nowak [ID](#)⁶⁶, C. Nunez [ID](#)⁸⁶, H. N. Nur [ID](#)⁶⁰, A. Oblakowska-Mucha [ID](#)⁴⁰, V. Obraztsov [ID](#)⁴⁴, T. Oeser [ID](#)¹⁷, A. Okhotnikov [ID](#)⁴⁴, O. Okhrimenko [ID](#)⁵³, R. Oldeman [ID](#)^{32,k}, F. Oliva [ID](#)^{59,49}, E. Olivart Pino [ID](#)⁴⁵, M. Olocco [ID](#)¹⁹, C.J.G. Onderwater [ID](#)⁸¹, R.H. O’Neil [ID](#)⁴⁹, D. Osthues [ID](#)¹⁹, J.M. Otalora Goicochea [ID](#)³, P. Owen [ID](#)⁵¹, A. Oyanguren [ID](#)⁴⁸, O. Ozcelik [ID](#)⁴⁹, F. Paciolla [ID](#)^{35,w}, A. Padee [ID](#)⁴², K.O. Padeken [ID](#)¹⁸, B. Pagare [ID](#)⁴⁷, T. Pajero [ID](#)⁴⁹, A. Palano [ID](#)²⁴, M. Palutan [ID](#)²⁸, C. Pan [ID](#)⁷⁴, X. Pan [ID](#)^{4,c}, S. Panebianco [ID](#)¹², G. Panshin [ID](#)⁵, L. Paolucci [ID](#)⁵⁷, A. Papanestis [ID](#)⁵⁸, M. Pappagallo [ID](#)^{24,h}, L.L. Pappalardo [ID](#)²⁶, C. Pappenheimer [ID](#)⁶⁶, C. Parkes [ID](#)⁶³, D. Parmar [ID](#)⁷⁷, B. Passalacqua [ID](#)^{26,l}, G. Passaleva [ID](#)²⁷, D. Passaro [ID](#)^{35,s,49}, A. Pastore [ID](#)²⁴, M. Patel [ID](#)⁶², J. Patoc [ID](#)⁶⁴, C. Patrignani [ID](#)^{25,j}, A. Paul [ID](#)⁶⁹, C.J. Pawley [ID](#)⁸¹, A. Pellegrino [ID](#)³⁸, J. Peng [ID](#)^{5,7}, X. Peng [ID](#)⁷³, M. Pepe Altarelli [ID](#)²⁸, S. Perazzini [ID](#)²⁵, D. Pereima [ID](#)⁴⁴, H. Pereira Da Costa [ID](#)⁶⁸, A. Pereiro Castro [ID](#)⁴⁷, C. Perez [ID](#)⁴⁶, P. Perret [ID](#)¹¹, A. Perrevoort [ID](#)⁸⁰, A. Perro [ID](#)^{49,13}, M.J. Peters [ID](#)⁶⁶, K. Petridis [ID](#)⁵⁵, A. Petrolini [ID](#)^{29,m}, J. P. Pfaller [ID](#)⁶⁶, H. Pham [ID](#)⁶⁹, L. Pica [ID](#)³⁵, M. Piccini [ID](#)³⁴, L. Piccolo [ID](#)³², B. Pietrzyk [ID](#)¹⁰, G. Pietrzyk [ID](#)¹⁴, R. N. Pilato [ID](#)⁶¹, D. Pinci [ID](#)³⁶, F. Pisani [ID](#)⁴⁹, M. Pizzichemi [ID](#)^{31,o,49}, V. M. Placinta [ID](#)⁴³, M. Plo Casasus [ID](#)⁴⁷, T. Poeschl [ID](#)⁴⁹, F. Polci [ID](#)¹⁶, M. Poli Lener [ID](#)²⁸, A. Poluektov [ID](#)¹³, N. Polukhina [ID](#)⁴⁴, I. Polyakov [ID](#)⁶³, E. Polycarpo [ID](#)³, S. Ponce [ID](#)⁴⁹, D. Popov [ID](#)^{7,49}, S. Poslavskii [ID](#)⁴⁴, K. Prasanth [ID](#)⁵⁹, C. Prouve [ID](#)⁸³, D. Provenzano [ID](#)^{32,k,49}, V. Pugatch [ID](#)⁵³, G. Punzi [ID](#)^{35,t}, S. Qasim [ID](#)⁵¹, Q. Q. Qian [ID](#)⁶, W. Qian [ID](#)⁷, N. Qin [ID](#)^{4,c}, S. Qu [ID](#)^{4,c}, R. Quagliani [ID](#)⁴⁹, R.I. Rabadan Trejo [ID](#)⁵⁷, J.H. Rademacker [ID](#)⁵⁵, M. Rama [ID](#)³⁵, M. Ramírez García [ID](#)⁸⁶, V. Ramos De Oliveira [ID](#)⁷⁰, M. Ramos Pernas [ID](#)⁵⁷, M.S. Rangel [ID](#)³, F. Ratnikov [ID](#)⁴⁴, G. Raven [ID](#)³⁹, M. Rebollo De Miguel [ID](#)⁴⁸, F. Redi [ID](#)^{30,i}, J. Reich [ID](#)⁵⁵, F. Reiss [ID](#)²⁰, Z. Ren [ID](#)⁷, P.K. Resmi [ID](#)⁶⁴, M. Ribalda Galvez [ID](#)⁴⁵, R. Ribatti [ID](#)⁵⁰, G. Ricart [ID](#)^{15,12}, D. Riccardi [ID](#)^{35,s}, S. Ricciardi [ID](#)⁵⁸, K. Richardson [ID](#)⁶⁵, M. Richardson-Slipper [ID](#)⁵⁹, K. Rinnert [ID](#)⁶¹, P. Robbe [ID](#)^{14,49}, G. Robertson [ID](#)⁶⁰, E. Rodrigues [ID](#)⁶¹, A. Rodriguez Alvarez [ID](#)⁴⁵, E. Rodriguez Fernandez [ID](#)⁴⁷, J.A. Rodriguez Lopez [ID](#)⁷⁶, E. Rodriguez Rodriguez [ID](#)⁴⁹, J. Roensch [ID](#)¹⁹, A. Rogachev [ID](#)⁴⁴, A. Rogovskiy [ID](#)⁵⁸, D.L. Rolf [ID](#)¹⁹, P. Roloff [ID](#)⁴⁹, V. Romanovskiy [ID](#)⁶⁶, A. Romero Vidal [ID](#)⁴⁷, G. Romolini [ID](#)^{26,49}, F. Ronchetti [ID](#)⁵⁰, T. Rong [ID](#)⁶, M. Rotondo [ID](#)²⁸, S. R. Roy [ID](#)²², M.S. Rudolph [ID](#)⁶⁹, M. Ruiz Diaz [ID](#)²², R.A. Ruiz Fernandez [ID](#)⁴⁷, J. Ruiz Vidal [ID](#)⁸¹, J. Saavedra-Arias [ID](#)⁹, J.J. Saborido Silva [ID](#)⁴⁷, R. Sadek [ID](#)¹⁵, N. Sagidova [ID](#)⁴⁴, D. Sahoo [ID](#)⁷⁸, N. Sahoo [ID](#)⁵⁴, B. Saitta [ID](#)^{32,k}, M. Salomoni [ID](#)^{31,49,o}, I. Sanderswood [ID](#)⁴⁸, R. Santacesaria [ID](#)³⁶, C. Santamarina Rios [ID](#)⁴⁷, M. Santimaria [ID](#)²⁸, L. Santoro [ID](#)², E. Santovetti [ID](#)³⁷, A. Saputi [ID](#), D. Saranin [ID](#)⁴⁴, A. Sarnatskiy [ID](#)⁸⁰, G. Sarpis [ID](#)⁵⁹, M. Sarpis [ID](#)⁷⁹, C. Satriano [ID](#)^{36,u}, M. Saur [ID](#)⁷³, D. Savrina [ID](#)⁴⁴, H. Sazak [ID](#)¹⁷, F. Sborzacchi [ID](#)^{49,28}, A. Scarabotto [ID](#)¹⁹, S. Schael [ID](#)¹⁷, S. Scherl [ID](#)⁶¹, M. Schiller [ID](#)²², H. Schindler [ID](#)⁴⁹, M. Schmelling [ID](#)²¹, B. Schmidt [ID](#)⁴⁹, S. Schmitt [ID](#)¹⁷, H. Schmitz [ID](#)¹⁸, O. Schneider [ID](#)⁵⁰, A. Schopper [ID](#)⁶², N. Schulte [ID](#)¹⁹, M.H. Schune [ID](#)¹⁴, G. Schwering [ID](#)¹⁷, B. Sciascia [ID](#)²⁸, A. Sciucati [ID](#)⁴⁹, I. Segal [ID](#)⁷⁷, S. Sellam [ID](#)⁴⁷, A. Semennikov [ID](#)⁴⁴, T. Senger [ID](#)⁵¹, M. Senghi Soares [ID](#)³⁹, A. Sergi [ID](#)^{29,m}, N. Serra [ID](#)⁵¹, L. Sestini [ID](#)²⁷, A. Seuthe [ID](#)¹⁹, B. Sevilla Sanjuan [ID](#)⁴⁶, Y. Shang [ID](#)⁶, D.M. Shangase [ID](#)⁸⁶, M. Shapkin [ID](#)⁴⁴, R. S. Sharma [ID](#)⁶⁹, I. Shchemerov [ID](#)⁴⁴, L. Shchutska [ID](#)⁵⁰, T. Shears [ID](#)⁶¹, L. Shekhtman [ID](#)⁴⁴, Z. Shen [ID](#)³⁸, S. Sheng [ID](#)^{5,7},

V. Shevchenko ⁴⁴, B. Shi ⁷, Q. Shi ⁷, W. S. Shi ⁷², Y. Shimizu ¹⁴, E. Shmanin ²⁵, R. Shorkin ⁴⁴, J.D. Shupperd ⁶⁹, R. Silva Coutinho ⁶⁹, G. Simi ^{33,q}, S. Simone ^{24,h}, M. Singha ⁷⁸, N. Skidmore ⁵⁷, T. Skwarnicki ⁶⁹, M.W. Slater ⁵⁴, E. Smith ⁶⁵, K. Smith ⁶⁸, M. Smith ⁶², L. Soares Lavra ⁵⁹, M.D. Sokoloff ⁶⁶, F.J.P. Soler ⁶⁰, A. Solomin ⁵⁵, A. Solovev ⁴⁴, N. S. Sommerfeld ¹⁸, R. Song ¹, Y. Song ⁵⁰, Y. Song ^{4,c}, Y. S. Song ⁶, F.L. Souza De Almeida ⁶⁹, B. Souza De Paula ³, E. Spadaro Norella ^{29,m}, E. Spedicato ²⁵, J.G. Speer ¹⁹, E. Spiridenkov ⁴⁴, P. Spradlin ⁶⁰, V. Sriskaran ⁴⁹, F. Stagni ⁴⁹, M. Stahl ⁷⁷, S. Stahl ⁴⁹, S. Stanislaus ⁶⁴, M. Stefaniak ⁸⁷, E.N. Stein ⁴⁹, O. Steinkamp ⁵¹, H. Stevens ¹⁹, D. Strelakina ⁴⁴, Y. Su ⁷, F. Suljik ⁶⁴, J. Sun ³², L. Sun ⁷⁴, D. Sundfeld ², W. Sutcliffe ⁵¹, K. Swientek ⁴⁰, F. Swystun ⁵⁶, A. Szabelski ⁴², T. Szumlak ⁴⁰, Y. Tan ^{4,c}, Y. Tang ⁷⁴, Y. T. Tang ⁷, M.D. Tat ²², A. Terentev ⁴⁴, F. Terzuoli ^{35,w}, F. Teubert ⁴⁹, E. Thomas ⁴⁹, D.J.D. Thompson ⁵⁴, A. R. Thomson-Strong ⁵⁹, H. Tilquin ⁶², V. Tisserand ¹¹, S. T’Jampens ¹⁰, M. Tobin ⁵, L. Tomassetti ^{26,l}, G. Tonani ³⁰, X. Tong ⁶, T. Tork ³⁰, D. Torres Machado ², L. Toscano ¹⁹, D.Y. Tou ^{4,c}, C. Trippel ⁴⁶, G. Tuci ²², N. Tuning ³⁸, L.H. Uecker ²², A. Ukleja ⁴⁰, D.J. Unverzagt ²², A. Upadhyay ⁴⁹, B. Urbach ⁵⁹, A. Usachov ³⁹, A. Ustyuzhanin ⁴⁴, U. Uwer ²², V. Vagnoni ²⁵, V. Valcarce Cadenas ⁴⁷, G. Valenti ²⁵, N. Valls Canudas ⁴⁹, J. van Eldik ⁴⁹, H. Van Hecke ⁶⁸, E. van Herwijnen ⁶², C.B. Van Hulse ^{47,z}, R. Van Laak ⁵⁰, M. van Veghel ³⁸, G. Vasquez ⁵¹, R. Vazquez Gomez ⁴⁵, P. Vazquez Regueiro ⁴⁷, C. Vázquez Sierra ⁸³, S. Vecchi ²⁶, J.J. Velthuis ⁵⁵, M. Veltri ^{27,x}, A. Venkateswaran ⁵⁰, M. Verdognia ³², M. Vesterinen ⁵⁷, W. Vetens ⁶⁹, D. Vico Benet ⁶⁴, P. Vidrier Villalba ⁴⁵, M. Vieites Diaz ⁴⁷, X. Vilasis-Cardona ⁴⁶, E. Vilella Figueras ⁶¹, A. Villa ²⁵, P. Vincent ¹⁶, B. Vivacqua ³, F.C. Volle ⁵⁴, D. vom Bruch ¹³, N. Voropaev ⁴⁴, K. Vos ⁸¹, C. Vrahas ⁵⁹, J. Wagner ¹⁹, J. Walsh ³⁵, E.J. Walton ^{1,57}, G. Wan ⁶, A. Wang ⁷, B. Wang ⁵, C. Wang ²², G. Wang ⁸, H. Wang ⁷³, J. Wang ⁶, J. Wang ⁵, J. Wang ^{4,c}, J. Wang ⁷⁴, M. Wang ⁴⁹, N. W. Wang ⁷, R. Wang ⁵⁵, X. Wang ⁸, X. Wang ⁷², X. W. Wang ⁶², Y. Wang ⁷⁵, Y. Wang ⁶, Y. W. Wang ⁷³, Z. Wang ¹⁴, Z. Wang ^{4,c}, Z. Wang ³⁰, J.A. Ward ^{57,1}, M. Waterlaat ⁴⁹, N.K. Watson ⁵⁴, D. Websdale ⁶², Y. Wei ⁶, J. Wendel ⁸³, B.D.C. Westhenry ⁵⁵, C. White ⁵⁶, M. Whitehead ⁶⁰, E. Whiter ⁵⁴, A.R. Wiederhold ⁶³, D. Wiedner ¹⁹, G. Wilkinson ^{64,49}, M.K. Wilkinson ⁶⁶, M. Williams ⁶⁵, M. J. Williams ⁴⁹, M.R.J. Williams ⁵⁹, R. Williams ⁵⁶, Z. Williams ⁵⁵, F.F. Wilson ⁵⁸, M. Winn ¹², W. Wislicki ⁴², M. Witek ⁴¹, L. Witola ¹⁹, T. Wolf ²², G. Wormser ¹⁴, S.A. Wotton ⁵⁶, H. Wu ⁶⁹, J. Wu ⁸, X. Wu ⁷⁴, Y. Wu ^{6,56}, Z. Wu ⁷, K. Wyllie ⁴⁹, S. Xian ⁷², Z. Xiang ⁵, Y. Xie ⁸, T. X. Xing ³⁰, A. Xu ^{35,s}, L. Xu ^{4,c}, L. Xu ^{4,c}, M. Xu ⁴⁹, Z. Xu ⁴⁹, Z. Xu ⁷, Z. Xu ⁵, K. Yang ⁶², X. Yang ⁶, Y. Yang ²⁹, Z. Yang ⁶, V. Yeroshenko ¹⁴, H. Yeung ⁶³, H. Yin ⁸, X. Yin ⁷, C. Y. Yu ⁶, J. Yu ⁷¹, X. Yuan ⁵, Y. Yuan ^{5,7}, E. Zaffaroni ⁵⁰, M. Zavertyaev ²¹, M. Zdybal ⁴¹, F. Zenesini ²⁵, C. Zeng ^{5,7}, M. Zeng ^{4,c}, C. Zhang ⁶, D. Zhang ⁸, J. Zhang ⁷, L. Zhang ^{4,c}, R. Zhang ⁸, S. Zhang ⁷¹, S. Zhang ⁶⁴, Y. Zhang ⁶, Y. Z. Zhang ^{4,c}, Z. Zhang ^{4,c}, Y. Zhao ²², A. Zhelezov ²², S. Z. Zheng ⁶, X. Z. Zheng ^{4,c}, Y. Zheng ⁷, T. Zhou ⁶, X. Zhou ⁸, Y. Zhou ⁷, V. Zhovkovska ⁵⁷, L. Z. Zhu ⁷, X. Zhu ^{4,c}, X. Zhu ⁸, Y. Zhu ¹⁷, V. Zhukov ¹⁷, J. Zhuo ⁴⁸, Q. Zou ^{5,7}, D. Zuliani ^{33,q}, G. Zunica ⁵⁰

¹ School of Physics and Astronomy, Monash University, Melbourne, Australia

² Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

³ Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

- ⁴ *Department of Engineering Physics, Tsinghua University, Beijing, China*
- ⁵ *Institute Of High Energy Physics (IHEP), Beijing, China*
- ⁶ *School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*
- ⁷ *University of Chinese Academy of Sciences, Beijing, China*
- ⁸ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China*
- ⁹ *Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica*
- ¹⁰ *Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*
- ¹¹ *Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*
- ¹² *Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Saclay, France, Gif-Sur-Yvette, France*
- ¹³ *Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*
- ¹⁴ *Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
- ¹⁵ *Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*
- ¹⁶ *LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*
- ¹⁷ *I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁸ *Universität Bonn — Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany*
- ¹⁹ *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ²⁰ *Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg, Germany*
- ²¹ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ²² *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ²³ *School of Physics, University College Dublin, Dublin, Ireland*
- ²⁴ *INFN Sezione di Bari, Bari, Italy*
- ²⁵ *INFN Sezione di Bologna, Bologna, Italy*
- ²⁶ *INFN Sezione di Ferrara, Ferrara, Italy*
- ²⁷ *INFN Sezione di Firenze, Firenze, Italy*
- ²⁸ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁹ *INFN Sezione di Genova, Genova, Italy*
- ³⁰ *INFN Sezione di Milano, Milano, Italy*
- ³¹ *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ³² *INFN Sezione di Cagliari, Monserrato, Italy*
- ³³ *INFN Sezione di Padova, Padova, Italy*
- ³⁴ *INFN Sezione di Perugia, Perugia, Italy*
- ³⁵ *INFN Sezione di Pisa, Pisa, Italy*
- ³⁶ *INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³⁷ *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³⁸ *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³⁹ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ⁴⁰ *AGH — University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ⁴¹ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ⁴² *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴³ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴⁴ *Authors affiliated with an institute formerly covered by a cooperation agreement with CERN.*
- ⁴⁵ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁶ *La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴⁷ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁸ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia — CSIC, Valencia, Spain*
- ⁴⁹ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁵⁰ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁵¹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵² *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵³ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*

- ⁵⁴ *School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ⁵⁵ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁶ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁷ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁸ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁹ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁶⁰ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁶¹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁶² *Imperial College London, London, United Kingdom*
- ⁶³ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁶⁴ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶⁵ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶⁶ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁷ *University of Maryland, College Park, MD, United States*
- ⁶⁸ *Los Alamos National Laboratory (LANL), Los Alamos, NM, United States*
- ⁶⁹ *Syracuse University, Syracuse, NY, United States*
- ⁷⁰ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ³*
- ⁷¹ *School of Physics and Electronics, Hunan University, Changsha City, China, associated to ⁸*
- ⁷² *Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ⁴*
- ⁷³ *Lanzhou University, Lanzhou, China, associated to ⁵*
- ⁷⁴ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ⁴*
- ⁷⁵ *Henan Normal University, Xinxiang, China, associated to ⁸*
- ⁷⁶ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹⁶*
- ⁷⁷ *Ruhr Universitaet Bochum, Fakultae f. Physik und Astronomie, Bochum, Germany, associated to ¹⁹*
- ⁷⁸ *Eotvos Lorand University, Budapest, Hungary, associated to ⁴⁹*
- ⁷⁹ *Faculty of Physics, Vilnius University, Vilnius, Lithuania, associated to ²⁰*
- ⁸⁰ *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³⁸*
- ⁸¹ *Universiteit Maastricht, Maastricht, Netherlands, associated to ³⁸*
- ⁸² *Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ⁴¹*
- ⁸³ *Universidade da Coruña, A Coruña, Spain, associated to ⁴⁶*
- ⁸⁴ *Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁶⁰*
- ⁸⁵ *Taras Schevchenko University of Kyiv, Faculty of Physics, Kyiv, Ukraine, associated to ¹⁴*
- ⁸⁶ *University of Michigan, Ann Arbor, MI, United States, associated to ⁶⁹*
- ⁸⁷ *Ohio State University, Columbus, United States, associated to ⁶⁸*

^a *Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil*

^b *Department of Physics and Astronomy, University of Victoria, Victoria, Canada*

^c *Center for High Energy Physics, Tsinghua University, Beijing, China*

^d *Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China*

^e *LIP6, Sorbonne Université, Paris, France*

^f *Lamarr Institute for Machine Learning and Artificial Intelligence, Dortmund, Germany*

^g *Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras*

^h *Università di Bari, Bari, Italy*

ⁱ *Università di Bergamo, Bergamo, Italy*

^j *Università di Bologna, Bologna, Italy*

^k *Università di Cagliari, Cagliari, Italy*

^l *Università di Ferrara, Ferrara, Italy*

^m *Università di Genova, Genova, Italy*

ⁿ *Università degli Studi di Milano, Milano, Italy*

^o *Università degli Studi di Milano-Bicocca, Milano, Italy*

^p *Università di Modena e Reggio Emilia, Modena, Italy*

^q *Università di Padova, Padova, Italy*

- ^r *Università di Perugia, Perugia, Italy*
^s *Scuola Normale Superiore, Pisa, Italy*
^t *Università di Pisa, Pisa, Italy*
^u *Università della Basilicata, Potenza, Italy*
^v *Università di Roma Tor Vergata, Roma, Italy*
^w *Università di Siena, Siena, Italy*
^x *Università di Urbino, Urbino, Italy*
^y *Universidad de Ingeniería y Tecnología (UTEC), Lima, Peru*
^z *Universidad de Alcalá, Alcalá de Henares, Spain*
^{aa} *Facultad de Ciencias Físicas, Madrid, Spain*
[†] *Deceased*