

ANALYSIS & PDE

Volume 13

No. 2

2020

CLARA ANTONUCCI, MASSIMO GOBBINO,
MATTEO MIGLIORINI AND NICOLA PICENNI

**OPTIMAL CONSTANTS FOR A NONLOCAL APPROXIMATION OF
SOBOLEV NORMS AND TOTAL VARIATION**

OPTIMAL CONSTANTS FOR A NONLOCAL APPROXIMATION OF SOBOLEV NORMS AND TOTAL VARIATION

CLARA ANTONUCCI, MASSIMO GOBBINO, MATTEO MIGLIORINI AND NICOLA PICENNI

We consider the family of nonlocal and nonconvex functionals proposed and investigated by J. Bourgain, H. Brezis and H.-M. Nguyen in a series of papers of the last decade. It was known that this family of functionals Gamma-converges to a suitable multiple of the Sobolev norm or the total variation, depending on the summability exponent, but the exact constants and the structure of recovery families were still unknown, even in dimension 1.

We prove a Gamma-convergence result with explicit values of the constants in any space dimension. We also show the existence of recovery families consisting of smooth functions with compact support.

The key point is reducing the problem first to dimension 1, and then to a finite combinatorial rearrangement inequality.

1. Introduction

Let $p \geq 1$ and $\delta > 0$ be real numbers, let d be a positive integer, and let $\Omega \subseteq \mathbb{R}^d$ be an open set. For every measurable function $u : \Omega \rightarrow \mathbb{R}$ we set

$$\Lambda_{\delta,p}(u, \Omega) := \iint_{I(\delta,u,\Omega)} \frac{\delta^p}{|y-x|^{d+p}} dx dy, \quad (1-1)$$

where

$$I(\delta, u, \Omega) := \{(x, y) \in \Omega^2 : |u(y) - u(x)| > \delta\}.$$

Nonconvex and nonlocal functionals of this type appeared in a paper by J. Bourgain, H. Brezis and P. Mironeanu [Bourgain et al. 2005]; see Open Problem 2 of that work. Subsequently, the family (1-1) was investigated in a series of papers by H.-M. Nguyen [2006; 2007; 2008; 2011; 2014], J. Bourgain and H.-M. Nguyen [2006], and H. Brezis and H.-M. Nguyen [2018]; see also [Brezis 2015; Brezis and Nguyen 2017].

We point out that the dependence on u is just on the integration set. The fixed integrand is divergent on the diagonal $y = x$, and the integration set is closer to the diagonal where the gradient of u is large. This suggests that $\Lambda_{\delta,p}(u, \Omega)$ is proportional, in the limit as $\delta \rightarrow 0^+$, to some norm of the gradient of u ,

MSC2010: 26B30, 46E35.

Keywords: Gamma-convergence, Sobolev spaces, bounded-variation functions, monotone rearrangement, nonlocal functional, nonconvex functional.

and more precisely to the functional

$$\Lambda_{0,p}(u, \Omega) := \begin{cases} \int_{\Omega} |\nabla u(x)|^p dx & \text{if } p > 1 \text{ and } u \in W^{1,p}(\Omega), \\ \text{total variation of } u \text{ in } \Omega & \text{if } p = 1 \text{ and } u \in \text{BV}(\Omega), \\ +\infty & \text{otherwise.} \end{cases} \tag{1-2}$$

It is natural to compare the family (1-1) with the classical approximations of Sobolev or BV norms, based on nonlocal convex functionals such as

$$G_{\varepsilon,p}(u, \Omega) := \iint_{\Omega} \frac{|u(y) - u(x)|^p}{|y - x|^p} \rho_{\varepsilon}(|y - x|) dx dy, \tag{1-3}$$

where gradients are replaced by finite differences weighted by a suitable family ρ_{ε} of mollifiers. The idea of approximating integrals of the gradient with double integrals of difference quotients, where all pairs of distinct points interact, has been considered independently by many authors in different contexts. For example, E. De Giorgi proposed an approximation of this kind to the Mumford–Shah functional in any space dimension, in order to overcome the anisotropy of the discrete approximation [Chambolle 1995]. The resulting theory appears in [Gobbino 1998] and was then extended in [Gobbino and Mora 2001] to more general free discontinuity problems, and in particular to Sobolev and BV spaces. In the same years, the case of Sobolev and BV norms was considered in detail in [Bourgain et al. 2001]; see also [Ponce 2004].

The result, as expected, is that the family $G_{\varepsilon,p}(u, \mathbb{R}^d)$ converges as $\varepsilon \rightarrow 0^+$ to a suitable multiple of $\Lambda_{0,p}(u, \mathbb{R}^d)$, both in the sense of pointwise convergence, and in the sense of De Giorgi’s Gamma-convergence. This provides a characterization of Sobolev functions (if $p > 1$), and of bounded variation functions (if $p = 1$), as those functions for which the pointwise limit or the Gamma-limit is finite.

From the heuristic point of view, the nonconvex approximating family (1-1) seems to follow a different paradigm. Indeed, it was observed by J.-M. Morel, as quoted on page 4 of the transparencies of the presentation [Brezis 2016], that this definition involves some sort of “vertical slicing” that evokes the definition of integral *à la Lebesgue*, in contrast to the definition *à la Riemann* that seems closer to the “horizontal slicing” of the finite differences in (1-3).

From the mathematical point of view, the asymptotic behavior of (1-1) exhibits some unexpected features. In order to state the precise results, let us introduce some notation. Let $\mathbb{S}^{d-1} := \{\sigma \in \mathbb{R}^d : |\sigma| = 1\}$ denote the unit sphere in \mathbb{R}^d . For every $p \geq 1$ we consider the geometric constant

$$G_{d,p} := \int_{\mathbb{S}^{d-1}} |\langle v, \sigma \rangle|^p d\sigma, \tag{1-4}$$

where v is any element of \mathbb{S}^{d-1} (of course the value of $G_{d,p}$ does not depend on the choice of v), and the integration is intended with respect to the $(d-1)$ -dimensional Hausdorff measure. The value of $G_{d,p}$ can be explicitly computed in terms of special functions through Beta integrals. It turns out that $G_{d,p} = 2$ for every p if $d = 1$, and

$$G_{d,p} = \text{meas}(\mathbb{S}^{d-2}) \int_{-\pi/2}^{\pi/2} (\cos \theta)^p \cdot |\sin \theta|^{d-2} d\theta = \frac{2\pi^{(d-1)/2} \Gamma((p+1)/2)}{\Gamma((p+d)/2)} \quad \text{for all } d \geq 2.$$

The main convergence results obtained so far can be summed up as follows.

- *Pointwise convergence for $p > 1$.* For every $p > 1$ it turns out that

$$\lim_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u, \mathbb{R}^d) = \frac{1}{p} G_{d,p} \Lambda_{0,p}(u, \mathbb{R}^d) \quad \text{for all } u \in L^p(\mathbb{R}^d). \tag{1-5}$$

- *Pointwise convergence for $p = 1$.* In the case $p = 1$, equality (1-5) holds true for every $u \in C_c^1(\mathbb{R}^d)$, but there do exist functions $u \in W^{1,1}(\mathbb{R}^d)$ for which the left-hand side is infinite (while of course the right-hand side is finite). A precise characterization of equality cases is still unknown.
- *Gamma-convergence for every $p \geq 1$.* For every $p \geq 1$ there exists a constant $C_{d,p}$ such that

$$\Gamma\text{-}\lim_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u, \mathbb{R}^d) = \frac{1}{p} G_{d,p} C_{d,p} \Lambda_{0,p}(u, \mathbb{R}^d) \quad \text{for all } u \in L^p(\mathbb{R}^d),$$

where the Gamma-limit is intended with respect to the usual metric of $L^p(\mathbb{R}^d)$ (but the result would be the same with respect to the convergence in $L^1(\mathbb{R}^d)$ or in measure). Moreover, it was proved that $C_{d,p} \in (0, 1)$; namely the Gamma-limit is always nontrivial but different from the pointwise limit.

As a consequence, again one can characterize the Sobolev space $W^{1,p}(\mathbb{R}^d)$ as the set of functions in $L^p(\mathbb{R}^d)$ for which the pointwise limit or the Gamma-limit is finite. As for $BV(\mathbb{R}^d)$, in this setting it can be characterized only through the Gamma-limit.

Some problems remained open, and were stated explicitly in [Nguyen 2011; Brezis and Nguyen 2018]:

Question 1. What is the exact value of $C_{d,p}$, at least in the case $d = 1$?

Question 2. Does $C_{d,p}$ depend on d ?

Question 3. Do there exist recovery families made up of continuous functions, or even of functions of class C^∞ ?

In this paper we answer these three questions. Concerning Questions 1 and 2, we prove that $C_{d,p}$ does not depend on d , and coincides with the value C_p conjectured in [Nguyen 2007] (see also [Nguyen 2011, Open question 2]) for the 1-dimensional case, namely

$$C_p := \begin{cases} \frac{1}{p-1} \left(1 - \frac{1}{2^{p-1}}\right) & \text{if } p > 1, \\ \log 2 & \text{if } p = 1. \end{cases} \tag{1-6}$$

Concerning the third question, we prove that smooth recovery families do exist. Our main result is the following.

Theorem 1.1 (Gamma-convergence). *Let us consider the functionals $\Lambda_{\delta,p}$ and $\Lambda_{0,p}$ defined in (1-1) and (1-2), respectively.*

Then for every positive integer d and every real number $p \geq 1$ it turns out that

$$\Gamma\text{-}\lim_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u, \mathbb{R}^d) = \frac{1}{p} G_{d,p} C_p \Lambda_{0,p}(u, \mathbb{R}^d) \quad \text{for all } u \in L^p(\mathbb{R}^d),$$

where $G_{d,p}$ is the geometric constant defined in (1-4), and C_p is the constant defined in (1-6). In particular, the following two statements hold true:

(1) (liminf inequality) For every family $\{u_\delta\}_{\delta>0} \subseteq L^p(\mathbb{R}^d)$, with $u_\delta \rightarrow u$ in $L^p(\mathbb{R}^d)$ as $\delta \rightarrow 0^+$, it turns out that

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, \mathbb{R}^d) \geq \frac{1}{p} G_{d,p} C_p \Lambda_{0,p}(u, \mathbb{R}^d). \tag{1-7}$$

(2) (limsup inequality) For every $u \in L^p(\mathbb{R}^d)$ there exists a family $\{u_\delta\}_{\delta>0} \subseteq L^p(\mathbb{R}^d)$, with $u_\delta \rightarrow u$ in $L^p(\mathbb{R}^d)$ as $\delta \rightarrow 0^+$, such that

$$\limsup_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, \mathbb{R}^d) \leq \frac{1}{p} G_{d,p} C_p \Lambda_{0,p}(u, \mathbb{R}^d).$$

We can also assume that the family $\{u_\delta\}$ consists of functions of class C^∞ with compact support.

The proof of this result requires a different approach to the problem, which we briefly sketch below. In previous literature, see [Nguyen 2011, formula (1.3)] or [Brezis and Nguyen 2018, formula (1.12)], the constant $C_{d,p}$ was defined through some sort of cell problem as

$$\frac{1}{p} G_{d,p} C_{d,p} := \inf \left\{ \liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (0, 1)^d) : u_\delta \rightarrow u_0 \text{ in } L^p((0, 1)^d) \right\},$$

where $u_0(x) = (x_1 + \dots + x_d)/\sqrt{d}$. Unfortunately, this definition is quite implicit and provides no information on the structure of the families that approach the optimal value. This lack of structure complicates things, in such a way that just proving that $C_{d,p} > 0$ requires extremely delicate estimates; this is the content of [Bourgain and Nguyen 2006]. On the Gamma-limsup side, since $\Lambda_{\delta,p}$ is quite sensitive to jumps, what is difficult is gluing together the recovery families corresponding to different slopes, even in the case of a piecewise affine function in dimension 1. This requires a delicate surgery near the junctions; see [Nguyen 2011]. Finally, as for Question 3, difficulties originate from the lack of convexity or continuity of the functionals (1-1), which do not seem to behave well under convolution or similar smoothing techniques.

The core of our approach consists in proving that $\Lambda_{\delta,p}$ in dimension 1 behaves well under vertical δ -segmentation and monotone rearrangement. We refer to Section 3A for the details, but roughly speaking this means that monotone step functions whose values are consecutive integer multiples of δ are the most efficient way to fill the gap between any two given levels. The argument is purely 1-dimensional, and it is carried out in Proposition 3.2. In turn, the proof relies on a discrete combinatorial rearrangement inequality, which we investigate in Theorem 2.2 under more general assumptions.

We observe that this strategy, namely estimating the asymptotic cost of oscillations by reducing ourselves to a discrete combinatorial minimum problem, is the same as that exploited in [Gobbino 1998; Gobbino and Mora 2001], with the remarkable difference that now the reduction to the discrete setting is achieved through vertical δ -segmentation, while in [Gobbino 1998; Gobbino and Mora 2001] it was obtained through a horizontal ε -segmentation (see Figure 1).

The asymptotic estimate on the cost of oscillations opens the door to the Gamma-liminf inequality in dimension 1, which at this point follows from well-established techniques. As for the Gamma-limsup inequality, in dimension 1 we just need to exhibit a family that realizes the given explicit multiple of $\Lambda_{0,p}(u, \mathbb{R})$, and this can be achieved through a vertical δ -segmentation *à la Lebesgue* (see Proposition 3.7). This produces a recovery family made up of step functions, and it is not difficult to modify them in




Figure 1. Vertical δ -segmentation vs. horizontal ε -segmentation (δ is the distance between the parallel lines on the left, ε is the distance between the parallel lines on the right).

order to obtain functions of class C^∞ with asymptotically the same energy (see [Proposition 3.9](#)). Finally, passing from dimension 1 to any dimension is just an application of the 1-dimensional result to all the 1-dimensional sections of a function of d variables.

At the end of the day, we have a completely self-contained proof of [Theorem 1.1](#) above, and a clear indication that the true difficulty of the problem lies in dimension 1, and actually in the discretized combinatorial model. We hope that these ideas can be extended to the more general functionals considered in [\[Brezis and Nguyen 2018\]](#). Some steps in this direction have already been done in [\[Antonucci et al. 2020\]](#); see also [\[Antonucci et al. 2018\]](#).

This paper is organized as follows. In [Section 2](#) we develop a theory of monotone rearrangements, first in a discrete, and then in a semidiscrete setting. In [Section 3](#) we prove our Gamma-convergence result in dimension 1. In [Section 4](#) we prove the Gamma-convergence result in any space dimension.

We would like to thank an anonymous referee for pointing out that the rearrangement inequality in our [Theorem 2.4](#) is equivalent to a rearrangement inequality proved in [\[Garsia and Rodemich 1974\]](#). This equivalence is not immediate (see [Remark 2.5](#) for further details), and for this reason the proofs follow different paths. However, in both cases the basic step consists in reducing the problem to a discrete combinatorial result, namely [Theorem 2.2](#) in this paper, and a variant of Taylor's lemma [\[1973\]](#) in [\[Garsia and Rodemich 1974\]](#).

2. An aggregation/segregation problem

In this section we study the minimum problem for two simplified versions of (1-1), which we interpret as optimizing the disposition of some objects of different types (actually dinosaurs of different species). The first problem is purely discrete, namely with a finite number of dinosaurs of a finite number of species. The second one is semidiscrete, namely with a continuum of dinosaurs belonging to a finite number of species.

2A. Discrete setting. Let us consider

- a positive integer n ,
- a function $u : \{1, \dots, n\} \rightarrow \mathbb{Z}$,
- a symmetric subset $E \subseteq \mathbb{Z}^2$ (namely any subset with the property that $(i, j) \in E$ if and only if $(j, i) \in E$),
- a nonincreasing function $h : \{0, 1, \dots, n-1\} \rightarrow \mathbb{R}$.

Let us introduce the discrete interaction set

$$J(E, u) := \{(x, y) \in \{1, \dots, n\}^2 : x \leq y, (u(x), u(y)) \in E\}, \quad (2-1)$$

and let us finally define

$$\mathcal{H}(h, E, u) := \sum_{(x, y) \in J(E, u)} h(y - x). \quad (2-2)$$

Just to help intuition, we think of u as an arrangement of n dinosaurs placed in the points $\{1, \dots, n\}$. There are different species of dinosaurs, indexed by integer numbers, so that $u(x)$ denotes the species of the dinosaur in position x . The subset $E \subseteq \mathbb{Z}^2$ is the list of all pairs of species that are hostile to each other. A pair of points (x, y) belongs to $J(E, u)$ if and only if $x \leq y$ and the two dinosaurs placed in x and y belong to hostile species, and in this case the real number $h(y - x)$ measures the “hostility” between the two dinosaurs. As expected, the closer the dinosaurs are, the larger their hostility.

Taking this Jurassic framework into account, sometimes in the sequel we call u a “discrete arrangement of n dinosaurs”, we call E an “enemy list”, we call h a “discrete hostility function”, and $\mathcal{H}(h, E, u)$ the “total hostility of the arrangement”. At this level of generality, we admit the possibility that $(i, i) \in E$ for some integer i , namely that a dinosaur is hostile to dinosaurs of the same species, including itself. For this reason, the hostility function $h(x)$ is defined also for $x = 0$. This generality turns out to be useful in the proof of the main result for discrete arrangements.

In the sequel we focus on the special case where E coincides with

$$E_k := \{(i, j) \in \mathbb{Z}^2 : |j - i| \geq k + 1\} \quad (2-3)$$

for some positive integer k . In this case it is quite intuitive that the arrangements that minimize the total hostility are the “monotone” ones, namely those in which all dinosaurs of the same species are close to each other, and the groups corresponding to different species are sorted in ascending or descending order. To this end, we introduce the following notion.

Definition 2.1 (nondecreasing rearrangement: discrete setting). Let n be a positive integer, and let $u : \{1, \dots, n\} \rightarrow \mathbb{Z}$ be a function. The *nondecreasing rearrangement* of u is the function $Mu : \{1, \dots, n\} \rightarrow \mathbb{Z}$ defined as

$$Mu(x) := \min\{j \in \mathbb{Z} : |\{y \in \{1, \dots, n\} : u(y) \leq j\}| \geq x\},$$

where $|A|$ denotes the number of elements of the set A .

As the name suggests, Mu is the unique nondecreasing function that can be represented in the form $Mu = u \circ \pi$, where $\pi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ is a suitable bijection. The nondecreasing rearrangement can also be uniquely characterized by the fact that the two level sets

$$\{x \in \{1, \dots, n\} : u(x) = j\}, \quad \{x \in \{1, \dots, n\} : Mu(x) = j\}$$

have the same number of elements for every $j \in \mathbb{Z}$.

As expected, the main result is that monotone arrangements minimize the total hostility with respect to the enemy list E_k .

Theorem 2.2 (total hostility minimization: discrete setting). *Let n and k be two positive integers, let $E_k \subseteq \mathbb{Z}^2$ be the subset defined by (2-3), and let $h : \{0, \dots, n-1\} \rightarrow \mathbb{R}$ be a nonincreasing function. Let $u : \{1, \dots, n\} \rightarrow \mathbb{Z}$ be any function, let Mu be the nondecreasing rearrangement of u introduced in Definition 2.1, and let $\mathcal{H}(h, E_k, u)$ be the total hostility defined in (2-2).*

Then it turns out that

$$\mathcal{H}(h, E_k, u) \geq \mathcal{H}(h, E_k, Mu). \quad (2-4)$$

Taylor's result [1973] is substantially equivalent to (2-4) in the special case where there are n dinosaurs of n different species indexed by n consecutive integers. It is likely that Taylor's approach based on the celebrated Hall's theorem, sometimes referred to as the "marriage theorem", could work even in the more general setting that we need here; see [Garsia and Rodemich 1974, Section 3]. The proof we present in Section 2C below follows a different path.

2B. Semidiscrete setting. Let us consider

- an interval $(a, b) \subseteq \mathbb{R}$,
- a measurable function $u : (a, b) \rightarrow \mathbb{Z}$ with finite image,
- a symmetric subset $E \subseteq \mathbb{Z}^2$,
- a nonincreasing function $c : (0, b-a) \rightarrow \mathbb{R}$ (note that $c(\sigma)$ might diverge as $\sigma \rightarrow 0^+$).

Let us introduce the semidiscrete interaction set

$$I(E, u) := \{(x, y) \in (a, b)^2 : (u(x), u(y)) \in E\}, \quad (2-5)$$

and let us finally define

$$\mathcal{F}(c, E, u) := \iint_{I(E, u)} c(|y-x|) dx dy. \quad (2-6)$$

In analogy with the discrete setting, we interpret $u(x)$ as a continuous arrangement of dinosaurs of a finite number of species, $c(y-x)$ as the hostility between two dinosaurs of hostile species placed in x and y , and we think of $\mathcal{F}(c, E, u)$ as the total hostility of the arrangement u with respect to the enemy list E .

Once again, we suspect that monotone arrangements minimize the total hostility with respect to the enemy list E_k . This leads to the following notion.

Definition 2.3 (nondecreasing rearrangement: semidiscrete setting). Let $u : (a, b) \rightarrow \mathbb{Z}$ be a measurable function with finite image. The *nondecreasing rearrangement* of u is the function $Mu : (a, b) \rightarrow \mathbb{Z}$ defined as

$$Mu(x) := \min\{j \in \mathbb{Z} : \text{meas}\{y \in (a, b) : u(y) \leq j\} \geq x-a\},$$

where $\text{meas}(A)$ denotes the Lebesgue measure of a subset $A \subseteq (a, b)$.

The function Mu is nondecreasing and satisfies

$$\text{meas}\{x \in (a, b) : u(x) = j\} = \text{meas}\{x \in (a, b) : Mu(x) = j\} \quad \text{for all } j \in \mathbb{Z}.$$

The following result is the semidiscrete counterpart of [Theorem 2.2](#).

Theorem 2.4 (total hostility minimization: semidiscrete setting). *Let $(a, b) \subseteq \mathbb{R}$ be an interval, let k be a positive integer, let $E_k \subseteq \mathbb{Z}^2$ be the subset defined by [\(2-3\)](#), and let $c : (0, b - a) \rightarrow \mathbb{R}$ be a nonincreasing function. Let $u : (a, b) \rightarrow \mathbb{Z}$ be any measurable function with finite image, let Mu be the nondecreasing rearrangement of u introduced in [Definition 2.3](#), and let $\mathcal{F}(c, E_k, u)$ be the total hostility defined in [\(2-6\)](#).*

Then it turns out that

$$\mathcal{F}(c, E_k, u) \geq \mathcal{F}(c, E_k, Mu). \tag{2-7}$$

Remark 2.5. [Theorem 2.4](#) above is stated in the form that we need in the proof of [Proposition 3.2](#). With a further approximation step in the proof, one can show that the same conclusion [\(2-7\)](#) holds true also without assuming that the image of u is finite and contained in \mathbb{Z} , and without assuming that k is a positive integer (but just a real number greater than -1).

It is interesting to compare this extended result with [\[Garsia and Rodemich 1974, Theorem 1.1\]](#), which states that for every nondecreasing function $\Phi : [0, +\infty) \rightarrow [0, +\infty)$, and every $t \in (0, b - a)$, it turns out that

$$\int_{D(t)} \Phi(|u(y) - u(x)|) dx dy \geq \int_{D(t)} \Phi(|Mu(y) - Mu(x)|) dx dy, \tag{2-8}$$

where $D(t) := \{(x, y) \in (a, b)^2 : |y - x| \leq t\}$. We observe that in [\(2-8\)](#) the integral involves only the pairs $(x, y) \in (a, b)^2$ that are close enough to the diagonal $y = x$, and the integrand Φ penalizes the pairs for which $|u(y) - u(x)|$ is large. On the contrary, in our total hostility the integral involves only the pairs with $|u(y) - u(x)|$ large enough, and the integrand c penalizes the pairs that are close to the diagonal. In this sense the two statements seem to be two sides of the same coin (again as the Riemann and the Lebesgue integral), and actually one can show that both statements are equivalent to saying that the inequality

$$\begin{aligned} \text{meas}\{(x, y) \in (a, b)^2 : |y - x| \leq t, |u(y) - u(x)| \geq \delta\} \\ \geq \text{meas}\{(x, y) \in (a, b)^2 : |y - x| \leq t, |Mu(y) - Mu(x)| \geq \delta\} \end{aligned} \tag{2-9}$$

holds true for every $t \in (0, b - a)$ and every $\delta > 0$.

The proof of [\(2-8\)](#) given in [\[Garsia and Rodemich 1974\]](#) relies on this equivalence, and establishes [\(2-9\)](#) through a variant of Taylor’s result. The proof of [\(2-7\)](#) that we present in [Section 2D](#) follows a more direct path, based on our [Theorem 2.2](#), which anyway is again discrete combinatorics.

2C. Proof of [Theorem 2.2](#). Since the hostility function h is fixed, in the sequel we simply write $\mathcal{H}(E, u)$ instead of $\mathcal{H}(h, E, u)$.

Our idea is to proceed by induction on the number of dinosaurs. In the case $n = 1$ there is nothing to prove. Let us assume now that [\(2-4\)](#) holds true for all arrangements of n dinosaurs, and let u be any arrangement of $n + 1$ dinosaurs. In order to obtain an arrangement of n dinosaurs, we remove from u the rightmost dinosaur of the species indexed by the highest integer, and we shift one position to the left all subsequent dinosaurs. More formally, we set

$$\mu := \max\{u(i) : i \in \{1, \dots, n + 1\}\},$$

we consider the largest index $m \in \{1, \dots, n+1\}$ such that $u(m) = \mu$, and we define the *reduction* of u to be the new arrangement $\text{Red}(u) : \{1, \dots, n\} \rightarrow \mathbb{Z}$ given by

$$[\text{Red}(u)](i) := \begin{cases} u(i) & \text{if } i < m, \\ u(i+1) & \text{if } i \geq m. \end{cases}$$

When passing from u to $\text{Red}(u)$, the total hostility changes by an amount that we call the *hostility gap*, defined as

$$\Delta(E, u) := \mathcal{H}(E, u) - \mathcal{H}(E, \text{Red}(u)).$$

Since $M(\text{Red}(u)) = \text{Red}(Mu)$, the inductive hypothesis reads as

$$\mathcal{H}(E_k, \text{Red}(u)) \geq \mathcal{H}(E_k, M(\text{Red}(u))) = \mathcal{H}(E_k, \text{Red}(Mu)),$$

and therefore

$$\begin{aligned} \mathcal{H}(E_k, u) &= \mathcal{H}(E_k, \text{Red}(u)) + \Delta(E_k, u) \\ &\geq \mathcal{H}(E_k, \text{Red}(Mu)) + \Delta(E_k, u) \\ &= \mathcal{H}(E_k, Mu) - \Delta(E_k, Mu) + \Delta(E_k, u). \end{aligned}$$

As a consequence, (2-4) is proved for the arrangement u if we can show that

$$\Delta(E_k, u) \geq \Delta(E_k, Mu), \quad (2-10)$$

namely that the monotone rearrangement decreases (or at least does not increase) the hostility gap.

In order to prove (2-10), we begin by deriving a formula for the hostility gap. Let us consider the removal that leads from u to $\text{Red}(u)$. We observe that interactions between any two dinosaurs placed on the same side of the removed one are equal before and after the removal, and therefore they cancel out when computing the gap. On the contrary, if two hostile dinosaurs are placed within distance d on opposite sides of the removed one, their hostility changes from $h(d)$ to $h(d-1)$ after the removal. It follows that the hostility gap can be written as

$$\Delta(E, u) = \sum_{i \in J_1(E, u, m)} h(|m-i|) - \sum_{(i,j) \in J_2(E, u, m)} (h(j-i-1) - h(j-i)), \quad (2-11)$$

where

$$J_1(E, u, m) := \{i \in \{1, \dots, n+1\} : (u(i), u(m)) \in E\},$$

$$J_2(E, u, m) := \{(i, j) \in \{1, \dots, n+1\}^2 : i < m < j, (u(i), u(j)) \in E\}.$$

The first sum in (2-11) takes into account the interactions of the removed dinosaur with the rest of the world, and the second sum represents the increment of the total hostility due to the reduction of distances among the others.

Now we introduce the new enemy list

$$E_{\langle \mu \rangle} := \mathbb{Z}^2 \setminus \{\mu, \mu-1, \dots, \mu-k\}^2,$$

and we claim that

$$\Delta(E_k, u) \geq \Delta(E_{\langle \mu \rangle}, u) \geq \Delta(E_{\langle \mu \rangle}, Mu) = \Delta(E_k, Mu), \quad (2-12)$$

which of course implies (2-10).

The equality between the last two terms of (2-12) follows from formula (2-11). Indeed, since Mu is nondecreasing, the removed dinosaur is the rightmost one, and therefore in both cases the second sum in (2-11) is void. Also the first sum in (2-11) is the same in both cases, because a dinosaur of the highest species is hostile to another dinosaur with respect to the enemy list E_k if and only if it is hostile to the same dinosaur with respect to the enemy list $E_{\langle\mu\rangle}$.

The inequality between the first two terms of (2-12) follows again from formula (2-11). Indeed, the first sum has the same terms both in the case of the enemy list E_k and in the case of the enemy list $E_{\langle\mu\rangle}$, as observed above. As for the second sum, the interactions with respect to E_k are also interactions with respect to $E_{\langle\mu\rangle}$, and therefore when passing from E_k to $E_{\langle\mu\rangle}$ the second sum cannot decrease. Since the second sum appears in (2-11) with negative sign, the hostility gap with respect to $E_{\langle\mu\rangle}$ is less than or equal to the hostility gap with respect to E_k .

It remains to prove that

$$\Delta(E_{\langle\mu\rangle}, u) \geq \Delta(E_{\langle\mu\rangle}, Mu). \tag{2-13}$$

To this end, we introduce the complement enemy list

$$E_{\langle\mu\rangle}^c := \{\mu, \mu - 1, \dots, \mu - k\}^2 = \mathbb{Z}^2 \setminus E_{\langle\mu\rangle}.$$

Since \mathbb{Z}^2 is the disjoint union of $E_{\langle\mu\rangle}$ and $E_{\langle\mu\rangle}^c$, and the total hostility is additive with respect to the enemy list, we deduce

$$\mathcal{H}(E_{\langle\mu\rangle}, w) = \mathcal{H}(\mathbb{Z}^2, w) - \mathcal{H}(E_{\langle\mu\rangle}^c, w)$$

for every arrangement w , and for the same reason

$$\Delta(E_{\langle\mu\rangle}, w) = \Delta(\mathbb{Z}^2, w) - \Delta(E_{\langle\mu\rangle}^c, w).$$

Moreover, we observe that the total hostility with respect to \mathbb{Z}^2 depends only on the number of dinosaurs, and in particular

$$\Delta(\mathbb{Z}^2, u) = \Delta(\mathbb{Z}^2, Mu).$$

As a consequence, proving (2-13) is equivalent to showing that

$$\Delta(E_{\langle\mu\rangle}^c, u) \leq \Delta(E_{\langle\mu\rangle}^c, Mu). \tag{2-14}$$

The advantage of this “complement formulation” is that hostility gaps with respect to $E_{\langle\mu\rangle}^c$ depend only on the relative positions of the removed dinosaur with respect to the other dinosaurs of the species with indices between $\mu - k$ and μ .

To be more precise, let us compute the left-hand side of (2-14). Let m denote as usual the position of the dinosaur that is removed from u to $\text{Red}(u)$, and let us set

$$\begin{aligned} R(u) &:= \{r \geq 1 : u(m+r) \in \{\mu, \mu - 1, \dots, \mu - k\}\}, \\ L(u) &:= \{\ell \geq 1 : u(m-\ell) \in \{\mu, \mu - 1, \dots, \mu - k\}\}. \end{aligned}$$

In other words, this means that

$$\{m - \ell : \ell \in L(u)\} \cup \{m\} \cup \{m + r : r \in R(u)\}$$

is the set of all integers $i \in \{1, \dots, n+1\}$ such that $u(i) \in \{\mu, \mu-1, \dots, \mu-k\}$, namely the set of positions where the dinosaurs of the last $k+1$ species are placed. With this notation, the first sum in (2-11) is

$$h(0) + \sum_{\ell \in L(u)} h(\ell) + \sum_{r \in R(u)} h(r)$$

(we recall that in this “complement formulation” the dinosaur in position m is also hostile to itself), while the second sum in (2-11) is

$$\sum_{(\ell, r) \in L(u) \times R(u)} (h(\ell+r-1) - h(\ell+r)).$$

Therefore, it turns out that

$$\Delta(E_{(\mu)}^c, u) = \mathcal{G}(L(u), R(u)),$$

where the function \mathcal{G} is defined by

$$\mathcal{G}(L, R) := h(0) + \sum_{\ell \in L} h(\ell) + \sum_{r \in R} h(r) - \sum_{(\ell, r) \in L \times R} (h(\ell+r-1) - h(\ell+r)) \quad (2-15)$$

for any two sets L and R of positive integers.

On the other hand, in the nondecreasing arrangement Mu the rightmost dinosaur has $|L(u)| + |R(u)|$ dinosaurs of the last $k+1$ species exactly on its left, and therefore

$$\Delta(E_{(\mu)}^c, Mu) = \sum_{i=0}^{|L(u)|+|R(u)|} h(i).$$

As a consequence, inequality (2-14) is proved if we show that

$$\mathcal{G}(L, R) \leq \sum_{i=0}^{|L|+|R|} h(i) \quad (2-16)$$

for every choice of the sets L and R . For this final step, we argue by induction on the number of elements of R . If $R = \emptyset$, from (2-15) we deduce

$$\mathcal{G}(L, R) := h(0) + \sum_{\ell \in L} h(\ell) \leq \sum_{i=0}^{|L|} h(i) = \sum_{i=0}^{|L|+|R|} h(i),$$

where the inequality is true term-by-term because h is nonincreasing.

Let us assume now that the conclusion holds true whenever R has n elements, and let us consider any pair (L, R) with $|R| = n+1$. Let us set

$$a := \max R, \quad b := \min\{n \in \mathbb{N} \setminus \{0\} : n \notin L\},$$

and let us consider the new pair (L_1, R_1) defined as

$$L_1 := L \cup \{b\}, \quad R_1 := R \setminus \{a\}.$$

In words, we have removed the largest element of R , and added the smallest possible element to L . We observe that $|R_1| = n$ and $|L_1| + |R_1| = |L| + |R|$. Therefore, if we show that

$$\mathcal{G}(L, R) \leq \mathcal{G}(L_1, R_1), \quad (2-17)$$

then (2-16) follows from the inductive assumption.

In order to prove (2-17), we expand the left- and right-hand sides according to (2-15). After canceling out the common terms, with some algebra we obtain that inequality (2-17) holds true if and only if

$$h(a) + \sum_{r \in R_1} (h(b+r-1) - h(b+r)) \leq h(b) + \sum_{\ell \in L} (h(\ell+a-1) - h(\ell+a)). \quad (2-18)$$

All terms in the sums are nonnegative because h is nonincreasing. Let us consider the left-hand side. If $a > 1$ we know that $R_1 \subseteq \{1, \dots, a-1\}$, and hence

$$\begin{aligned} h(a) + \sum_{r \in R_1} (h(b+r-1) - h(b+r)) &\leq h(a) + \sum_{r=1}^{a-1} (h(b+r-1) - h(b+r)) \\ &= h(a) + h(b) - h(a+b-1). \end{aligned} \quad (2-19)$$

The same inequality is true for trivial reasons also if $a = 1$.

Let us consider now the right-hand side of (2-18). If $b > 1$ we know that $L \supseteq \{1, \dots, b-1\}$, and hence

$$\begin{aligned} h(b) + \sum_{\ell \in L} (h(\ell+a-1) - h(\ell+a)) &\geq h(b) + \sum_{\ell=1}^{b-1} (h(\ell+a-1) - h(\ell+a)) \\ &= h(b) + h(a) - h(a+b-1). \end{aligned} \quad (2-20)$$

As before, the same inequality is true for trivial reasons also if $b = 1$.

Combining (2-20) and (2-19) we obtain (2-18), which in turn is equivalent to (2-17). This completes the proof of (2-16). \square

2D. Proof of Theorem 2.4. The proof relies on the following approximation result (we omit the proof, which is an exercise in basic measure theory).

Lemma 2.6. *Let m be a positive integer, and let D_1, \dots, D_m be disjoint measurable subsets of $(0, 1)$ such that*

$$\bigcup_{i=1}^m D_i = (0, 1).$$

Then for every $\varepsilon > 0$ there exist disjoint subsets $D_{1,\varepsilon}, \dots, D_{m,\varepsilon}$ of $[0, 1]$ such that

$$\bigcup_{i=1}^m D_{i,\varepsilon} = (0, 1)$$

and such that for every $i = 1, \dots, m$ it turns out that

- $D_{i,\varepsilon}$ is a finite union of intervals with rational endpoints,
- the Lebesgue measure of the symmetric difference between D_i and $D_{i,\varepsilon}$ is less than or equal to ε .

We are now ready to prove [Theorem 2.4](#). First of all, we observe that (2-7) is invariant by translations and homotheties. As a consequence, there is no loss of generality in assuming that $(a, b) = (0, 1)$ and $c : (0, 1) \rightarrow \mathbb{R}$. Then we proceed in three steps. To begin with, we prove (2-7) in the special case where the hostility function c is bounded and the arrangement u has a very rigid structure, then for general u but again bounded hostility function, and finally in the general setting.

Step 1: We prove (2-7) under the additional assumption that the hostility function $c : (0, 1) \rightarrow \mathbb{R}$ is bounded, and that there exists a positive integer d such that $u(x)$ is constant in each interval of the form $((i-1)/d, i/d)$ with $i = 1, \dots, d$.

Indeed, this is actually the discrete setting. To be more precise, we introduce the discrete arrangement $v : \{1, \dots, d\} \rightarrow \mathbb{Z}$ defined as

$$v(i) := u\left(\frac{i - \frac{1}{2}}{d}\right) \quad \text{for all } i \in \{1, \dots, d\}$$

and the discrete hostility function $h : \{0, \dots, d-1\} \rightarrow \mathbb{R}$ defined as

$$h(i) := \int_0^{1/d} dx \int_{i/d}^{(i+1)/d} c(|y-x|) dy \quad \text{for all } i \in \{0, \dots, d-1\},$$

which represents the contribution to the total hostility of two intervals of length $1/d$ occupied by hostile dinosaurs, and placed at distance i/d from each other. Then for every enemy list E_k it turns out that

$$\mathcal{F}(c, E_k, u) = 2\mathcal{H}(h, E_k, v),$$

where $\mathcal{H}(h, E_k, v)$ is the discrete total hostility defined in (2-2), and the factor 2 takes into account that both (x, y) and (y, x) are included in the semidiscrete interaction set $I(E_k, u)$, while only one of them is included in the discrete counterpart $J(E_k, v)$; see (2-1) and (2-5). Moreover, the monotone rearrangement Mv of v is related to the monotone rearrangement Mu of u by the formula

$$Mv(i) = Mu\left(\frac{i - \frac{1}{2}}{d}\right) \quad \text{for all } i \in \{1, \dots, d\},$$

and again it turns out that

$$\mathcal{F}(c, E_k, Mu) = 2\mathcal{H}(h, E_k, Mv)$$

for every enemy list E_k . At this point, (2-7) is equivalent to

$$\mathcal{H}(h, E_k, v) \geq \mathcal{H}(h, E_k, Mv),$$

which in turn is true because of [Theorem 2.2](#).

Step 2: We prove (2-7) for a general arrangement $u : (0, 1) \rightarrow \mathbb{Z}$, but again under the additional assumption that the hostility function $c : (0, 1) \rightarrow \mathbb{R}$ is bounded.

To this end, let $z_1 < z_2 < \dots < z_m$ denote the elements in the image of u , and let

$$D_i := \{x \in (0, 1) : u(x) = z_i\} \quad \text{for all } i \in \{1, \dots, m\}$$

denote the set of positions of dinosaurs of the species z_i . For every $\varepsilon > 0$, let us consider the sets $D_{1,\varepsilon}, \dots, D_{m,\varepsilon}$ given by [Lemma 2.6](#), and the function $u_\varepsilon : (0, 1) \rightarrow \mathbb{Z}$ defined as

$$u_\varepsilon(x) = z_i \quad \text{for all } x \in D_{i,\varepsilon}.$$

Since the hostility function c is bounded, and the symmetric difference between D_i and $D_{i,\varepsilon}$ has measure less than or equal to ε , there exists a constant Γ (depending on m and c , but independent of ε) such that

$$|\mathcal{F}(c, E_k, u) - \mathcal{F}(c, E_k, u_\varepsilon)| \leq \Gamma\varepsilon \quad \text{and} \quad |\mathcal{F}(c, E_k, Mu) - \mathcal{F}(c, E_k, Mu_\varepsilon)| \leq \Gamma\varepsilon.$$

On the other hand, the function u_ε satisfies the assumptions of the previous step, and therefore

$$\mathcal{F}(c, E_k, u_\varepsilon) \geq \mathcal{F}(c, E_k, Mu_\varepsilon).$$

From all these inequalities it follows that

$$\mathcal{F}(c, E_k, u) \geq \mathcal{F}(c, E_k, Mu) - 2\Gamma\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, [\(2-7\)](#) is proved in this case.

Step 3: We prove [\(2-7\)](#) without assuming that the hostility function $c(x)$ is bounded.

To this end, for every $n \in \mathbb{N}$ we consider the truncated hostility function

$$c_n(x) := \min\{c(x), n\} \quad \text{for all } x \in (0, 1).$$

We observe that

$$\mathcal{F}(c, E_k, u) \geq \mathcal{F}(c_n, E_k, u) \quad \text{for all } n \in \mathbb{N}$$

because $c(x) \geq c_n(x)$ for every $x \in (0, 1)$, and

$$\mathcal{F}(c_n, E_k, u) \geq \mathcal{F}(c_n, E_k, Mu) \quad \text{for all } n \in \mathbb{N}$$

because of the result of the previous step applied to the bounded hostility function $c_n(x)$. As a consequence, we obtain

$$\mathcal{F}(c, E_k, u) \geq \mathcal{F}(c_n, E_k, Mu) \quad \text{for all } n \in \mathbb{N}. \tag{2-21}$$

On the other hand, by monotone convergence we deduce

$$\mathcal{F}(c, E_k, Mu) = \sup_{n \in \mathbb{N}} \mathcal{F}(c_n, E_k, Mu),$$

and therefore [\(2-7\)](#) follows from [\(2-21\)](#). □

3. Gamma-convergence in dimension 1

In this section we prove [Theorem 1.1](#) for $d = 1$, in which case

$$G_{1,p} = 2 \quad \text{for all } p \geq 1. \tag{3-1}$$

To begin with, we introduce the notion of vertical δ -segmentation, which is going to play a crucial role in many parts of the proof.

Definition 3.1 (vertical δ -segmentation). Let \mathbb{X} be any set, let $w : \mathbb{X} \rightarrow \mathbb{R}$ be any function, and let $\delta > 0$. The vertical δ -segmentation of w is the function $S_\delta w : \mathbb{X} \rightarrow \mathbb{R}$ defined by

$$S_\delta w(x) := \delta \left\lfloor \frac{w(x)}{\delta} \right\rfloor \quad \text{for all } x \in \mathbb{X}. \quad (3-2)$$

The function $S_\delta w$ takes its values in $\delta\mathbb{Z}$, and it is uniquely characterized by the fact that $S_\delta w(x) = k\delta$ for some $k \in \mathbb{Z}$ if and only if $k\delta \leq w(x) < (k+1)\delta$.

3A. Asymptotic cost of oscillations. Let us assume that a function $u_\delta(x)$ oscillates between two values A and B in some interval (a, b) . Does this provide an estimate from below for $\Lambda_{\delta,p}(u_\delta, (a, b))$, at least when δ is small enough? The following proposition and the subsequent corollaries give a sharp quantitative answer to this question. They are the fundamental tool in the proof of the liminf inequality.

Proposition 3.2 (limit cost of vertical oscillations). *Let $p \geq 1$ be a real number, let $(a, b) \subseteq \mathbb{R}$ be an interval, and let $\{u_\delta\}_{\delta>0} \subseteq L^p((a, b))$ be a family of functions.*

Let us assume that there exist two real numbers $A \leq B$ such that

$$\liminf_{\delta \rightarrow 0^+} \text{meas}\{x \in (a, b) : u_\delta(x) \leq A + \varepsilon\} > 0 \quad \text{for all } \varepsilon > 0, \quad (3-3)$$

$$\liminf_{\delta \rightarrow 0^+} \text{meas}\{x \in (a, b) : u_\delta(x) \geq B - \varepsilon\} > 0 \quad \text{for all } \varepsilon > 0. \quad (3-4)$$

Then it turns out that

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (a, b)) \geq \frac{2}{p} \cdot C_p \cdot \frac{(B-A)^p}{(b-a)^{p-1}}, \quad (3-5)$$

where C_p is the constant defined in (1-6).

Proof. To begin with, we observe that (3-5) is trivial if $A = B$, and therefore in the sequel we assume that $A < B$.

Let us fix $\varepsilon > 0$ such that $4\varepsilon < B - A$. Due to assumptions (3-3) and (3-4), there exist $\eta > 0$ and $\delta_0 > 0$ such that

$$\text{meas}\{x \in (a, b) : u_\delta(x) \leq A + \varepsilon\} \geq \eta \quad \text{for all } \delta \in (0, \delta_0), \quad (3-6)$$

$$\text{meas}\{x \in (a, b) : u_\delta(x) \geq B - \varepsilon\} \geq \eta \quad \text{for all } \delta \in (0, \delta_0). \quad (3-7)$$

Truncation, δ -segmentation and monotone rearrangement: In this section of the proof, we replace $\{u_\delta\}$ with a new family $\{\hat{u}_\delta\}$ of monotone piecewise constant functions that still satisfies (3-3) and (3-4), without increasing the left-hand side of (3-5). To this end, we perform three operations on $u_\delta(x)$.

The first operation is a truncation between A and B . To be more precise, we define $T_{A,B}u_\delta : (a, b) \rightarrow \mathbb{R}$ by setting

$$T_{A,B}u_\delta(x) := \begin{cases} A & \text{if } u_\delta(x) < A, \\ u_\delta(x) & \text{if } A \leq u_\delta(x) \leq B, \\ B & \text{if } u_\delta(x) > B. \end{cases}$$

We observe that the implication

$$|T_{A,Bu_\delta}(y) - T_{A,Bu_\delta}(x)| > \delta \implies |u_\delta(y) - u_\delta(x)| > \delta$$

holds true for every x and y in (a, b) , and hence

$$\Lambda_{\delta,p}(T_{A,Bu_\delta}, (a, b)) \leq \Lambda_{\delta,p}(u_\delta, (a, b)) \quad \text{for all } \delta > 0.$$

We also observe that (3-6) and (3-7) remain true if we replace $u_\delta(x)$ by $T_{A,Bu_\delta}(x)$.

The second operation is a vertical δ -segmentation; namely we replace T_{A,Bu_δ} by the function $S_\delta T_{A,Bu_\delta}$ defined according to (3-2). Again we observe that the implications

$$\begin{aligned} |S_\delta T_{A,Bu_\delta}(y) - S_\delta T_{A,Bu_\delta}(x)| > \delta &\implies |S_\delta T_{A,Bu_\delta}(y) - S_\delta T_{A,Bu_\delta}(x)| \geq 2\delta \\ &\implies |T_{A,Bu_\delta}(y) - T_{A,Bu_\delta}(x)| > \delta \end{aligned}$$

hold true for every x and y in (a, b) , and hence

$$\Lambda_{\delta,p}(S_\delta T_{A,Bu_\delta}, (a, b)) \leq \Lambda_{\delta,p}(T_{A,Bu_\delta}, (a, b)) \quad \text{for all } \delta > 0.$$

As for (3-6) and (3-7), we set $\delta_1 := \min\{\varepsilon, \delta_0\}$, and we observe that now

$$\text{meas}\{x \in (a, b) : S_\delta T_{A,Bu_\delta}(x) \leq A + 2\varepsilon\} \geq \eta \quad \text{for all } \delta \in (0, \delta_1), \quad (3-8)$$

$$\text{meas}\{x \in (a, b) : S_\delta T_{A,Bu_\delta}(x) \geq B - 2\varepsilon\} \geq \eta \quad \text{for all } \delta \in (0, \delta_1). \quad (3-9)$$

The third and last operation we perform is monotone rearrangement; namely we replace $S_\delta T_{A,Bu_\delta}$ with the nondecreasing function $MS_\delta T_{A,Bu_\delta}$ in (a, b) whose level sets have the same measure of the level sets of $S_\delta T_{A,Bu_\delta}$ (see Definition 2.3).

From (3-8) and (3-9) we deduce that now

$$MS_\delta T_{A,Bu_\delta}(x) \leq A + 2\varepsilon \quad \text{for all } x \in (a, a + \eta), \quad \text{for all } \delta \in (0, \delta_1), \quad (3-10)$$

$$MS_\delta T_{A,Bu_\delta}(x) \geq B - 2\varepsilon \quad \text{for all } x \in (b - \eta, b), \quad \text{for all } \delta \in (0, \delta_1). \quad (3-11)$$

Moreover, we claim that

$$\Lambda_{\delta,p}(MS_\delta T_{A,Bu_\delta}, (a, b)) \leq \Lambda_{\delta,p}(S_\delta T_{A,Bu_\delta}, (a, b)) \quad \text{for all } \delta > 0. \quad (3-12)$$

This is a straightforward consequence of Theorem 2.4. To be more formal, let us consider the semidiscrete arrangement $v_\delta : (a, b) \rightarrow \mathbb{Z}$ defined by

$$v_\delta(x) := \frac{1}{\delta} S_\delta T_{A,Bu_\delta}(x) \quad \text{for all } x \in (a, b)$$

(we recall that $S_\delta T_{A,Bu_\delta}$ takes its values in $\delta\mathbb{Z}$, and hence $v_\delta(x)$ is integer-valued) and the hostility function $c : (0, b - a) \rightarrow \mathbb{R}$ defined as $c(\sigma) := \delta^p \sigma^{-1-p}$. We observe that

$$MS_\delta T_{A,Bu_\delta}(x) = \delta Mv_\delta(x) \quad \text{for all } x \in (a, b),$$

where Mv_δ is the nondecreasing rearrangement of v_δ according to Definition 2.3.

We observe also that for every pair of points x and y in (a, b) it turns out that

$$(x, y) \in I(\delta, S_\delta T_{A,B} u_\delta, (a, b)) \iff |v_\delta(y) - v_\delta(x)| \geq 2 \iff (x, y) \in I(E_1, v_\delta),$$

where E_1 is the enemy list defined in (2-3), and $I(E_1, v_\delta)$ is the semidiscrete interaction set defined according to (2-5). It follows that

$$\Lambda_{\delta,p}(S_\delta T_{A,B} u_\delta, (a, b)) = \mathcal{F}(c, E_1, v_\delta), \quad \Lambda_{\delta,p}(M S_\delta T_{A,B} u_\delta, (a, b)) = \mathcal{F}(c, E_1, M v_\delta),$$

and therefore (3-12) is equivalent to (2-7).

In conclusion, the three operations described so far delivered us a family

$$\hat{u}_\delta := M S_\delta T_{A,B} u_\delta$$

of nondecreasing functions such that the image of \hat{u}_δ is contained in $\delta\mathbb{Z}$. This family satisfies (3-10) and (3-11), and

$$\Lambda_{\delta,p}(u_\delta, (a, b)) \geq \Lambda_{\delta,p}(\hat{u}_\delta, (a, b)) \quad \text{for all } \delta > 0. \quad (3-13)$$

In the sequel we are going to show that any such family satisfies

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(\hat{u}_\delta, (a, b)) \geq \frac{2}{p} \cdot C_p \cdot \frac{(B - A - 4\varepsilon)^p}{(b - a)^{p-1}}. \quad (3-14)$$

Due to (3-13) and the arbitrariness of $\varepsilon > 0$, this is enough to prove (3-5).

Extension of the integrals to a vertical strip: In this section of the proof we modify the domain of integration in order to simplify the computation of $\Lambda_{\delta,p}(\hat{u}_\delta, (a, b))$. To begin with, we observe that

$$\Lambda_{\delta,p}(\hat{u}_\delta, (a, b)) = \iint_{A_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy \geq \iint_{B_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy,$$

where

$$\begin{aligned} A_\delta &:= I(\delta, \hat{u}_\delta, (a, b)) = \{(x, y) \in (a, b)^2 : |\hat{u}_\delta(y) - \hat{u}_\delta(x)| > \delta\}, \\ B_\delta &:= \{(x, y) \in (a + \eta, b - \eta) \times (a, b) : |\hat{u}_\delta(y) - \hat{u}_\delta(x)| > \delta\}. \end{aligned}$$

Then we write the last integral in the form

$$\iint_{B_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy = \iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy - \iint_{C_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy,$$

where

$$C_\delta := (a + \eta, b - \eta) \times (\mathbb{R} \setminus (a, b)).$$

In other words, the set $B_\delta \cup C_\delta$ consists of the vertical strip $(a + \eta, b - \eta) \times \mathbb{R}$ minus the set of points $(x, y) \in (a + \eta, b - \eta) \times (a, b)$ such that $|\hat{u}_\delta(y) - \hat{u}_\delta(x)| \leq \delta$. Now we observe that

$$\iint_{C_\delta} \frac{\delta^p}{|y - x|^{1+p}} dx dy = 2\delta^p \int_{a+\eta}^{b-\eta} dx \int_b^{+\infty} \frac{1}{|y - x|^{1+p}} dy.$$

From the convergence of the last double integral it follows that

$$\lim_{\delta \rightarrow 0^+} \iint_{C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy = 0,$$

and therefore

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(\hat{u}_\delta, (a, b)) \geq \liminf_{\delta \rightarrow 0^+} \iint_{B_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy = \liminf_{\delta \rightarrow 0^+} \iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy. \quad (3-15)$$

Computing the integrals: In this last part of the proof we show that

$$\liminf_{\delta \rightarrow 0^+} \iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy \geq \frac{2}{p} \cdot C_p \cdot \frac{(B-A-4\varepsilon)^p}{(b-a)^{p-1}}. \quad (3-16)$$

Recalling (3-15), this proves (3-14), and hence also (3-5).

To this end, we need to introduce some notation. We know that \hat{u}_δ is a nondecreasing function with finite image. Let us consider the partition

$$a = x_0 < x_1 < \dots < x_n = b$$

of (a, b) with the property that $\hat{u}_\delta(x)$ is constant in each interval of the form (x_{i-1}, x_i) , and different intervals correspond to different constants. Let us set

$$\begin{aligned} h &:= \min\{i \in \{1, \dots, n\} : x_i \geq a + \eta\}, \\ k &:= \max\{i \in \{0, \dots, n-1\} : x_i \leq b - \eta\}. \end{aligned}$$

Of course n, h, k , as well as the partition, do depend on δ . Now we claim that

$$\iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy \geq \frac{2}{p} \cdot C_p \cdot \frac{\delta^p (k-h-1)^p}{(b-a)^{p-1}} \quad \text{for all } \delta \in (0, \delta_1). \quad (3-17)$$

To this end we can limit ourselves, without loss of generality, to the case where the values of $\hat{u}_\delta(x)$ in neighboring intervals are consecutive multiples of δ ; namely if $\hat{u}_\delta(x) = m\delta$ in (x_{i-1}, x_i) for some $m \in \mathbb{Z}$, then $\hat{u}_\delta(x) = (m+1)\delta$ in (x_i, x_{i+1}) . Indeed, if $\hat{u}_\delta(x) \geq (m+2)\delta$ in (x_i, x_{i+1}) , then it turns out that

$$\iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy \geq \int_{x_{i-1}}^{x_i} dx \int_{x_i}^{x_{i+1}} \frac{\delta^p}{(y-x)^{1+p}} dy.$$

Since the integral in the right-hand side is divergent, the left-hand side is divergent as well, and in this case (3-17) is trivially true.

Therefore, in the sequel we treat the case where the values of $\hat{u}_\delta(x)$ in neighboring intervals are consecutive multiples of δ . Under this assumption it turns out that

$$\begin{aligned} \iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy &\geq \sum_{i=h+1}^{k-1} \left(\int_{x_{i-1}}^{x_i} dx \int_{x_{i+1}}^{+\infty} \frac{\delta^p}{|y-x|^{1+p}} dy + \int_{x_i}^{x_{i+1}} dx \int_{-\infty}^{x_{i-1}} \frac{\delta^p}{|y-x|^{1+p}} dy \right) \\ &= \frac{\delta^p}{p} \sum_{i=h+1}^{k-1} \left(\int_{x_{i-1}}^{x_i} \frac{1}{(x_{i+1}-x)^p} dx + \int_{x_i}^{x_{i+1}} \frac{1}{(x-x_{i-1})^p} dx \right). \end{aligned}$$

Now we distinguish two cases.

- If $p = 1$, computing the integrals we obtain

$$\iint_{B_\delta \cup C_\delta} \frac{\delta}{(y-x)^2} dx dy \geq \delta \sum_{i=h+1}^{k-1} \log \left(\frac{x_{i+1} - x_{i-1}}{x_{i+1} - x_i} \cdot \frac{x_{i+1} - x_{i-1}}{x_i - x_{i-1}} \right).$$

If $\ell_i := x_i - x_{i-1}$ denotes the length of the i -th interval of the partition, and we apply the inequality between arithmetic and geometric mean, we obtain

$$\iint_{B_\delta \cup C_\delta} \frac{\delta}{(y-x)^2} dx dy \geq \delta \sum_{i=h+1}^{k-1} \log \frac{(\ell_i + \ell_{i+1})^2}{\ell_i \cdot \ell_{i+1}} \geq \delta \sum_{i=h+1}^{k-1} \log 4 = 2 \log 2 \cdot \delta(k-h-1),$$

which proves (3-17) in this case.

- If $p > 1$, computing the integrals we obtain

$$\iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy \geq \frac{\delta^p}{p(p-1)} \sum_{i=h+1}^{k-1} \left(\frac{1}{\ell_{i+1}^{p-1}} + \frac{1}{\ell_i^{p-1}} - \frac{2}{(\ell_{i+1} + \ell_i)^{p-1}} \right),$$

where we set $\ell_i := x_i - x_{i-1}$ as before. Therefore, with two applications of Jensen's inequality to the convex function $t \rightarrow t^{1-p}$, we obtain

$$\begin{aligned} \iint_{B_\delta \cup C_\delta} \frac{\delta^p}{|y-x|^{1+p}} dx dy &\geq \frac{\delta^p}{p(p-1)} \sum_{i=h+1}^{k-1} \frac{2^p - 2}{(\ell_{i+1} + \ell_i)^{p-1}} \\ &\geq \frac{\delta^p (2^p - 2)}{p(p-1)} \cdot \frac{(k-h-1)^p}{\left(\sum_{i=h+1}^{k-1} (\ell_{i+1} + \ell_i) \right)^{p-1}} \\ &\geq \frac{\delta^p (2^p - 2)}{p(p-1)} \cdot \frac{(k-h-1)^p}{(2(b-a))^{p-1}} = \frac{2}{p} \cdot C_p \cdot \frac{\delta^p (k-h-1)^p}{(b-a)^{p-1}}, \end{aligned}$$

which proves (3-17) also in this case.

Now it remains to estimate $\delta(k-h-1)$. To this end, from (3-10) and the minimality of h we deduce

$$A + 2\varepsilon \geq \hat{u}_\delta(x) =: m_A \delta \quad \text{for all } x \in (x_{h-1}, x_h).$$

Similarly, from (3-11) and the maximality of k we deduce

$$B - 2\varepsilon \leq \hat{u}_\delta(x) =: m_B \delta \quad \text{for all } x \in (x_k, x_{k+1}).$$

Since the values of \hat{u}_δ in consecutive intervals are consecutive multiples of δ , it turns out that

$$m_B = m_A + (k-h+1),$$

and therefore

$$(k-h-1)\delta = (k-h+1)\delta - 2\delta = (m_B - m_A)\delta - 2\delta \geq B - A - 4\varepsilon - 2\delta.$$

Plugging this inequality into (3-17), and letting $\delta \rightarrow 0^+$, we obtain (3-16), which completes the proof. \square

The following result is a straightforward consequence of [Proposition 3.2](#).

Corollary 3.3. *Let us assume that $u_\delta \rightarrow u$ in $L^p(\mathbb{R})$, and let $(a, b) \subseteq \mathbb{R}$ be an interval whose endpoints a and b are Lebesgue points of u .*

Then it turns out that

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (a, b)) \geq \frac{2}{p} \cdot C_p \cdot \frac{|u(b) - u(a)|^p}{(b - a)^{p-1}}.$$

Proof. It is enough to apply [Proposition 3.2](#) with $A := \min\{u(a), u(b)\}$ and $B := \max\{u(a), u(b)\}$. Assumptions (3-3) and (3-4) are satisfied because a and b are Lebesgue points of the limit of the sequence u_δ . \square

We conclude with another variant of [Proposition 3.2](#). We do not need this statement in the sequel, but we think that it clarifies once more the relation between oscillations of u_δ and values of $\Lambda_{\delta,p}(u_\delta, (a, b))$.

Corollary 3.4. *Let $(a, b) \subseteq \mathbb{R}$ be an interval, let $\{u_\delta\}_{\delta>0} \subseteq L^p((a, b))$ be a family of functions, and let $\text{osc}(u_\delta, (a, b))$ denote the essential oscillation of u_δ in (a, b) .*

Then it turns out that

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (a, b)) \geq \frac{2}{p} C_p \frac{1}{(b - a)^{p-1}} \left(\liminf_{\delta \rightarrow 0^+} \text{osc}(u_\delta, (a, b)) \right)^p.$$

Proof. Let i_δ and s_δ denote the essential infimum and the essential supremum of $u_\delta(x)$ in (a, b) , respectively. Let us assume that i_δ and s_δ are real numbers (otherwise an analogous argument works with standard minor changes). Let us set $w_\delta(x) := u_\delta(x) - i_\delta$, and let us observe that

$$\Lambda_{\delta,p}(u_\delta, (a, b)) = \Lambda_{\delta,p}(w_\delta, (a, b)) \quad \text{for all } \delta > 0.$$

Now it is enough to apply [Proposition 3.2](#) with $A := 0$ and

$$B := \liminf_{\delta \rightarrow 0^+} (s_\delta - i_\delta) = \liminf_{\delta \rightarrow 0^+} \text{osc}(u_\delta, (a, b)). \quad \square$$

3B. Piecewise affine approximation. The value of $\Lambda_{0,p}(u, \mathbb{R})$ is the supremum of $\Lambda_{0,p}(v, \mathbb{R})$ as v ranges over a sequence of piecewise affine functions that approximate u . The formal statement is the following (we omit the standard proof, based on the convexity of the norm).

Lemma 3.5 (piecewise affine horizontal segmentation). *Let $p \geq 1$ be a real number, and let $u \in L^p(\mathbb{R})$.*

Then there exists $c \in \mathbb{R}$ such that $c + q$ is a Lebesgue point of u for every $q \in \mathbb{Q}$.

Moreover, if for every positive integer k we consider the piecewise affine function $v_k : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$v_k\left(c + \frac{i}{k}\right) = u\left(c + \frac{i}{k}\right) \quad \text{for all } i \in \mathbb{Z},$$

then it turns out that

$$\Lambda_{0,p}(u, \mathbb{R}) = \lim_{k \rightarrow +\infty} \int_{\mathbb{R}} |v'_k(x)|^p dx = \sup_{k \geq 1} \int_{\mathbb{R}} |v'_k(x)|^p dx.$$

3C. Proof of Gamma-liminf inequality in dimension 1. We are now ready to prove (1-7) in the case $d = 1$. The idea is that Corollary 3.3 represents a “localized” version of the liminf inequality (1-7), which now follows from well-established techniques; see for example [Gobbino 1998; Gobbino and Mora 2001]. To this end, let $u_\delta \rightarrow u$ be any family converging in $L^p(\mathbb{R})$, and let c and v_k be as in Lemma 3.5. For every $i \in \mathbb{Z}$, we set $c_{k,i} := c + i/k$, and we apply Corollary 3.3 in the interval $(c_{k,i}, c_{k,i+1})$. We obtain

$$\liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (c_{k,i}, c_{k,i+1})) \geq \frac{2}{p} C_p \frac{|u(c_{k,i+1}) - u(c_{k,i})|^p}{(1/k)^{p-1}} = \frac{2}{p} C_p \int_{c_{k,i}}^{c_{k,i+1}} |v'_k(x)|^p dx.$$

Since

$$\Lambda_{\delta,p}(u_\delta, \mathbb{R}) \geq \sum_{i \in \mathbb{Z}} \Lambda_{\delta,p}(u_\delta, (c_{k,i}, c_{k,i+1})) \quad \text{for all } \delta > 0,$$

we deduce

$$\begin{aligned} \liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, \mathbb{R}) &\geq \liminf_{\delta \rightarrow 0^+} \sum_{i \in \mathbb{Z}} \Lambda_{\delta,p}(u_\delta, (c_{k,i}, c_{k,i+1})) \\ &\geq \sum_{i \in \mathbb{Z}} \liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, (c_{k,i}, c_{k,i+1})) \\ &\geq \frac{2}{p} C_p \sum_{i \in \mathbb{Z}} \int_{c_{k,i}}^{c_{k,i+1}} |v'_k(x)|^p dx = \frac{2}{p} C_p \int_{\mathbb{R}} |v'_k(x)|^p dx. \end{aligned}$$

Letting $k \rightarrow +\infty$, and recalling (3-1), we obtain exactly (1-7). □

3D. Proof of Gamma-limsup inequality in dimension 1. This subsection is devoted to a proof of statement (2) of Theorem 1.1 in the case $d = 1$.

It is well known that we can limit ourselves to showing the existence of recovery families for every u belonging to a subset of $L^p(\mathbb{R})$ that is dense in energy with respect to $\Lambda_{0,p}(u, \mathbb{R})$. Classical examples of subsets that are dense in energy are the space $C_c^\infty(\mathbb{R})$ of functions of class C^∞ with compact support and the space of piecewise affine functions with compact support. Here for the sake of generality we consider the space $PC_c^1(\mathbb{R})$ of piecewise C^1 functions with compact support, defined as follows.

Definition 3.6. Let $u : \mathbb{R} \rightarrow \mathbb{R}$ be a function. We say that $u \in PC_c^1(\mathbb{R})$ if u has compact support, it is Lipschitz continuous, and there exists a finite subset $S \subseteq \mathbb{R}$ such that $u \in C^1(\mathbb{R} \setminus S)$.

We show that for every $u \in PC_c^1(\mathbb{R})$ the family $S_\delta u$ of vertical δ -segmentations of u is a recovery family. This proves the Gamma-limsup inequality in dimension 1.

Proposition 3.7 (existence of recovery families). *Let $p \geq 1$ be a real number, and let $u \in PC_c^1(\mathbb{R})$ be a piecewise C^1 function with compact support according to Definition 3.6. For every $\delta > 0$, let $S_\delta u$ denote the vertical δ -segmentation of u according to Definition 3.1.*

Then it turns out that

$$\limsup_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(S_\delta u, \mathbb{R}) \leq \frac{2}{p} C_p \int_{\mathbb{R}} |u'(x)|^p dx. \tag{3-18}$$

Proof. To begin with, we introduce some notation. Let $R_0 \geq 1$ be any real number such that the support of u is contained in $[-R_0 + 1, R_0 - 1]$. Let L be the Lipschitz constant of u in \mathbb{R} , and let $S \subseteq \mathbb{R}$ be a finite set such that $u \in C^1(\mathbb{R} \setminus S)$. For every $x \in \mathbb{R}$ and every $\delta > 0$ we set

$$J(\delta, u, x) := \{y \in \mathbb{R} : |S_\delta u(y) - S_\delta u(x)| > \delta\}, \tag{3-19}$$

and

$$H_{\delta,p}(x) := \int_{J(\delta,u,x)} \frac{\delta^p}{|y-x|^{1+p}} dy,$$

so that

$$\Lambda_{\delta,p}(S_\delta u, \mathbb{R}) = \int_{\mathbb{R}} H_{\delta,p}(x) dx \quad \text{for all } \delta > 0. \tag{3-20}$$

In the sequel we call $H_{\delta,p}(x)$ the ‘‘pointwise hostility function’’. It represents the contribution of each point x to the double integral defining $\Lambda_{\delta,p}(S_\delta u, \mathbb{R})$.

Strategy of the proof: The outline of the proof is the following. First of all, we show that

$$\lim_{\delta \rightarrow 0^+} \int_{-\infty}^{-R_0} H_{\delta,p}(x) dx = \lim_{\delta \rightarrow 0^+} \int_{R_0}^{+\infty} H_{\delta,p}(x) dx = 0. \tag{3-21}$$

Then we define an averaged pointwise hostility function $\widehat{H}_{\delta,p}(x)$ with the property that

$$\int_{-R_0}^{R_0} H_{\delta,p}(x) dx = \int_{-R_0}^{R_0} \widehat{H}_{\delta,p}(x) dx. \tag{3-22}$$

We also show that the averaged pointwise hostility function satisfies the uniform bound

$$\widehat{H}_{\delta,p}(x) \leq \frac{2}{p} L^p \quad \text{for all } x \in [-R_0, R_0], \text{ for all } \delta > 0, \tag{3-23}$$

and the asymptotic estimate

$$\limsup_{\delta \rightarrow 0^+} \widehat{H}_{\delta,p}(x) \leq \frac{2}{p} C_p |u'(x)|^p \quad \text{for all } x \in [-R_0, R_0] \setminus S. \tag{3-24}$$

At this point, from Fatou’s lemma we deduce

$$\limsup_{\delta \rightarrow 0^+} \int_{-R_0}^{R_0} H_{\delta,p}(x) dx = \limsup_{\delta \rightarrow 0^+} \int_{-R_0}^{R_0} \widehat{H}_{\delta,p}(x) dx \leq \int_{-R_0}^{R_0} \limsup_{\delta \rightarrow 0^+} \widehat{H}_{\delta,p}(x) dx \leq \frac{2}{p} C_p \int_{-R_0}^{R_0} |u'(x)|^p dx.$$

Keeping (3-20) and (3-21) into account, this estimate implies (3-18).

Reducing integration to a bounded interval: We prove (3-21).

To this end, let us consider any $x \leq -R_0$. We observe that in this case the set $J(\delta, u, x)$ defined in (3-19) is contained in the support of u , and hence

$$\int_{-\infty}^{-R_0} H_{\delta,p}(x) dx \leq \delta^p \int_{-\infty}^{-R_0} dx \int_{-R_0+1}^{R_0-1} \frac{1}{|y-x|^{1+p}} dy.$$

At this point the first limit in (3-21) follows from the convergence of the double integral. The proof of the second limit is analogous.

Uniform bound on the pointwise hostility function: We prove that

$$H_{\delta,p}(x) \leq \frac{2}{p} L^p \quad \text{for all } x \in [-R_0, R_0], \text{ for all } \delta > 0. \quad (3-25)$$

To this end, we observe that the implication

$$|S_\delta u(y) - S_\delta u(x)| > \delta \quad \implies \quad |u(y) - u(x)| > \delta$$

holds true for every $(x, y) \in \mathbb{R}^2$. Since u is Lipschitz continuous, we deduce that

$$|S_\delta u(y) - S_\delta u(x)| > \delta \quad \implies \quad |y - x| \geq \frac{\delta}{L},$$

and hence

$$H_{\delta,p}(x) \leq \int_{|y-x| \geq \delta/L} \frac{\delta^p}{|y-x|^{1+p}} dy = 2 \int_{\delta/L}^{+\infty} \frac{\delta^p}{z^{1+p}} dz = \frac{2}{p} L^p,$$

as required.

Averaged pointwise hostility function: In this part of the proof we introduce the averaged pointwise hostility function. To this end, we consider the open set

$$A(u, \delta) := \{x \in (-R_0, R_0) : u(x) \notin \delta\mathbb{Z}\}.$$

A connected component (a, b) of $A(u, \delta)$ is called *monotone* if $[a, b] \cap S = \emptyset$, and $|u'(x)| \geq \delta$ for every $x \in [a, b]$. In this case there exists $k \in \mathbb{Z}$ such that $u(a) = k\delta$ and $u(b) = k\delta \pm \delta$, where the sign depends on the sign of $u'(x)$ in (a, b) . From the Lipschitz continuity of u we deduce that $A(u, \delta)$ has only a finite number of monotone connected components.

The averaged pointwise hostility function $\widehat{H}_{\delta,p} : \mathbb{R} \rightarrow \mathbb{R}$ is defined as

$$\widehat{H}_{\delta,p}(x) := \frac{1}{b-a} \int_a^b H_{\delta,p}(s) ds$$

if $x \in [a, b)$ for some monotone connected component of $A(\delta, u)$, and $\widehat{H}_{\delta,p}(x) := H_{\delta,p}(x)$ otherwise.

At this point, inequality (3-23) follows from (3-25), while (3-22) is true because the integrals of $H_{\delta,p}(x)$ and $\widehat{H}_{\delta,p}(x)$ are the same both in all monotone connected components, and in the complement set.

Asymptotic estimate in stationary points: We prove that (3-24) holds true for every $x \in (-R_0, R_0) \setminus S$ with $|u'(x)| = 0$.

To begin with, we observe that in this case $x \notin [a, b)$ for every monotone connected component (a, b) of $A(\delta, u)$ (because $|u'(x)|$ is strictly positive in the closure of every monotone connected component), and therefore $\widehat{H}_{\delta,p}(x) = H_{\delta,p}(x)$ for every $\delta > 0$.

If $J(\delta, u, x) = \emptyset$ for every $\delta > 0$, then u is identically null, and the conclusion is trivial. Otherwise $J(\delta, u, x) \neq \emptyset$ when δ is small enough. In this case, let r_δ be the largest positive real number such that

$$(x - r_\delta, x + r_\delta) \cap J(\delta, u, x) = \emptyset,$$

so that

$$H_{\delta,p}(x) \leq \int_{-\infty}^{x-r_\delta} \frac{\delta^p}{|y-x|^{1+p}} dy + \int_{x+r_\delta}^{+\infty} \frac{\delta^p}{|y-x|^{1+p}} dy = \frac{2}{p} \left(\frac{\delta}{r_\delta} \right)^p.$$

Let $\delta_k \rightarrow 0^+$ be any sequence such that

$$\limsup_{\delta \rightarrow 0^+} \frac{\delta}{r_\delta} = \lim_{k \rightarrow +\infty} \frac{\delta_k}{r_{\delta_k}}. \quad (3-26)$$

Up to subsequences, we can also assume that r_{δ_k} tends to some r_0 . If $r_0 > 0$, then the limit in the right-hand side of (3-26) is 0, which proves (3-24) in this case. If $r_0 = 0$, then from the maximality of r_{δ_k} we deduce that $|u(x \pm r_{\delta_k}) - u(x)| = \delta_k$ for a suitable choice of the sign, which might depend on k . In any case, the limit in the right-hand side of (3-26) turns out to be

$$\lim_{k \rightarrow +\infty} \frac{\delta_k}{r_{\delta_k}} = \lim_{k \rightarrow +\infty} \frac{|u(x \pm r_{\delta_k}) - u(x)|}{r_{\delta_k}} = |u'(x)| = 0,$$

which proves (3-24) also in this case.

Asymptotic estimate in nonstationary points: We prove that (3-24) holds true for every $x \in (-R_0, R_0) \setminus S$ with $|u'(x)| > 0$.

Let us assume, without loss of generality, that $u'(x) > 0$ (the other case is analogous). Then for every $\delta > 0$ small enough it turns out that x lies in the closure of a monotone connected component of $A(\delta, u)$. More precisely, there exist four real numbers $a_\delta, b_\delta, c_\delta, d_\delta$ with

$$a_\delta < b_\delta \leq x < c_\delta < d_\delta,$$

and $k_\delta \in \mathbb{Z}$ such that

$$u(a_\delta) = (k_\delta - 1)\delta, \quad u(b_\delta) = k_\delta\delta, \quad u(c_\delta) = (k_\delta + 1)\delta, \quad u(d_\delta) = (k_\delta + 2)\delta,$$

and

$$u(y) \in ((k_\delta - 1)\delta, k_\delta\delta) \quad \text{for all } y \in (a_\delta, b_\delta), \quad (3-27)$$

$$u(y) \in (k_\delta\delta, (k_\delta + 1)\delta) \quad \text{for all } y \in (b_\delta, c_\delta), \quad (3-28)$$

$$u(y) \in ((k_\delta + 1)\delta, (k_\delta + 2)\delta) \quad \text{for all } y \in (c_\delta, d_\delta). \quad (3-29)$$

We observe that $a_\delta, b_\delta, c_\delta$, and d_δ tend to x as $\delta \rightarrow 0^+$, and hence

$$\lim_{\delta \rightarrow 0^+} \frac{\delta}{b_\delta - a_\delta} = \lim_{\delta \rightarrow 0^+} \frac{u(b_\delta) - u(a_\delta)}{b_\delta - a_\delta} = u'(x). \quad (3-30)$$

Similarly it turns out that

$$\lim_{\delta \rightarrow 0^+} \frac{\delta}{c_\delta - b_\delta} = \lim_{\delta \rightarrow 0^+} \frac{\delta}{d_\delta - c_\delta} = u'(x), \quad (3-31)$$

$$\lim_{\delta \rightarrow 0^+} \frac{\delta}{c_\delta - a_\delta} = \lim_{\delta \rightarrow 0^+} \frac{\delta}{d_\delta - b_\delta} = \frac{u'(x)}{2}. \quad (3-32)$$

From (3-27) through (3-29) we deduce that

$$J(\delta, u, s) \subseteq (-\infty, a_\delta] \cup [d_\delta, +\infty) \quad \text{for all } s \in (b_\delta, c_\delta).$$

It follows that

$$H_{\delta,p}(s) \leq \int_{\mathbb{R} \setminus (a_\delta, d_\delta)} \frac{\delta^p}{|y-s|^{1+p}} dy = \frac{\delta^p}{p} \left(\frac{1}{(d_\delta-s)^p} + \frac{1}{(s-a_\delta)^p} \right) \quad \text{for all } s \in [b_\delta, c_\delta),$$

and hence

$$\widehat{H}_{\delta,p}(x) = \frac{1}{c_\delta - b_\delta} \int_{b_\delta}^{c_\delta} H_{\delta,p}(s) ds \leq \frac{\delta^p}{p} \frac{1}{c_\delta - b_\delta} \int_{b_\delta}^{c_\delta} \left(\frac{1}{(d_\delta-s)^p} + \frac{1}{(s-a_\delta)^p} \right) ds \quad (3-33)$$

for every $x \in [b_\delta, c_\delta)$. Now we distinguish two cases.

- If $p = 1$, computing the integrals in (3-33) we obtain

$$\widehat{H}_{\delta,p}(x) \leq \frac{\delta}{c_\delta - b_\delta} \log \left(\frac{d_\delta - b_\delta}{\delta} \cdot \frac{\delta}{d_\delta - c_\delta} \cdot \frac{c_\delta - a_\delta}{\delta} \cdot \frac{\delta}{b_\delta - a_\delta} \right),$$

and therefore (3-24) follows from (3-30) through (3-32).

- If $p > 1$, computing the integrals in (3-33) we obtain

$$\widehat{H}_{\delta,p}(x) \leq \frac{1}{p(p-1)} \frac{\delta}{c_\delta - b_\delta} \left\{ \frac{\delta^{p-1}}{(d_\delta - c_\delta)^{p-1}} + \frac{\delta^{p-1}}{(b_\delta - a_\delta)^{p-1}} - \frac{\delta^{p-1}}{(d_\delta - b_\delta)^{p-1}} - \frac{\delta^{p-1}}{(c_\delta - a_\delta)^{p-1}} \right\},$$

and therefore also in this case (3-24) follows from (3-30) through (3-32). \square

3E. Smooth recovery families. The aim of this subsection is refining the Gamma-limsup inequality by showing the existence of recovery families consisting of C^∞ functions with compact support. To this end, we introduce the following notion.

Definition 3.8 (δ -step functions). Let δ be a positive real number. A function $u : \mathbb{R} \rightarrow \mathbb{R}$ is called a δ -step function if there exists a positive integer n , an $(n+1)$ -tuple $x_0 < x_1 < \dots < x_n$ of real numbers, and $(k_1, \dots, k_n) \in \mathbb{Z}^n$ such that

- $u(x) = 0$ for every $x \in (-\infty, x_0) \cup (x_n, +\infty)$,
- $u(x) = k_i \delta$ in (x_{i-1}, x_i) for every $i = 1, \dots, n$,
- $|k_1| = |k_n| = 1$ and $|k_i - k_{i-1}| = 1$ for every $i = 2, \dots, n$.

The values of $u(x)$ for $x \in \{x_0, x_1, \dots, x_n\}$ are not relevant (just to fix ideas, we can define $u(x_i)$ as the maximum between the limit of $u(x)$ as $x \rightarrow x_i^+$ and the limit of $u(x)$ as $x \rightarrow x_i^-$).

Now we show that, for every fixed $\delta > 0$, every δ -step function can be approximated in energy by functions of class C^∞ with compact support. Roughly speaking, this is possible because the rigid structure of δ -step functions allows us to control the effect of convolutions, which otherwise is unpredictable due to the sensitivity of the integration region in (1-1) to small perturbations.

Proposition 3.9 (smooth approximation of δ -step functions). Let $\delta > 0$ and $p \geq 1$ be real numbers, and let $u : \mathbb{R} \rightarrow \mathbb{R}$ be a δ -step function.

Then there exists a family $\{u_\varepsilon\}_{\varepsilon>0} \subseteq C_c^\infty(\mathbb{R})$ such that

$$\lim_{\varepsilon \rightarrow 0^+} u_\varepsilon = u \quad \text{in } L^p(\mathbb{R}),$$

and

$$\lim_{\varepsilon \rightarrow 0^+} \Lambda_{\delta,p}(u_\varepsilon, \mathbb{R}) = \Lambda_{\delta,p}(u, \mathbb{R}).$$

Proof. Let n , x_i and k_i be as in the definition of δ -step functions, and let

$$\tau := \min\{x_i - x_{i-1} : i = 1, \dots, n\}$$

be the length of the smallest interval of the partition. We observe that points in neighboring intervals do not contribute to the computation of $\Lambda_{\delta,p}(u, \mathbb{R})$. In particular, if we write as usual

$$\Lambda_{\delta,p}(u, \mathbb{R}) := \iint_{I(\delta, u, \mathbb{R})} \frac{\delta^p}{|y-x|^{1+p}} dx dy,$$

then it turns out that

$$|y-x| \geq \tau \quad \text{for all } (x, y) \in I(\delta, u, \mathbb{R}). \quad (3-34)$$

Let us fix a mollifier $\rho \in C_c^\infty(\mathbb{R})$ with

- $\rho(x) \geq 0$ for every $x \in \mathbb{R}$,
- $\rho(x) = 0$ for every $x \in \mathbb{R}$ with $|x| \geq 1$,
- $\int_{\mathbb{R}} \rho(x) dx = 1$,

and let us consider the usual regularization by convolution

$$u_\varepsilon(x) := \int_{\mathbb{R}} u(x + \varepsilon y) \rho(y) dy.$$

It is well known that $u_\varepsilon \in C_c^\infty(\mathbb{R})$ for every $\varepsilon > 0$, and that for every $p \geq 1$ it turns out that $u_\varepsilon \rightarrow u$ in $L^p(\mathbb{R})$ as $\varepsilon \rightarrow 0^+$.

Let us assume that $2\varepsilon < \tau$, let us consider the two open sets

$$A_\varepsilon := \bigcup_{i=0}^n (x_i - \varepsilon, x_i + \varepsilon) \subseteq \mathbb{R}, \quad B_\varepsilon := (A_\varepsilon \times \mathbb{R}) \cup (\mathbb{R} \times A_\varepsilon) \subseteq \mathbb{R}^2,$$

and let us write

$$\Lambda_{\delta,p}(u_\varepsilon, \mathbb{R}) = \iint_{I(\delta, u_\varepsilon, \mathbb{R}) \cap B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy + \iint_{I(\delta, u_\varepsilon, \mathbb{R}) \setminus B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy.$$

Since the support of ρ is contained in $[-1, 1]$, it turns out that $u_\varepsilon(x) = u(x)$ for every $x \in \mathbb{R} \setminus A_\varepsilon$. It follows that

$$I(\delta, u_\varepsilon, \mathbb{R}) \setminus B_\varepsilon = I(\delta, u, \mathbb{R}) \setminus B_\varepsilon,$$

and therefore

$$\lim_{\varepsilon \rightarrow 0^+} \iint_{I(\delta, u_\varepsilon, \mathbb{R}) \setminus B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy = \lim_{\varepsilon \rightarrow 0^+} \iint_{I(\delta, u, \mathbb{R}) \setminus B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy = \Lambda_{\delta,p}(u, \mathbb{R}),$$

where the last equality follows from Lebesgue’s dominated convergence theorem because B_ε shrinks to a set of null measure. So it remains to show that

$$\lim_{\varepsilon \rightarrow 0^+} \iint_{I(\delta, u_\varepsilon, \mathbb{R}) \cap B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy = 0. \tag{3-35}$$

To this end, from (3-34) and the properties of the support of the mollifier, we deduce that now

$$|y-x| \geq \tau - 2\varepsilon \quad \text{for all } (x, y) \in I(\delta, u_\varepsilon, \mathbb{R}),$$

and therefore

$$\begin{aligned} \iint_{I(\delta, u_\varepsilon, \mathbb{R}) \cap B_\varepsilon} \frac{\delta^p}{|y-x|^{1+p}} dx dy &\leq 2 \sum_{i=0}^n \int_{x_i-\varepsilon}^{x_i+\varepsilon} dx \int_{|z| \geq \tau-2\varepsilon} \frac{\delta^p}{|z|^{1+p}} dz \\ &\leq 2 \sum_{i=0}^n \int_{x_i-\varepsilon}^{x_i+\varepsilon} \frac{2}{p} \frac{\delta^p}{|\tau-2\varepsilon|^p} dx = \frac{4}{p} \frac{\delta^p}{|\tau-2\varepsilon|^p} \cdot 2\varepsilon(n+1), \end{aligned}$$

which implies (3-35). □

We are now ready to show the existence of smooth recovery families. As usual, it is enough to show the existence of such a family for every u in a subset of $L^p(\mathbb{R})$ which is dense in energy for $\Lambda_{0,p}(u, \mathbb{R})$. In this case we consider the space $PA_c(\mathbb{R})$ of piecewise affine functions with compact support.

Since piecewise affine functions are piecewise C^1 , we know from Proposition 3.7 that the family $S_\delta u$ of vertical δ -segmentations of u is a (nonsmooth) recovery family for u . The key point is that the vertical δ -segmentation of a piecewise affine function with compact support is a δ -step function according to Definition 3.8. Thus from Proposition 3.9 we deduce the existence of a function $u_\delta \in C_c^\infty(\mathbb{R})$ such that

$$\|u_\delta - S_\delta u\|_{L^p(\mathbb{R})} \leq \delta \quad \text{and} \quad \Lambda_{\delta,p}(u_\delta, \mathbb{R}) \leq \Lambda_{\delta,p}(S_\delta u, \mathbb{R}) + \delta$$

for every $\delta > 0$. This implies that $\{u_\delta\}$ is a smooth recovery family for u . □

4. Gamma-convergence in any dimension

It remains to prove Theorem 1.1 in any space dimension. This follows from well-established sectioning techniques. For every $\sigma \in \mathbb{S}^{d-1}$, let $\langle \sigma \rangle^\perp$ denote the hyperplane orthogonal to σ , namely

$$\langle \sigma \rangle^\perp := \{z \in \mathbb{R}^d : \langle z, \sigma \rangle = 0\}.$$

Given any $u : \mathbb{R}^d \rightarrow \mathbb{R}$, for every $\sigma \in \mathbb{S}^{d-1}$ and every $z \in \langle \sigma \rangle^\perp$, we consider the 1-dimensional section $u_{\sigma,z} : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$u_{\sigma,z}(x) := u(z + \sigma x) \quad \text{for all } x \in \mathbb{R}.$$

The main idea is that Sobolev norms, total variation, and functionals such as $\Lambda_{\delta,p}$ computed in u are a sort of average of the same quantities computed on the 1-dimensional sections $u_{\sigma,z}$. The result is the following.

Proposition 4.1 (integral-geometric representation). *Let $u : \mathbb{R}^d \rightarrow \mathbb{R}$ be any measurable function. Let $\Lambda_{\delta,p}$ and $\Lambda_{0,p}$ be the functionals defined in (1-1) and (1-2), respectively.*

(1) For every $p \geq 1$ it turns out that

$$\int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} \Lambda_{0,p}(u_{\sigma,z}, \mathbb{R}) dz = G_{d,p} \Lambda_{0,p}(u, \mathbb{R}^d),$$

where $G_{d,p}$ is the geometric constant defined in (1-4).

(2) For every $\delta > 0$ and every $p \geq 1$ it turns out that

$$\int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} \Lambda_{\delta,p}(u_{\sigma,z}, \mathbb{R}) dz = 2\Lambda_{\delta,p}(u, \mathbb{R}^d). \quad \square$$

We skip the details of the proof of Proposition 4.1, which is a simple application of variable changes in multiple integrals. More generally, for every $\sigma \in \mathbb{S}^{d-1}$ and every $g \in L^1(\mathbb{R}^d)$ it turns out that

$$\int_{\mathbb{R}^d} g(y) dy = \int_{\langle \sigma \rangle^\perp} dz \int_{\mathbb{R}} g(z + \sigma x) dx,$$

and this is the main ingredient in the proof of statement (1).

Similarly, for every $g \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$ it turns out that

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} g(u, v) du dv = \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} dz \iint_{\mathbb{R} \times \mathbb{R}} g(z + \sigma x, z + \sigma y) \cdot |y - x|^{d-1} dx dy,$$

and this is the main ingredient in the proof of statement (2).

We are now ready to prove Theorem 1.1.

Proof. Gamma-liminf: Let us assume that $u_\delta \rightarrow u$ in $L^1(\mathbb{R}^d)$. Then for every $\sigma \in \mathbb{S}^{d-1}$ it turns out that

$$(u_\delta)_{\sigma,z} \rightarrow u_{\sigma,z} \quad \text{in } L^1(\mathbb{R})$$

for almost every $z \in \langle \sigma \rangle^\perp$. Therefore, from the integral-geometric representations of Proposition 4.1, Fatou’s lemma, and the 1-dimensional result, we obtain

$$\begin{aligned} \liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(u_\delta, \mathbb{R}^d) &= \liminf_{\delta \rightarrow 0^+} \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} \Lambda_{\delta,p}((u_\delta)_{\sigma,z}, \mathbb{R}) dz \\ &\geq \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} \liminf_{\delta \rightarrow 0^+} \Lambda_{\delta,p}((u_\delta)_{\sigma,z}, \mathbb{R}) dz \\ &\geq \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{\langle \sigma \rangle^\perp} \frac{2}{p} C_p \Lambda_{0,p}(u_{\sigma,z}, \mathbb{R}) dz \\ &= \frac{1}{p} G_{d,p} C_p \Lambda_{0,p}(u, \mathbb{R}^d). \end{aligned}$$

Gamma-limsup: Let $u \in C_c^\infty(\mathbb{R}^d)$ be any function with compact support. For every $\delta > 0$ we consider the vertical δ -segmentation $S_\delta u$ of u , and we observe that this operation commutes with the 1-dimensional sections, in the sense that

$$(S_\delta u)_{\sigma,z} = S_\delta(u_{\sigma,z}) \quad \text{for all } \sigma \in \mathbb{S}^{d-1}, \text{ for all } z \in \langle \sigma \rangle^\perp.$$

Therefore, from the integral-geometric representations of [Proposition 4.1](#), Fatou's lemma, and the 1-dimensional result, we obtain

$$\begin{aligned}
\limsup_{\delta \rightarrow 0^+} \Lambda_{\delta,p}(S_\delta u, \mathbb{R}^d) &= \limsup_{\delta \rightarrow 0^+} \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{(\sigma)^\perp} \Lambda_{\delta,p}((S_\delta u)_{\sigma,z}, \mathbb{R}) dz \\
&\leq \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{(\sigma)^\perp} \limsup_{\delta \rightarrow 0^+} \Lambda_{\delta,p}((S_\delta u)_{\sigma,z}, \mathbb{R}) dz \\
&\leq \frac{1}{2} \int_{\mathbb{S}^{d-1}} d\sigma \int_{(\sigma)^\perp} \frac{2}{p} C_p \Lambda_{0,p}(u_{\sigma,z}, \mathbb{R}) dz \\
&= \frac{1}{p} G_{d,p} C_p \Lambda_{0,p}(u, \mathbb{R}^d).
\end{aligned}$$

The δ -independent bounds on $\Lambda_{\delta,p}((S_\delta u)_{\sigma,z}, \mathbb{R})$ needed in order to apply Fatou's lemma follow from the Lipschitz continuity of u and the boundedness of its support.

Smooth recovery families: It remains to show the existence of smooth recovery families. The strategy is analogous to the 1-dimensional case, and therefore we limit ourselves to outlining the argument, sparing the reader all technicalities.

To begin with, we observe that the space $PA_c(\mathbb{R}^d)$ of piecewise affine functions with compact support is a subspace of $L^p(\mathbb{R}^d)$ that is dense in energy for $\Lambda_{0,p}(u, \mathbb{R}^d)$. This is true because $C_c^\infty(\mathbb{R}^d)$ is dense in energy, and in turn any function in $C_c^\infty(\mathbb{R}^d)$ can be approximated in $W^{1,\infty}(\mathbb{R}^d)$ by functions in $PA_c(\mathbb{R}^d)$; see for example Chapter 4 in [\[Brenner and Scott 1994\]](#), and in particular Corollary 4.4.24.

As a consequence, it is enough to show the existence of a recovery family for every $u \in PA_c(\mathbb{R}^d)$, in which case a nonsmooth recovery family is provided by the vertical δ -segmentations $S_\delta u$ of u . On the other hand, vertical δ -segmentations of piecewise affine functions with compact support are δ -step functions, and these functions can be approximated in energy by smooth functions. It follows that for every $\delta > 0$ there exists $u_\delta \in C_c^\infty(\mathbb{R}^d)$ such that

$$\|u_\delta - S_\delta u\|_{L^p(\mathbb{R}^d)} \leq \delta \quad \text{and} \quad \Lambda_{\delta,p}(u_\delta, \mathbb{R}^d) \leq \Lambda_{\delta,p}(S_\delta u, \mathbb{R}^d) + \delta,$$

and therefore $\{u_\delta\}$ is the required recovery family.

The last approximation step can be proved by convolution as we did in [Proposition 3.9](#). To be more precise, a δ -step function in dimension d is a function $v : \mathbb{R}^d \rightarrow \mathbb{R}$ with the property that there exist a finite set $\{P_1, \dots, P_m\}$ of disjoint open polytopes (bounded intersections of half-spaces) and integers k_1, \dots, k_m such that

- $v(x) = k_i \delta$ in P_i for every $i = 1, \dots, m$,
- $v(x) = 0$ in the open set P_0 defined as the complement set of the closure of $P_1 \cup \dots \cup P_m$,
- $|k_i - k_j| \leq 1$ whenever the closure of P_i intersects the closure of P_j ,
- $|k_i| \leq 1$ whenever the closure of P_i intersects the closure of P_0 .

In words, the level sets of a δ -step function are finite unions of polytopes, and values in adjacent regions differ by δ .

The key point is that for every δ -step function v there exists a positive real number τ such that

$$(x, y) \in I(\delta, v, \mathbb{R}^d) \implies |y - x| \geq \tau.$$

As a consequence, when we define v_ε as the convolution of v with a mollifier whose support is contained in the ball with center in the origin and radius ε , we obtain

$$(x, y) \in I(\delta, v_\varepsilon, \mathbb{R}^d) \implies |y - x| \geq \tau - 2\varepsilon,$$

and at this point the conclusion follows exactly as in the proof of [Proposition 3.9](#). □

Acknowledgments

Gobbino was introduced to this family of nonlocal functionals by an inspiring talk given by H. Brezis [\[2016\]](#) during the congress “A mathematical tribute to Ennio De Giorgi”, held in Pisa in September 2016 on the 20th anniversary of his death. Gobbino was introduced to nonlocal approximations of free discontinuity problems by E. De Giorgi himself in the last year of his life.

We are all deeply grateful to both of them.

References

- [Antonucci et al. 2018] C. Antonucci, M. Gobbino, M. Migliorini, and N. Picenni, “On the shape factor of interaction laws for a non-local approximation of the Sobolev norm and the total variation”, *C. R. Math. Acad. Sci. Paris* **356**:8 (2018), 859–864. [MR](#) [Zbl](#)
- [Antonucci et al. 2020] C. Antonucci, M. Gobbino, and N. Picenni, “On the gap between the Gamma-limit and the pointwise limit for a nonlocal approximation of the total variation”, *Anal. PDE* **13**:3 (2020), 627–649.
- [Bourgain and Nguyen 2006] J. Bourgain and H.-M. Nguyen, “A new characterization of Sobolev spaces”, *C. R. Math. Acad. Sci. Paris* **343**:2 (2006), 75–80. [MR](#) [Zbl](#)
- [Bourgain et al. 2001] J. Bourgain, H. Brezis, and P. Mironescu, “Another look at Sobolev spaces”, pp. 439–455 in *Optimal control and partial differential equations* (Paris, 2000), edited by J. L. Menaldi et al., IOS, Amsterdam, 2001. [MR](#) [Zbl](#)
- [Bourgain et al. 2005] J. Bourgain, H. Brezis, and P. Mironescu, “Lifting, degree, and distributional Jacobian revisited”, *Comm. Pure Appl. Math.* **58**:4 (2005), 529–551. [MR](#) [Zbl](#)
- [Brenner and Scott 1994] S. C. Brenner and L. R. Scott, *The mathematical theory of finite element methods*, Texts in Appl. Math. **15**, Springer, 1994. [MR](#) [Zbl](#)
- [Brezis 2015] H. Brezis, “New approximations of the total variation and filters in imaging”, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **26**:2 (2015), 223–240. [MR](#) [Zbl](#)
- [Brezis 2016] H. Brezis, “Another triumph for De Giorgi’s Gamma convergence”, video lecture, 2016, available at <https://youtu.be/1Y6fvZX1fx8>.
- [Brezis and Nguyen 2017] H. Brezis and H.-M. Nguyen, “Non-convex, non-local functionals converging to the total variation”, *C. R. Math. Acad. Sci. Paris* **355**:1 (2017), 24–27. [MR](#) [Zbl](#)
- [Brezis and Nguyen 2018] H. Brezis and H.-M. Nguyen, “Non-local functionals related to the total variation and connections with image processing”, *Ann. PDE* **4**:1 (2018), art. id. 9. [MR](#) [Zbl](#)
- [Chambolle 1995] A. Chambolle, “Image segmentation by variational methods: Mumford and Shah functional and the discrete approximations”, *SIAM J. Appl. Math.* **55**:3 (1995), 827–863. [MR](#) [Zbl](#)
- [Garsia and Rodemich 1974] A. M. Garsia and E. Rodemich, “Monotonicity of certain functionals under rearrangement”, *Ann. Inst. Fourier (Grenoble)* **24**:2 (1974), 67–116. [MR](#) [Zbl](#)

- [Gobbino 1998] M. Gobbino, “Finite difference approximation of the Mumford–Shah functional”, *Comm. Pure Appl. Math.* **51**:2 (1998), 197–228. [MR](#) [Zbl](#)
- [Gobbino and Mora 2001] M. Gobbino and M. G. Mora, “Finite-difference approximation of free-discontinuity problems”, *Proc. Roy. Soc. Edinburgh Sect. A* **131**:3 (2001), 567–595. [MR](#) [Zbl](#)
- [Nguyen 2006] H.-M. Nguyen, “Some new characterizations of Sobolev spaces”, *J. Funct. Anal.* **237**:2 (2006), 689–720. [MR](#) [Zbl](#)
- [Nguyen 2007] H.-M. Nguyen, “ Γ -convergence and Sobolev norms”, *C. R. Math. Acad. Sci. Paris* **345**:12 (2007), 679–684. [MR](#) [Zbl](#)
- [Nguyen 2008] H.-M. Nguyen, “Further characterizations of Sobolev spaces”, *J. Eur. Math. Soc.* **10**:1 (2008), 191–229. [MR](#) [Zbl](#)
- [Nguyen 2011] H.-M. Nguyen, “ Γ -convergence, Sobolev norms, and BV functions”, *Duke Math. J.* **157**:3 (2011), 495–533. [MR](#) [Zbl](#)
- [Nguyen 2014] H.-M. Nguyen, “Estimates for the topological degree and related topics”, *J. Fixed Point Theory Appl.* **15**:1 (2014), 185–215. [MR](#) [Zbl](#)
- [Ponce 2004] A. C. Ponce, “A new approach to Sobolev spaces and connections to Γ -convergence”, *Calc. Var. Partial Differential Equations* **19**:3 (2004), 229–255. [MR](#) [Zbl](#)
- [Taylor 1973] H. Taylor, “Rearrangements of incidence tables”, *J. Combinatorial Theory Ser. A* **14** (1973), 30–36. [MR](#) [Zbl](#)

Received 27 May 2018. Revised 23 Dec 2018. Accepted 7 Mar 2019.

CLARA ANTONUCCI: clara.antonucci@sns.it
Scuola Normale Superiore, Pisa, Italy

MASSIMO GOBBINO: massimo.gobbino@unipi.it
Università degli Studi di Pisa, Pisa, Italy

MATTEO MIGLIORINI: matteo.migliorini@sns.it
Scuola Normale Superiore, Pisa, Italy

NICOLA PICENNI: nicola.picenni@sns.it
Scuola Normale Superiore, Pisa, Italy

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2020 is US \$340/year for the electronic version, and \$550/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 13 No. 2 2020

Regularity estimates for elliptic nonlocal operators	317
BARTŁOMIEJ DYDA and MORITZ KASSMANN	
On solvability and ill-posedness of the compressible Euler system subject to stochastic forces	371
DOMINIC BREIT, EDUARD FEIREISL and MARTINA HOFMANOVÁ	
Variable coefficient Wolff-type inequalities and sharp local smoothing estimates for wave equations on manifolds	403
DAVID BELTRAN, JONATHAN HICKMAN and CHRISTOPHER D. SOGGE	
On the Hölder continuous subsolution problem for the complex Monge–Ampère equation, II	435
NGOC CUONG NGUYEN	
The Calderón problem for the fractional Schrödinger equation	455
TUHIN GHOSH, MIKKO SALO and GUNTHER UHLMANN	
Sharp Strichartz inequalities for fractional and higher-order Schrödinger equations	477
GIANMARCO BROCCHI, DIOGO OLIVEIRA E SILVA and RENÉ QUILODRÁN	
A bootstrapping approach to jump inequalities and their applications	527
MARIUSZ MIREK, ELIAS M. STEIN and PAVEL ZORIN-KRANICH	
On the trace operator for functions of bounded \mathbb{A} -variation	559
DOMINIC BREIT, LARS DIENING and FRANZ GMEINER	
Optimal constants for a nonlocal approximation of Sobolev norms and total variation	595
CLARA ANTONUCCI, MASSIMO GOBBINO, MATTEO MIGLIORINI and NICOLA PICENNI	