

# Measurement of the shape of the $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$ differential decay rate

---



## The LHCb collaboration

*E-mail:* [rvazquez@cern.ch](mailto:rvazquez@cern.ch)

**ABSTRACT:** The shape of the  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  differential decay rate is obtained as a function of the hadron recoil parameter using proton-proton collision data at a centre-of-mass energy of 13 TeV, corresponding to an integrated luminosity of  $1.7 \text{ fb}^{-1}$  collected by the LHCb detector. The  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decay is reconstructed through the decays  $D_s^{*-} \rightarrow D_s^- \gamma$  and  $D_s^- \rightarrow K^- K^+ \pi^-$ . The differential decay rate is fitted with the Caprini-Lellouch-Neubert (CLN) and Boyd-Grinstein-Lebed (BGL) parametrisations of the form factors, and the relevant quantities for both are extracted.

**KEYWORDS:** B physics, Flavor physics, Hadron-Hadron scattering (experiments)

**ARXIV EPRINT:** [2003.08453](https://arxiv.org/abs/2003.08453)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Formalism of the <math>B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu</math> decay</b>	<b>2</b>
2.1	CLN form-factor parametrisation	4
2.2	BGL form-factor parametrisation	5
<b>3</b>	<b>Detector and simulation</b>	<b>5</b>
<b>4</b>	<b>Data selection</b>	<b>6</b>
<b>5</b>	<b>Signal yield</b>	<b>7</b>
<b>6</b>	<b>Efficiency correction</b>	<b>9</b>
<b>7</b>	<b>Unfolded yields</b>	<b>11</b>
7.1	Systematic uncertainties	12
7.2	Results	13
<b>8</b>	<b>Form factor fits</b>	<b>14</b>
8.1	Systematic uncertainties	15
8.2	Results	15
<b>9</b>	<b>Conclusions</b>	<b>18</b>
<b>A</b>	<b>Fitted yields and efficiency</b>	<b>19</b>
<b>B</b>	<b>Covariance and response matrices</b>	<b>19</b>
<b>C</b>	<b>Additional information BGL fit</b>	<b>21</b>
<b>D</b>	<b>Comparison with <i>Phys. Rev. D</i> 101 (2020) 072004</b>	<b>22</b>
	<b>The LHCb collaboration</b>	<b>26</b>

---

## 1 Introduction

Semileptonic decays of heavy hadrons are commonly used to measure the parameters of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1, 2], as they involve only one hadronic current that can be parametrised in terms of scalar functions known as form factors. The number of form factors needed to describe a particular decay depends upon the spin of the initial- and final-state hadrons [3–5]. For the decay of a pseudoscalar  $B$  meson to a vector

$D^*$  meson, four form factors are required. The determination of the CKM matrix element  $|V_{cb}|$  using  $B \rightarrow D^{(*)}\ell\nu_\ell$  decays or via the inclusive sum of all hadronic  $B \rightarrow X_c\ell\nu_\ell$  decay channels has been giving inconsistent results during the last thirty years [6]. The exclusive determination relies heavily on the parametrisation of the form factors, as it requires an extrapolation of the differential decay rate to the zero recoil point, where the momentum transfer to the lepton system is maximum.

Recently, the LHCb collaboration has measured  $|V_{cb}|$  using  $B_s^0 \rightarrow D_s^{(*)-}\mu^+\nu_\mu$  decays<sup>1</sup> with two form-factor parametrisations, giving consistent results [7]. The determination of the form factors in  $B_s^0 \rightarrow D_s^{*-}\ell^+\nu_\ell$  decays obtained using different parametrisations can help to clarify the  $|V_{cb}|$  inconsistency between the exclusive and inclusive approaches. It can also be used to improve the Standard Model (SM) predictions of the  $B_s^0 \rightarrow D_s^{*-}\tau^+\nu_\tau$  branching fraction and the ratio  $\mathcal{R}(D_s^*) = \mathcal{B}(B_s^0 \rightarrow D_s^{*-}\tau^+\nu_\tau)/\mathcal{B}(B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu)$ . A measurement and precise prediction of the latter could increase the understanding of the current tension between experimental and theoretical values of the equivalent ratio  $\mathcal{R}(D^{(*)}) = \mathcal{B}(B \rightarrow D^{(*)}\tau^+\nu_\tau)/\mathcal{B}(B \rightarrow D^{(*)}\mu^+\nu_\mu)$  [6]. Theoretical predictions on  $B_s^0$  semileptonic decays are expected to be more precise than those on  $B^0$  or  $B^+$  decays. For example, the Lattice QCD calculations of the form factors are computationally easier due to the larger mass of the spectator  $s$  quark compared to that of  $u$  or  $d$  quarks [8, 9]. Despite these advantages, the study of semileptonic  $B_s^0$  decays has received less theoretical attention than the equivalent  $B^0$  and  $B^+$  decays due to the lack of experimental results.


This paper reports the first measurement of the shape of the differential decay rate of the  $B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu$  decay as a function of the hadronic recoil parameter  $w = v_{B_s^0} \cdot v_{D_s^{*-}}$ , where  $v_{B_s^0}$  and  $v_{D_s^{*-}}$  are the four-vector velocities of the  $B_s^0$  and  $D_s^{*-}$  mesons, respectively. The spectrum of  $w$  is unfolded accounting for the detector resolution on  $w$  and corrected for the reconstruction and selection efficiency. The  $D_s^{*-}$  meson is reconstructed in the  $D_s^{*-} \rightarrow D_s^-\gamma$  mode, where the  $D_s^-$  meson subsequently decays via the  $D_s^- \rightarrow \phi(\rightarrow K^+K^-)\pi^-$  or  $D_s^- \rightarrow K^{*0}(\rightarrow \pi^-K^+)K^-$  mode. The data used correspond to an integrated luminosity of  $1.7 \text{ fb}^{-1}$  collected by the LHCb experiment in 2016 at a centre-of-mass energy of 13 TeV.

The  $B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu$  decay is described by four form factors. The most commonly used parametrisations to model these form factors are by Caprini-Lellouch-Neubert (CLN) [10] and by Boyd-Grinstein-Lebed (BGL) [11–13]. This paper also describes how the relevant parameters of these parametrisations can be extracted by fitting the measured differential decay rate.

## 2 Formalism of the $B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu$ decay

The  $B_s^0 \rightarrow D_s^{*-}\mu^+\nu_\mu$  decay, with the subsequent  $D_s^{*-} \rightarrow D_s^-\gamma$  decay, can be described by three angular variables and the squared momentum transfer to the lepton system, defined as  $q^2 = (p_{B_s^0} - p_{D_s^{*-}})^2$ , where  $p_{B_s^0}$  and  $p_{D_s^{*-}}$  are the four-momenta of the  $B_s^0$  and  $D_s^{*-}$  mesons, respectively. The three angular variables, indicated in figure 1, are two helicity angles  $\theta_\mu$  and  $\theta_{D_s}$ , and the angle  $\chi$ . In this figure the direction of the  $z$ -axis is defined in the

<sup>1</sup>The inclusion of charge-conjugate processes is implied throughout this paper.



**Figure 1.** Schematic overview of the  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decay, introducing the angles  $\theta_{D_s}$ ,  $\theta_\mu$  and  $\chi$ .

$B_s^0$  rest frame as  $\hat{z} = \vec{p}_{D_s^{*-}} / |\vec{p}_{D_s^{*-}}|$ . The angle between the muon direction in the virtual  $W$  rest frame and the  $z$  direction is called  $\theta_\mu$ , while the angle between the  $D_s^{*-}$  meson direction in the  $D_s^{*-}$  rest frame and the  $z$  direction is called  $\theta_{D_s}$ . Finally,  $\chi$  is the angle between the plane formed by the  $D_s^{*-}$  decay products and that formed by the two leptons in the  $B_s^0$  rest frame [14]. The angular basis is designed such that the angular definition for the  $B_s^0$  decay is a  $CP$  transformation of that of the  $B_s^0$  decay.

The measurement is performed by integrating the full decay rate over the decay angles. Thus, the expression of the  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decay rate is given by

$$\frac{d\Gamma(B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu)}{dq^2} = \frac{G_F^2 |V_{cb}|^2 |\eta_{EW}|^2 |\vec{p}| q^2}{96 \pi^3 m_{B_s^0}^2} \left(1 - \frac{m_\mu^2}{q^2}\right)^2 \times \left[ (|H_+|^2 + |H_-|^2 + |H_0|^2) \left(1 + \frac{m_\mu^2}{2q^2}\right) + \frac{3}{2} \frac{m_\mu^2}{q^2} |H_t|^2 \right]. \quad (2.1)$$

In this equation,  $G_F$  is the Fermi constant,  $V_{cb}$  is the CKM matrix element describing the  $b$  to  $c$  transition,  $\eta_{EW} = 1.0066$  is the electroweak correction to  $V_{cb}$  [15],  $m_\mu$  is the muon mass [16], and  $H_0, H_+, H_-, H_t$  are the helicity amplitudes. The magnitude of the  $D_s^{*-}$  momentum in the  $B_s^0$  rest frame is given by  $|\vec{p}|$ . The hadronic recoil,  $w$ , is related to the squared momentum transfer to the lepton pair,  $q^2$ , by

$$w = \frac{p_{B_s^0}}{m_{B_s^0}} \cdot \frac{p_{D_s^{*-}}}{m_{D_s^{*-}}} = \frac{m_{B_s^0}^2 + m_{D_s^{*-}}^2 - q^2}{2 m_{B_s^0} m_{D_s^{*-}}}, \quad (2.2)$$

where  $m_{B_s^0}$  and  $m_{D_s^{*-}}$  are the masses of the  $B_s^0$  and  $D_s^{*-}$  mesons, respectively. The minimal value,  $w = 1$ , corresponds to the situation in which the  $D_s^{*-}$  meson has zero recoil in the  $B_s^0$  rest frame. It is also the value for which  $q^2$  is maximal.

The dependence of the helicity amplitudes on  $w$  can be expressed in different ways, most commonly described in either the CLN or BGL parametrisations, as discussed further

in section 2.1 and section 2.2. This analysis is only sensitive to a single form-factor contribution while the other form factors are fixed to existing measurements from  $B^+$  and  $B^0$  semileptonic decays [6, 17]. This is supported by ref. [18], where when imposing unitarity and analyticity the differences in form factors for semileptonic  $B \rightarrow D$  and  $B_s^0 \rightarrow D_s^+$  decays are found to be within  $\mathcal{O}(1\%)$  over the entire kinematic range. Also, a simultaneous analysis of the  $B_q \rightarrow D_q^{(*)}$  form factors for both light ( $q = u, d$ ) and strange ( $q = s$ ) spectator quarks within the Heavy-Quark-Expansion framework to order  $\mathcal{O}(\alpha_s, 1/m_b, 1/m_c^2)$  [19] does not show any significant SU(3) symmetry breaking. Moreover, Lattice QCD calculations indicate that there is also good agreement of the form factors at zero recoil [9, 20].

## 2.1 CLN form-factor parametrisation

For the CLN parametrisation [10], the helicity amplitudes  $H_0$ ,  $H_+$ ,  $H_-$  and  $H_t$  can be written in terms of the form factors  $A_1(w)$ ,  $V(w)$ ,  $A_2(w)$  and  $A_0(w)$  as

$$\begin{aligned} H_{\pm}(w) &= m_{B_s^0} (1+r) A_1(w) \mp \frac{2}{1+r} |\vec{p}| V(w), \\ H_0(w) &= \frac{m_{B_s^0} m_{D_s^{*-}} (w-r) (1+r)^2 A_1(w) - 2 |\vec{p}|^2 A_2(w)}{m_{D_s^{*-}} (1+r) \sqrt{1+r^2 - 2wr}}, \\ H_t(w) &= \frac{2 |\vec{p}|}{\sqrt{1+r^2 - 2wr}} A_0(w), \end{aligned} \tag{2.3}$$

where  $r = m_{D_s^{*-}}/m_{B_s^0}$ . The form factors are rewritten in terms of a single leading form factor

$$h_{A_1}(w) = A_1(w) \frac{1}{R_{D_s^{*-}}} \frac{2}{w+1}, \tag{2.4}$$

and three ratios of form factors

$$R_0(w) = \frac{A_0(w)}{h_{A_1}(w)} R_{D_s^{*-}}, \quad R_1(w) = \frac{V(w)}{h_{A_1}(w)} R_{D_s^{*-}}, \quad R_2(w) = \frac{A_2(w)}{h_{A_1}(w)} R_{D_s^{*-}}, \tag{2.5}$$

where

$$R_{D_s^{*-}} = \frac{2\sqrt{r}}{1+r}. \tag{2.6}$$

In the CLN parametrisation, the leading form factor and the three ratios are parametrised in terms of  $w$  as

$$\begin{aligned} h_{A_1}(w) &= h_{A_1}(1)[1 - 8\rho^2 z(w) + (53\rho^2 - 15)z^2(w) - (231\rho^2 - 91)z^3(w)], \\ R_0(w) &= R_0(1) - 0.11(w-1) + 0.01(w-1)^2, \\ R_1(w) &= R_1(1) - 0.12(w-1) + 0.05(w-1)^2, \\ R_2(w) &= R_2(1) + 0.11(w-1) - 0.06(w-1)^2, \end{aligned} \tag{2.7}$$

where the coefficients, originally calculated for  $B$  decays, are assumed to be the same for  $B_s^0$  decays. The function  $z(w)$  is defined as

$$z(w) = \frac{\sqrt{w+1} - \sqrt{2}}{\sqrt{w+1} + \sqrt{2}}. \tag{2.8}$$

As this analysis is only sensitive to the shape of the form-factor parametrisation the term  $h_{A_1}(1)$  is absorbed in the normalisation. The values of  $R_1(1)$  and  $R_2(1)$  are taken from the HFLAV average of the corresponding parameters, obtained from  $B \rightarrow D^* \ell \nu_\ell$  decays [6]. The  $R_0(1)$  parameter is suppressed by  $m_\ell^2/q^2$  in the helicity amplitude and its contribution to the total rate is negligible. The value predicted by the exact heavy quark limit of  $R_0(1) = 1$  [21] is used, as no measurement of  $R_0(1)$  has been performed. The slope,  $\rho^2$ , of  $h_{A_1}(w)$  is the only parameter fitted in this parametrisation.

## 2.2 BGL form-factor parametrisation

In the BGL parametrisation [11–13], the helicity amplitudes are parametrised as

$$\begin{aligned}
 H_0(w) &= \frac{\mathcal{F}_1(w)}{m_{B_s^0} \sqrt{1+r^2+2wr}}, \\
 H_\pm(w) &= f(w) \mp m_{B_s^0} m_{D_s^{*-}} \sqrt{w^2-1} g(w), \\
 H_t(w) &= m_{B_s^0} \frac{\sqrt{r(1+r)} \sqrt{w^2-1}}{\sqrt{1+r^2-2wr}} \mathcal{F}_2(w),
 \end{aligned} \tag{2.9}$$

where the form factors are defined as

$$\begin{aligned}
 f(z) &= \frac{1}{P_{1+}(z) \phi_f(z)} \sum_{n=0}^{\infty} a_n^f z^n, & \mathcal{F}_1(z) &= \frac{1}{P_{1+}(z) \phi_{\mathcal{F}_1}(z)} \sum_{n=0}^{\infty} a_n^{\mathcal{F}_1} z^n, \\
 g(z) &= \frac{1}{P_{1-}(z) \phi_g(z)} \sum_{n=0}^{\infty} a_n^g z^n, & \mathcal{F}_2(z) &= \frac{\sqrt{r}}{(1+r) P_{0-}(z) \phi_{\mathcal{F}_2}(z)} \sum_{n=0}^{\infty} a_n^{\mathcal{F}_2} z^n.
 \end{aligned} \tag{2.10}$$

The functions  $\phi_i$  are the so-called outer functions,  $P_{1\pm,0-}$  are Blaschke factors, and the coefficients  $a_n^i$ , where  $i = \{f, g, \mathcal{F}_1, \mathcal{F}_2\}$ , are parameters to be fit from data.

As the form-factor parametrisation is given through analytic functions, they must satisfy the unitarity condition in the  $z$  expansion

$$\sum_{n=0}^{\infty} (a_n^g)^2 \leq 1, \quad \sum_{n=0}^{\infty} (a_n^f)^2 + \sum_{n=0}^{\infty} (a_n^{\mathcal{F}_1})^2 \leq 1, \quad \sum_{n=0}^{\infty} (a_n^{\mathcal{F}_2})^2 \leq 1. \tag{2.11}$$

This analysis is only sensitive to the form factor  $f(z)$ , and its series is truncated at  $n = 2$ , following refs. [17, 22–25]. The shapes for  $\mathcal{F}_1(z)$  and  $g(z)$  are constrained using the results in ref. [17], where the  $a_n^i$  coefficients are fit using recent Belle measurements with  $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$  decays [26, 27]. The value of  $a_0^f$  in ref. [17] is determined from the combination of lattice calculations in ref. [28]. The parameters  $a_n^{\mathcal{F}_2}$  for  $\mathcal{F}_2(z)$  are fixed from predictions in ref. [24], where they are called  $P_1$ . As this analysis represents the first measurement of form factors in  $B_s^0 \rightarrow D_s^{*-}$  transitions, the choice of the input parameters is driven by having as much experimental input as possible. An overview of the fit inputs is given in table 8 in appendix C.

## 3 Detector and simulation

The LHCb detector [29, 30] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$

quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region [31], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [32] placed downstream of the magnet. The tracking system provides a measurement of the momentum,  $p$ , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ $c$ . The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of  $(15 + 29/p_T)$   $\mu\text{m}$ , where  $p_T$  is the component of the momentum transverse to the beam, in GeV/ $c$ . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [33]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [34]. The online event selection is performed by a trigger [35], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. The hardware muon trigger selects events containing a high- $p_T$  muon candidate. The software trigger requires three tracks with a significant displacement from any primary  $pp$  interaction vertex.

Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. In the simulation,  $pp$  collisions are generated using PYTHIA [36, 37] with a specific LHCb configuration [38]. Decays of unstable particles are described by EVTGEN [39], in which final-state radiation is generated using PHOTOS [40]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [41, 42] as described in ref. [43].

The simulation is corrected for mismodeling of the kinematic properties of the generated  $B_s^0$  mesons and of the photons from the  $D_s^{*-}$  decays, as well as for data-simulation differences in the muon trigger efficiency and tracking efficiencies of the final-state particles. Corrections to the  $B_s^0$  and  $\gamma$  kinematic distributions are determined by comparing data and simulated samples of  $B^+ \rightarrow J/\psi K^+$  and  $B_s^0 \rightarrow D_s^{*-} \pi^+$  decays, respectively. Kinematic differences between  $B_s^0$  and  $B^+$  mesons due to their production mechanisms are small and considered to be negligible [44, 45]. Corrections to the trigger and tracking efficiencies are evaluated using data and simulated samples of  $B^+ \rightarrow J/\psi K^+$  decays [46]. In the simulated signal sample, the form factors are described following the CLN parametrisation with numerical values  $\rho^2 = 1.205$ ,  $R_1(1) = 1.404$  and  $R_2(1) = 0.854$ .

## 4 Data selection

Candidate  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decays are selected by pairing  $D_s^{*-}$  and  $\mu^+$  candidates, where the  $D_s^{*-}$  candidate is reconstructed through the  $D_s^- \gamma$  decay. The  $D_s^-$  mesons are reconstructed requiring two opposite-sign kaons and a pion inconsistent with coming from a PV, and forming a common vertex that is displaced from every PV. The final-state hadrons and muon must satisfy strict particle identification (PID) criteria, consistent with the assigned particle hypothesis.

To suppress the combinatorial background in the  $D_s^-$  mass spectrum, only the regions of the  $D_s^- \rightarrow K^+K^-\pi^-$  Dalitz plot compatible with originating from the  $\phi\pi^-$  and  $K^{*0}K^-$  decay modes are retained by requiring the  $K^+K^-$  mass to be within  $20 \text{ MeV}/c^2$  of the known  $\phi$  mass, or the reconstructed  $K^+\pi^-$  mass to be within  $90 \text{ MeV}/c^2$  of the average  $K^*(892)^0$  mass [16]. Possible backgrounds arising from the misidentification of one of the  $D_s^-$  decay products are removed with explicit vetoes which apply more stringent PID requirements in a small window of invariant mass of the corresponding particle combination. The main contributions that are removed come from  $\bar{A}_c^- \rightarrow K^+\bar{p}\pi^-$ ,  $D^- \rightarrow K^+\pi^-\pi^-$ ,  $D_s^- \rightarrow K^-\pi^+\pi^-$ , and misidentified or partially reconstructed multibody  $D$  decays, all originating from semileptonic  $b$ -hadron decays.

Due to the small mass difference between the  $D_s^{*-}$  and  $D_s^-$  mesons, the photon must be emitted close to the  $D_s^-$  flight direction. Photons are selected inside a narrow cone surrounding the  $D_s^-$  candidate, defined in pseudorapidity and azimuthal angle. Only the highest  $p_T$  photon inside the cone is combined with the  $D_s^-$  candidate. Potential contamination from neutral pions reconstructed as a single merged cluster in the electromagnetic calorimeter is suppressed by employing a neural network classifier trained to separate  $\pi^0$  mesons from photons [47].

A fit to the  $D_s^-\gamma$  invariant-mass distribution, with the reconstructed  $D_s^-$  mass constrained to the known value [16], is performed as shown in figure 2. The signal is described by a Gaussian function with a power-law tail on the right hand side of the distribution and the background by an exponential distribution. The power-law tail accounts for cases where additional activity in the calorimeter is mistakenly included in the photon cluster. The *sPlot* technique [48] is employed to subtract the combinatorial background from random photons. Weighted signal is used to create the templates described in section 5. The correlation between the weights and  $w$  is below 4%.

The muon candidate is required to have  $p_T$  in excess of  $1.2 \text{ GeV}/c$ . Background arising from  $b$ -hadrons decaying into final states containing two charmed hadrons,  $H_b \rightarrow D_s^{*-}H_c$ , followed by a semileptonic decay of the charmed hadron  $H_c \rightarrow \mu^+\nu_\mu X$ , where  $X$  is one or more hadrons, are suppressed by using a multivariate algorithm based on the isolation of the muon [49]. Finally, the  $B_s^0$  meson candidates are formed by combining  $\mu^+$  and  $D_s^{*-}$  candidates which are consistent with coming from a common vertex.

## 5 Signal yield


The signal yield is determined using a template fit to the distribution of the corrected mass [50],

$$m_{\text{corr}} = \sqrt{m_{D_s^{*-}\mu^+}^2 + |p_\perp|^2} + |p_\perp|, \quad (5.1)$$

where  $m_{D_s^{*-}\mu^+}$  is the measured mass of the  $D_s^{*-}\mu^+$  candidate, and  $p_\perp$  is the momentum of the candidate transverse to the  $B_s^0$  flight direction. When only one massless final-state particle is missing from the decay,  $m_{\text{corr}}$  peaks at the  $B_s^0$  mass. Only candidates in the range  $3500 < m_{\text{corr}} < 5367 \text{ MeV}/c^2$  are considered.

Extended binned maximum-likelihood fits to the  $m_{\text{corr}}$  distribution are performed independently in seven bins of the reconstructed hadronic recoil,  $w$ , to obtain the raw yields





**Figure 2.** Distribution of the reconstructed  $D_s^- \gamma$  mass,  $m(D_s^- \gamma)$ , with the fit overlaid. The fit is performed constraining the  $D_s^-$  mass to the world-average value [16]. The signal and background components are shown separately with dashed red and dotted green lines, respectively.

bin	1	2	3	4	5	6	7
$w$	1.1087	1.1688	1.2212	1.2717	1.3226	1.3814	1.4667

**Table 1.** Binning scheme used for this measurement. Only the upper bound for each bin is presented. The lower bound on the first bin corresponds to  $w = 1$ .

$N_{\text{meas}}$  per bin. The binning scheme, detailed in table 1, is chosen such that each  $w$  bin has roughly the same signal yield, based on simulation. Obtaining the value of  $w$  requires the knowledge of the momentum of the  $B_s^0$  meson, which in the decays under study can be solved up to a quadratic ambiguity. By imposing momentum balance against the visible system with respect to the flight direction, and assuming the mass of the  $B_s^0$  meson, the momentum of the  $B_s^0$  meson can be estimated. To resolve the ambiguity in the solutions, a multivariate regression algorithm based on the flight direction is used [51] yielding a purity on the solutions of around 70%. The  $m_{\text{corr}}$  distribution is fitted using shapes (templates) of signal and of background distributions mostly obtained from simulation. These simulated decays are selected as described in section 4, and are corrected for the simulation mismodeling as described in section 3.

The largest contribution to the background is due to  $B_s^0 \rightarrow D_s^{*-} \tau^+ \nu_\tau$  decays, with  $\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau$ . A small source of background is formed by excited  $D_s^-$  mesons decaying into a  $D_s^{*-}$  resonance. The only such excited state is the  $D_{s1}(2460)^-$  meson, and hence templates for  $B_s^0 \rightarrow D_{s1}(2460)^- \mu^+ \nu_\mu$  and  $B_s^0 \rightarrow D_{s1}(2460)^- \tau^+ \nu_\tau$  decays are included in the fit. The background arising from  $b$  hadrons decaying into final states containing two charmed hadrons,  $H_b \rightarrow D_s^{*-} H_c$ , is also addressed. The template for this process is gener-

ated using simulated events of  $B_s^0$ ,  $B^0$ ,  $B^+$  and  $A_b^0$  decays, with an appropriate admixture of final states, based on their production rates, branching ratios and relative reconstruction efficiencies taken from simulation. The last background considered in the fit is the combinatorial background, arising from random combinations of tracks. This template is obtained from a data sample where the  $D_s^-$  meson and the muon have the same charge.

The free parameters in the fit are the signal yield, the relative abundances of  $B_s^0 \rightarrow D_s^{*-} \tau^+ \nu_\tau$  and  $B_s^0 \rightarrow D_{s1}(2460)^- \mu^+ \nu_\mu$  candidates with respect to that of the signal, and the fraction of combinatorial background. The total fraction of backgrounds from  $H_c \rightarrow \mu^+ \nu_\mu X$  decays is fixed to the expected value using the measured branching fractions and selection efficiencies obtained from simulation. A 40% uncertainty is assigned to this component to account for the uncertainties on the branching fractions [16]. The  $B_s^0 \rightarrow D_{s1}(2460)^- \tau^+ \nu_\tau$  contribution is also fixed assuming a value of its ratio with respect to the muonic mode equal to the SM prediction for  $\mathcal{B}(B^+ \rightarrow D^{*0} \tau^+ \nu_\tau) / \mathcal{B}(B^+ \rightarrow D^{*0} \mu^+ \nu_\mu)$  [21] under the assumption that this ratio is identical for  $B_s^0$  meson decays. The contribution of this decay to the fit is negligible. The Barlow-Beeston “lite” technique [52, 53] is applied to account for the limited size of the simulation samples. The distributions of  $m_{\text{corr}}$  with the fit overlaid are shown in figure 3.


Using the fractions obtained from the fit, data and simulated distributions of the angular variables  $\cos(\theta_\mu)$ ,  $\cos(\theta_{D_s})$ , and  $\chi$ , as defined in section 2, are shown in figure 4. All distributions show good agreement between data and simulation, indicating that integrating over the angles does not introduce biases.

## 6 Efficiency correction


This analysis requires a precise measurement of all contributions to the efficiency as a function of the true value of the hadronic recoil  $w_{\text{true}}$  extracted from simulation. However, the overall normalisation of the efficiency is not determined as only its dependency with  $w_{\text{true}}$  is relevant.

The total efficiency is the product of the geometrical acceptance of the detector, the efficiency of reconstructing all tracks, the trigger requirements, and the full set of kinematic, PID and background rejection requirements. Most of the contributions to the total efficiency are obtained using simulation. Only the particle identification and the  $D_s^-$  selection efficiencies are derived from data using control samples. The muon and hadron PID efficiencies are taken from large data samples of  $J/\psi \rightarrow \mu^+ \mu^-$  and  $D^{*+} \rightarrow D^0 \pi^+$  decays, respectively [54]. These samples are then used to determine the PID efficiencies in bins of  $p$ ,  $p_T$  and number of tracks in the event. The  $D_s^-$  selection efficiency accounts for selecting the regions in the Dalitz plot, as well as the vetoes described in section 4. This efficiency is determined from a sample of fully reconstructed  $B_s^0 \rightarrow D_s^- \pi^+$  decays as a function of the  $D_s^-$   $p_T$ . The efficiencies extracted from data are convolved with the simulation to obtain their dependency on  $w_{\text{true}}$ .

The efficiencies derived from simulation are extracted by comparing the generator-level simulation, based on PYTHIA [36, 37] and EVTGEN [39], to the final reconstructed



**Figure 3.** Distribution of the corrected mass,  $m_{\text{corr}}$ , for the seven bins of  $w$ , overlaid with the fit results. The  $B_s^0 \rightarrow D_{s1}(2460)^- \tau^+ \nu_\tau$  and the  $B_s^0 \rightarrow D_{s1}(2460)^- \mu^+ \nu_\mu$  components are combined in  $B_s^0 \rightarrow D_{s1}(2460)^- \ell^+ \nu_\ell$ . Below each plot, differences between the data and fit are shown, normalised by the uncertainty in the data.



**Figure 4.** Distribution of (top right)  $\chi$ , (bottom left)  $\cos(\theta_{D_s})$  and (bottom right)  $\cos(\theta_\mu)$  integrating over  $w$  and the other decay angles from data (black points) compared to the distribution from simulation with their relative size extracted from the fit to the corrected mass. The  $B_s^0 \rightarrow D_{s1}(2460)^- \tau^+ \nu_\tau$  and the  $B_s^0 \rightarrow D_{s1}(2460)^- \mu^+ \nu_\mu$  components are combined in  $B_s^0 \rightarrow D_{s1}(2460)^- \ell^+ \nu_\ell$ . The uncertainties on the templates, indicated by the hashed areas in the figures, are a combination from all templates.

and selected simulation sample used for the template fit, omitting the particle identification and the  $D_s^-$  selection criteria.

## 7 Unfolded yields

The measured  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  spectrum from section 5 must be unfolded to account for the resolution on the  $w$  variable, which is 0.07. The unfolding procedure uses a migration matrix determined from simulation, defined as the probability that a candidate generated in bin  $j$  of the  $w_{\text{true}}$  distribution appears in bin  $i$  of the  $w$  distribution. The unfolded spectrum is then corrected bin-by-bin using the efficiency described in section 6. The combination of the migration matrix and the total efficiency, called the response matrix, is shown in appendix B.

The unfolding procedure adopted is based on the singular value decomposition (SVD) method [55] using the RooUnfold package implemented in the Root package [56]. The SVD method includes a regularisation procedure that depends upon a parameter  $k$ , ranging between unity and the number of degrees of freedom, seven in this case. Using simulation, the optimal value for  $k$  is found to be  $k = 5$ , which minimises the difference between the

yield from the unfolding procedure and the expected yield in each bin. The final yields, labelled  $N_{\text{corr}}^{\text{unf}}$ , are normalised to unity and presented in table 2.

### 7.1 Systematic uncertainties

Systematic uncertainties on  $N_{\text{corr}}^{\text{unf}}$  originate from the fitted  $D_s^{*-}$  and  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  yields, and the efficiency corrections. By varying the determination of the unfolded yields, systematic uncertainties are quantified. Since this analysis is sensitive only to the shape of the decay distribution and the absolute normalisation is unknown, every such variation is normalised to unity. After normalising, the values are compared to those from the default normalised unfolded yields, and from this the uncertainties are extracted.

The size of the simulated samples, which are very CPU intensive to generate, is the dominating systematic uncertainty on the unfolded yields. The simulated sample size is accounted for in the fit by applying the Barlow-Beeston “lite” technique [52, 53] when determining the signal yield. Its relative contribution to the systematic uncertainty is assessed by not applying this technique and comparing the obtained uncertainties. The uncertainties due to the size of the control samples used to determine the efficiencies and corrections are obtained by varying each of the efficiency and correction inputs within their uncertainty, repeating this 1000 times, and taking the spread as the uncertainty on  $N_{\text{corr}}^{\text{unf}}$ .

The uncertainty on the SVD unfolding procedure is determined by repeating the regularisation procedure with a different regularisation parameter,  $k$ . The nominal value used is  $k = 5$ , which is changed to  $k = 4$  and  $k = 6$ , and the difference with the nominal value is assigned as the systematic uncertainty.

Two systematic uncertainties are determined to account for assumptions in the simulation. Radiative corrections simulated by the PHOTOS package are known to be incomplete [40, 57]. The difference in  $N_{\text{corr}}^{\text{unf}}$  from simulated samples with and without PHOTOS is evaluated and a third of the difference is assigned following ref. [58]. The efficiency due to the detector acceptance, and thus the shape of the efficiency correction, may be affected by the form factors in the HQET model used to generate the simulation, which are based on the 2016 HFLAV averages [59]. This is studied by weighting both the generator level and fully reconstructed simulated samples to the 2019 HFLAV averages [6]:  $\rho^2 = 1.122 \pm 0.024$ ,  $R_1(1) = 1.270 \pm 0.026$ , and  $R_2(1) = 0.852 \pm 0.018$ , with correlations  $\text{corr}[\rho^2, R_1(1)] = -0.824$ ,  $\text{corr}[\rho^2, R_2(1)] = 0.566$ , and  $\text{corr}[R_1(1), R_2(1)] = -0.715$ . The values of each pair are varied within one standard deviation of their mean, taking into account their correlation. The value of  $R_0(1)$  is varied by a 20% uncertainty accounting for finite  $b$ - and  $c$ -quark masses [21]. These variations result in small changes of the total efficiency and the average difference is taken as the uncertainty.

The effect of the  $B_s^0$  and  $\gamma$  kinematic corrections is assessed by changing the kinematic binning schemes in which the corrections are evaluated. The large effect induced by this change has been checked for statistical fluctuations of the calibration samples. The sample is split randomly into two, after which new corrections and  $N_{\text{corr}}^{\text{unf}}$  yields are calculated. No relevant differences between the  $N_{\text{corr}}^{\text{unf}}$  values of these two samples are found in any  $w$  bin. Hence, the systematic uncertainty is based on the change of binning schemes alone.

	$w$ bin						
	1	2	3	4	5	6	7
Fraction of $N_{\text{corr},i}^{\text{unf}}$	0.183	0.144	0.148	0.128	0.117	0.122	0.158
Uncertainties (%)							
Simulation sample size	3.5	3.0	2.8	3.1	3.4	3.0	3.7
Sample sizes for effs and corrections	3.6	3.2	3.0	2.8	2.8	2.7	2.8
SVD unfolding regularisation	0.5	0.5	0.1	0.7	1.2	0.0	0.5
Radiative corrections	0.1	0.2	0.1	0.3	0.4	0.2	0.2
Simulation FF parametrisation	0.3	0.1	0.1	0.1	0.2	0.4	0.2
Kinematic corrections	2.4	1.0	1.1	0.1	0.2	0.1	0.9
Hardware-trigger efficiency	0.3	0.3	0.0	0.2	0.2	0.3	0.1
Software-trigger efficiency	0.0	0.1	0.0	0.0	0.1	0.0	0.0
$D_s^-$ selection efficiency	0.5	0.2	0.3	0.3	0.2	0.1	0.3
Photon background subtraction	0.0	2.3	0.8	2.9	2.0	0.9	0.4
Total systematic uncertainty	5.6	5.1	4.4	5.2	5.0	4.2	4.8
Statistical uncertainty	3.4	2.9	2.7	3.1	3.2	2.9	3.4

**Table 2.** Fraction of the unfolded yields corrected for the global efficiencies,  $N_{\text{corr}}^{\text{unf}}$ , for each  $w$  bin. Also shown in this table is the breakdown of the systematic and statistical uncertainties on  $N_{\text{corr}}^{\text{unf}}$ . These are shown as a fraction of the unfolded yield.

The corrections to the hardware and software trigger efficiencies applied to the simulated samples depend on the kinematics and PID of the candidates. The systematic uncertainty is evaluated by changing the binning scheme and the PID selection of the control sample.

The systematic uncertainty due to the kinematic dependence of the  $D_s^-$  selection efficiency is assessed by extracting the efficiency as a function of  $p$  instead of  $p_T$  from the  $B_s^0 \rightarrow D_s^- \pi^+$  control sample.

The systematic uncertainty due to the photon background subtraction, performed through the *sPlot* method with fits to the  $D_s^{*-}$  invariant mass, is assessed by implementing the fit with a third-order Chebyshev polynomial for the background description, and repeating the background subtraction process.

Systematic uncertainties induced by the tracking corrections, detector occupancy and PID efficiencies are found to be negligible as they do not affect the corrected mass distribution nor the shape of the efficiency correction.

## 7.2 Results

The  $N_{\text{corr}}^{\text{unf}}$  yields and corresponding systematic and statistical uncertainties per  $w$  bin are shown in table 2. The correlations between the  $N_{\text{corr}}^{\text{unf}}$  yields including statistical and systematic uncertainties are given in table 3, and the covariance matrix is presented in table 6 in

$w$ bin	1	2	3	4	5	6	7
1	1						
2	0.44	1					
3	0.13	0.60	1				
4	0.19	0.32	0.48	1			
5	0.30	0.30	0.15	0.60	1		
6	0.34	0.38	0.33	0.22	0.54	1	
7	0.27	0.34	0.34	0.27	0.07	0.32	1

**Table 3.** Correlation matrix for the unfolded data set in bins of  $w$ , including both statistical and systematic uncertainties.

appendix B. The detector response combined with the reconstruction efficiency is presented in appendix B. Together these can be used to constrain form-factor parametrisations.

### 8 Form factor fits

The yields  $N_{\text{corr}}^{\text{unf}}$  with corresponding correlation matrix presented in section 7 can be fit using various form-factor parametrisations. Fits using the commonly used CLN and BGL parametrisations, with the assumptions described in section 2, are presented in the following.

The values of the form-factor parameters are derived from a  $\chi^2$  fit with

$$\chi^2 = \sum_{i,j} \left( N_{\text{corr},i}^{\text{unf}} - N_{\text{exp},i} \right) C_{ij}^{-1} \left( N_{\text{corr},j}^{\text{unf}} - N_{\text{exp},j} \right). \tag{8.1}$$

In this expression,  $N_{\text{corr},i(j)}^{\text{unf}}$  is the normalised, unfolded and efficiency-corrected yield in bin  $i(j)$ ,  $N_{\text{exp},i(j)}$  is the expected yield in bin  $i(j)$  obtained from integrating  $d\Gamma_{i(j)}/dw$  from the CLN or BGL parametrisation over the bin, and  $C_{ij}$  is the covariance matrix describing the statistical uncertainties from the yields and efficiency corrections. This  $\chi^2$  function is minimised for the CLN and BGL parametrisations separately. For the CLN parametrisation, the fitted value is  $\rho^2 = 1.16 \pm 0.05$ , where the uncertainty is only statistical in nature.

For the BGL parametrisation, the unitarity constraint is considered in the minimisation by adding a Gaussian penalty function [60] to the  $\chi^2$  defined in eq. (8.1). This function is of the form

$$\theta(U - 1) \left( \frac{U - 1}{\sigma} \right)^2, \tag{8.2}$$

where  $\theta$  is the Heaviside function,  $U$  is the unitarity constraint  $\sum_{n=0}^2 (a_n^f)^2 + \sum_{n=0}^2 (a_n^{\mathcal{F}1})^2$ , and  $\sigma$  is the theoretical uncertainty associated with the bound [61]. The correlation between the external parameter  $a_0^f$  and the fitted parameters  $a_1^f$  and  $a_2^f$  is not considered due to its small uncertainty. To assess the impact of this choice, the value of  $a_0^f$  has been increased (decreased) by  $+1(-1)\sigma$ . The change in the fitted parameters is observed to be negligible

compared with the overall systematic uncertainty which covers for it for any value of the correlation between  $a_0^f$  and the rest of the parameters. This can be explained as  $a_0^f$  only enters as a nuisance parameter in the unitarity bounds, which is the only source of correlation between these parameters. As the two scaled parameters  $a_1^f/a_0^f$  and  $a_2^f/a_0^f$  can be much larger than one (as shown in figure 5) the average magnitude of the correlation diminishes.

The fitted values are  $a_1^f = -0.005 \pm 0.034$ , and  $a_2^f = 1.00_{-0.19}^{+0.00}$ , where the uncertainties are only statistical in nature.

### 8.1 Systematic uncertainties

The systematic uncertainties on the parameters  $\rho^2$ ,  $a_1^f$  and  $a_2^f$  originate from the same sources as those described in section 7.1. Additional systematic uncertainties originate from the external parameters used in the form-factor fits. A summary of all systematic uncertainties for  $\rho^2$ ,  $a_1^f$  and  $a_2^f$  is shown in table 4.

The impact of changes in signal yields or efficiencies has been assessed by repeating the fit with different conditions and comparing the obtained values to the nominal ones. In the  $\chi^2$  fit, the parameters  $R_1(1)$  and  $R_2(1)$  are fixed to the HFLAV averages [6]. The uncertainties on these values are propagated to the CLN fit outcome by changing  $R_1(1)$  and  $R_2(1)$  within one standard deviation from their average, while accounting for the correlation between these values. For the BGL fit, the values of the external parameters of the  $f(z)$ ,  $g(z)$  and  $\mathcal{F}_1(z)$  functions are varied simultaneously within their uncertainty. When the uncertainties are asymmetric the largest is chosen. This process is repeated 1000 times applying the unitarity constraint and the difference between the average of the variations and the nominal value is assigned as a systematic uncertainty.

### 8.2 Results

An analysis to extract the leading parameters of the form factor describing the semileptonic transition  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  has been performed. Using the CLN parametrisation the result obtained is

$$\rho^2 = 1.16 \pm 0.05 \text{ (stat)} \pm 0.07 \text{ (syst)},$$

where the mass of the muon has not been neglected. To compare with other published results, the fit is repeated assuming a massless muon, resulting in a small shift of the central value of the  $\rho^2$  parameter of about 1.5%, as shown in table 5. The world-average value of  $\rho^2$  for the equivalent  $B^0 \rightarrow D^{*+} \mu^- \nu_\mu$  decay is  $\rho^2 = 1.122 \pm 0.015 \text{ (stat)} \pm 0.019 \text{ (syst)}$  [6]. Both values of  $\rho^2$  are consistent within their uncertainties. The measurement is also in agreement with the value obtained in ref. [7],  $\rho^2 = 1.23 \pm 0.17 \text{ (stat)} \pm 0.05 \text{ (syst)} \pm 0.01 \text{ (ext)}$ , where the last uncertainty comes from external inputs. That analysis uses  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decays from an independent data set, and where the photon from the  $D_s^{*-}$  decay is not reconstructed. A comparison with the normalised  $\Delta\Gamma/\Delta w$  spectra inferred from the CLN and BGL parametrisations in ref. [7] gives consistent results with the measured  $w$  spectrum in this paper, as shown in appendix D.



Source	$\sigma(\rho^2)$	$\sigma(a_1^f)$	$\sigma(a_2^f)$
Simulation sample size	0.053	0.036	+0.00 -0.35
Sample sizes for efficiencies and corrections	0.020	0.016	+0.00 -0.15
SVD unfolding regularisation	0.008	0.004	–
Radiative corrections	0.004	–	–
Simulation FF parametrisation	0.007	0.005	–
Kinematic corrections	0.024	0.012	–
Hardware-trigger efficiency	0.001	0.008	–
Software-trigger efficiency	0.004	0.002	–
$D_s^-$ selection efficiency	–	0.008	–
Photon background subtraction	0.002	0.015	–
External parameters in fit	0.024	0.002	+0.00 -0.04
Total systematic uncertainty	0.068	0.046	+0.00 -0.38
Statistical uncertainty	0.052	0.034	+0.00 -0.19


**Table 4.** Summary of the systematic and statistical uncertainties on the parameters  $\rho^2$ ,  $a_1^f$  and  $a_2^f$  from the unfolded CLN and BGL fits. The total systematic uncertainty is obtained by adding the individual components in quadrature.

Using the BGL parametrisation, the results obtained are


$$\begin{aligned}
 a_1^f &= -0.005 \pm 0.034 \text{ (stat)} \pm 0.046 \text{ (syst)}, \\
 a_2^f &= 1.00_{-0.19}^{+0.00} \text{ (stat)}_{-0.38}^{+0.00} \text{ (syst)}.
 \end{aligned}$$

In figure 5, the  $\Delta\chi^2$  contours for the scaled parameters  $a_1^f/a_0^f$  versus  $a_2^f/a_0^f$  are shown; figure 8 in appendix C shows the contours of the unscaled  $a_1^f$  versus  $a_2^f$  parameters. The unitarity constraint results in a non-gaussian distribution of the uncertainty on the  $a_2^f/a_0^f$  parameter. The fits to the differential decay rate using both parametrisations are shown in figure 6. The  $p$ -values are 8.2% and 1.3% for the CLN and BGL parametrisations, respectively. The low  $p$ -values are found to be caused by the third bin in  $w$ , which is higher than expected for both parametrisations. When artificially decreasing the central value of this bin by one standard deviation, the  $p$ -values increase to 69.7% and 8.3% for the CLN and BGL parametrisations, respectively. The low  $p$ -value for the latter fit is explained by the fact that the minimum of the  $\chi^2$  function without the unitarity constraint lies in the region excluded by this constraint.

The prediction of the decay rate can also be transformed to a prediction of the expected normalised event yields taking into account the efficiency and resolution, which then is fit to the experimental spectrum. Both procedures provide similar results with small differences induced by slightly different bin-by-bin correlations shown in table 5.



**Figure 5.**  $\Delta\chi^2$  contours for the scaled parameters  $a_1^f/a_0^f$  versus  $a_2^f/a_0^f$ . The black cross marks the best-fit central value. The solid (dashed) contour encloses the  $\Delta\chi^2 = 2.3$  (6.17) region. The observed shape is due to the applied unitarity condition, see eq. (2.11).



**Figure 6.** Unfolded normalised differential decay rate with the fit superimposed for the CLN parametrisation (green), and BGL (red). The band in the fit results includes both the statistical and systematic uncertainty on the data yields.

CLN fit	
Unfolded fit	$\rho^2 = 1.16 \pm 0.05 \pm 0.07$
Unfolded fit with massless leptons	$\rho^2 = 1.17 \pm 0.05 \pm 0.07$
Folded fit	$\rho^2 = 1.14 \pm 0.04 \pm 0.07$
BGL fit	
Unfolded fit	$a_1^f = -0.005 \pm 0.034 \pm 0.046$
	$a_2^f = 1.00_{-0.19}^{+0.00+0.00}$
Folded fit	$a_1^f = 0.039 \pm 0.029 \pm 0.046$
	$a_2^f = 1.00_{-0.13}^{+0.00+0.00}$


**Table 5.** Results from different fit configurations, where the first uncertainty is statistical and the second systematic.

## 9 Conclusions

In conclusion, this paper presents for the first time the unfolded normalised differential decay rate for  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decays as a function of the recoil parameter  $w$ . The unfolded spectrum as a function of  $w$  with the systematic uncertainty per bin is given in table 2 and the correlations between these bins in table 3. This result allows to constrain  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  form-factor parametrisations. The CLN and BGL form-factor parametrisations have been used to fit the measured spectrum with additional input from  $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$  decays. Both fits give consistent results when compared to data.

## Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (U.S.A.). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (U.S.A.). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).



**Figure 7.** Total efficiency as a function of  $w_{\text{true}}$ , including the acceptance of the LHCb detector as well as the reconstruction and selection efficiencies.

### A Fitted yields and efficiency

Figure 7 shows the total efficiency applied to the unfolded signal yields, as a function of  $w_{\text{true}}$ . It is the combination of the reconstruction and selection efficiencies, including the acceptance of the LHCb detector.

### B Covariance and response matrices

This section contains the information needed to reproduce a form-factor fit. To perform the fit using the unfolded, efficiency-corrected and normalised yields given in table 2, the corresponding covariance matrix with the combined statistical uncertainties is given in table 6.

To transform theoretical predictions into expected signal yields, the response matrix, given in table 7 is needed. This contains the migration matrix (from the true value of  $w$  to the reconstructed one) combined with the reconstruction efficiency. The migration matrix is normalised such that the entries within a given bin of  $w$  sum up to unity. The absolute efficiencies have not been measured for this analysis.

$w$ bin [ $10^{-5}$ ]	1	2	3	4	5	6	7
1	16.10						
2	4.73	7.05					
3	1.21	3.81	5.63				
4	1.87	2.12	2.81	6.10			
5	2.74	1.80	0.78	3.37	5.12		
6	2.42	1.82	1.38	0.98	2.17	3.19	
7	3.24	2.69	2.43	2.02	0.44	1.69	8.95


**Table 6.** Covariance matrix for the unfolded data set in bins of  $w$ , including both statistical and systematic uncertainties in units of  $10^{-5}$ .

$w$	$w_{\text{true}}$						
	1	2	3	4	5	6	7
1	132.0	29.9	11.0	6.1	2.7	2.4	1.0
2	22.4	111.0	36.3	11.1	5.0	3.8	1.4
3	6.0	28.7	109.0	35.9	12.3	6.6	4.8
4	4.6	9.8	27.0	102.0	34.6	12.3	5.7
5	1.4	4.4	8.9	30.3	98.0	33.7	10.3
6	0.8	0.7	5.0	8.5	34.5	97.0	30.9
7	-0.1	0.7	2.2	5.7	11.0	33.5	98.5

**Table 7.** Response matrix, containing the migration from  $w_{\text{true}}$  to  $w$  bins together with the total efficiency in units of  $10^{-4}$ .

BGL parameter	Value
$a_0^f$	$0.01221 \pm 0.00016$
$a_1^{\mathcal{F}1}$	$0.0042 \pm 0.0022$
$a_2^{\mathcal{F}1}$	$-0.069^{+0.041}_{-0.037}$
$a_0^g$	$0.024^{+0.021}_{-0.009}$
$a_1^g$	$0.05^{+0.39}_{-0.72}$
$a_2^g$	$1.0^{+0.0}_{-2.0}$
$a_0^{\mathcal{F}2}$	$0.0595 \pm 0.0093$
$a_1^{\mathcal{F}2}$	$-0.318 \pm 0.170$


**Table 8.** Fit inputs used for the BGL fit, taken from ref. [17] and ref. [24].



**Figure 8.**  $\Delta\chi^2$  contours for the scaled parameters  $a_1^f$  versus  $a_2^f$ . The black cross marks the best-fit central value. The solid (dashed) contour encloses the  $\Delta\chi^2 = 2.3$  (6.17) region. The observed shape is due to the applied unitarity condition, see eq. (2.11).

### C Additional information BGL fit

Table 8 gives an overview of the fit inputs for the BGL fit. Figure 8 shows the unscaled  $a_1^f$  versus  $a_2^f$  contours, equivalent to figure 5.



**Figure 9.** Comparison between the  $w$  spectrum measured in this paper to the normalised  $\Delta\Gamma/\Delta w$  spectra inferred from the CLN and BGL parametrisations in ref. [7].

#### D Comparison with *Phys. Rev. D* 101 (2020) 072004

The  $w$  spectrum measured in this analysis can be compared with the results obtained in ref. [7] where the form-factor parameters of the  $B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu$  decay are measured using a version of the CLN and BGL parametrisations. From this, the normalised  $\Delta\Gamma/\Delta w$  spectrum can be inferred, which is shown in figure 9. The spectrum measured in this paper is consistent with the normalised spectra inferred from both CLN and BGL parametrisations used in ref. [7].

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

#### References

- [1] N. Cabibbo, *Unitary symmetry and leptonic decays*, *Phys. Rev. Lett.* **10** (1963) 531 [[INSPIRE](#)].
- [2] M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[INSPIRE](#)].
- [3] D. Fakirov and B. Stech, *F and D decays*, *Nucl. Phys. B* **133** (1978) 315 [[INSPIRE](#)].
- [4] M. Bauer, B. Stech and M. Wirbel, *Exclusive nonleptonic decays of D, D<sub>s</sub>, and B mesons*, *Z. Phys. C* **34** (1987) 103 [[INSPIRE](#)].
- [5] M. Wirbel, B. Stech and M. Bauer, *Exclusive semileptonic decays of heavy mesons*, *Z. Phys. C* **29** (1985) 637 [[INSPIRE](#)].

- [6] HFLAV collaboration, *Averages of b-hadron, c-hadron, and  $\tau$ -lepton properties as of 2018*, [arXiv:1909.12524](#) [INSPIRE].
- [7] LHCb collaboration, *Measurement of  $|V_{cb}|$  with  $B_s^0 \rightarrow D_s^{(*)-} \mu^+ \nu_\mu$  decays*, *Phys. Rev. D* **101** (2020) 072004 [[arXiv:2001.03225](#)] [INSPIRE].
- [8] C.J. Monahan, H. Na, C.M. Bouchard, G.P. Lepage and J. Shigemitsu,  *$B_s \rightarrow D_s \ell \nu$  form factors and the fragmentation fraction ratio  $f_s/f_d$* , *Phys. Rev. D* **95** (2017) 114506 [[arXiv:1703.09728](#)] [INSPIRE].
- [9] HPQCD collaboration, *Lattice QCD calculation of the  $B_{(s)} \rightarrow D_{(s)}^* \ell \nu$  form factors at zero recoil and implications for  $|V_{cb}|$* , *Phys. Rev. D* **97** (2018) 054502 [[arXiv:1711.11013](#)] [INSPIRE].
- [10] I. Caprini, L. Lellouch and M. Neubert, *Dispersive bounds on the shape of  $B \rightarrow D^{(*)} \ell \nu$  form factors*, *Nucl. Phys. B* **530** (1998) 153 [[hep-ph/9712417](#)] [INSPIRE].
- [11] C. Boyd, B. Grinstein and R.F. Lebed, *Constraints on form-factors for exclusive semileptonic heavy to light meson decays*, *Phys. Rev. Lett.* **74** (1995) 4603 [[hep-ph/9412324](#)] [INSPIRE].
- [12] C. Boyd, B. Grinstein and R.F. Lebed, *Model-independent extraction of  $|V_{cb}|$  using dispersion relations*, *Phys. Lett. B* **353** (1995) 306 [[hep-ph/9504235](#)] [INSPIRE].
- [13] C. Boyd, B. Grinstein and R.F. Lebed, *Model-independent determinations of  $\bar{B} \rightarrow D \ell \nu$ ,  $D^* \ell \nu$  form factors*, *Nucl. Phys. B* **461** (1996) 493 [[hep-ph/9508211](#)] [INSPIRE].
- [14] P. Colangelo and F. De Fazio, *Scrutinizing  $\bar{B} \rightarrow D^*(D\pi)\ell^-\bar{\nu}_\ell$  and  $\bar{B} \rightarrow D^*(D\gamma)\ell^-\bar{\nu}_\ell$  in search of new physics footprints*, *JHEP* **06** (2018) 082 [[arXiv:1801.10468](#)] [INSPIRE].
- [15] A. Sirlin, *Large  $m_W$ ,  $m_Z$  behavior of the  $O(\alpha)$  corrections to semileptonic processes mediated by  $W$* , *Nucl. Phys. B* **196** (1982) 83 [INSPIRE].
- [16] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **98** (2018) 030001 and 2019 updates [INSPIRE].
- [17] P. Gambino, M. Jung and S. Schacht, *The  $V_{cb}$  puzzle: an update*, *Phys. Lett. B* **795** (2019) 386 [[arXiv:1905.08209](#)] [INSPIRE].
- [18] A. Kobach, *Continuity and semileptonic  $B_{(s)} \rightarrow D_{(s)}$  form factors*, *Phys. Lett. B* **809** (2020) 135708 [[arXiv:1910.13024](#)] [INSPIRE].
- [19] M. Bordone, N. Gubernari, D. van Dyk and M. Jung, *Heavy-quark expansion for  $\bar{B}_s \rightarrow D_s^{(*)}$  form factors and unitarity bounds beyond the  $SU(3)_F$  limit*, *Eur. Phys. J. C* **80** (2020) 347 [[arXiv:1912.09335](#)] [INSPIRE].
- [20] E. McLean, C.T.H. Davies, A.T. Lytle and J. Koponen, *Lattice QCD form factor for  $B_s \rightarrow D_s^* \ell \nu$  at zero recoil with nonperturbative current renormalisation*, *Phys. Rev. D* **99** (2019) 114512 [[arXiv:1904.02046](#)] [INSPIRE].
- [21] S. Fajfer, J.F. Kamenik and I. Nišandžić, *On the  $B \rightarrow D^* \tau \bar{\nu}_\tau$  sensitivity to new physics*, *Phys. Rev. D* **85** (2012) 094025 [[arXiv:1203.2654](#)] [INSPIRE].
- [22] D. Bigi, P. Gambino and S. Schacht, *A fresh look at the determination of  $|V_{cb}|$  from  $B \rightarrow D^* \ell \nu$* , *Phys. Lett. B* **769** (2017) 441 [[arXiv:1703.06124](#)] [INSPIRE].
- [23] B. Grinstein and A. Kobach, *Model-independent extraction of  $|V_{cb}|$  from  $\bar{B} \rightarrow D^* \ell \bar{\nu}$* , *Phys. Lett. B* **771** (2017) 359 [[arXiv:1703.08170](#)] [INSPIRE].
- [24] D. Bigi, P. Gambino and S. Schacht,  *$R(D^*)$ ,  $|V_{cb}|$ , and the heavy quark symmetry relations between form factors*, *JHEP* **11** (2017) 061 [[arXiv:1707.09509](#)] [INSPIRE].



- [25] S. Jaiswal, S. Nandi and S.K. Patra, *Updates on extraction of  $|V_{cb}|$  and SM prediction of  $R(D^*)$  in  $B \rightarrow D^* \ell \nu_\ell$  decays*, *JHEP* **06** (2020) 165 [[arXiv:2002.05726](#)] [[INSPIRE](#)].
- [26] BELLE collaboration, *Precise determination of the CKM matrix element  $|V_{cb}|$  with  $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}_\ell$  decays with hadronic tagging at Belle*, [arXiv:1702.01521](#) [[INSPIRE](#)].
- [27] BELLE collaboration, *Measurement of the CKM matrix element  $|V_{cb}|$  from  $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$  at Belle*, *Phys. Rev. D* **100** (2019) 052007 [[arXiv:1809.03290](#)] [[INSPIRE](#)].
- [28] FLAVOUR LATTICE AVERAGING GROUP collaboration, *FLAG review 2019*, *Eur. Phys. J. C* **80** (2020) 113 [[arXiv:1902.08191](#)] [[INSPIRE](#)].
- [29] LHCb collaboration, *The LHCb detector at the LHC*, 2008 *JINST* **3** S08005 [[INSPIRE](#)].
- [30] LHCb collaboration, *LHCb detector performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [[arXiv:1412.6352](#)] [[INSPIRE](#)].
- [31] R. Aaij et al., *Performance of the LHCb vertex locator*, 2014 *JINST* **9** P09007 [[arXiv:1405.7808](#)] [[INSPIRE](#)].
- [32] LHCb OUTER TRACKER GROUP collaboration, *Improved performance of the LHCb outer tracker in LHC run 2*, 2017 *JINST* **12** P11016 [[arXiv:1708.00819](#)] [[INSPIRE](#)].
- [33] LHCb RICH GROUP collaboration, *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [[arXiv:1211.6759](#)] [[INSPIRE](#)].
- [34] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, 2013 *JINST* **8** P02022 [[arXiv:1211.1346](#)] [[INSPIRE](#)].
- [35] R. Aaij et al., *The LHCb trigger and its performance in 2011*, 2013 *JINST* **8** P04022 [[arXiv:1211.3055](#)] [[INSPIRE](#)].
- [36] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)] [[INSPIRE](#)].
- [37] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[INSPIRE](#)].
- [38] I. Belyaev et al., *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [[INSPIRE](#)].
- [39] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [40] P. Golonka and Z. Was, *PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays*, *Eur. Phys. J. C* **45** (2006) 97 [[hep-ph/0506026](#)] [[INSPIRE](#)].
- [41] GEANT4 collaboration, *GEANT4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [42] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [43] M. Clemencic et al., *The LHCb simulation application, Gauss: design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [[INSPIRE](#)].
- [44] LHCb collaboration, *Measurement of the fragmentation fraction ratio  $f_s/f_d$  and its dependence on B meson kinematics*, *JHEP* **04** (2013) 001 [[arXiv:1301.5286](#)] [[INSPIRE](#)].
- [45] LHCb collaboration, *Measurement of  $f_s/f_u$  variation with proton-proton collision energy and B-meson kinematics*, *Phys. Rev. Lett.* **124** (2020) 122002 [[arXiv:1910.09934](#)] [[INSPIRE](#)].

- [46] S. Tolk, J. Albrecht, F. Dettori and A. Pellegrino, *Data driven trigger efficiency determination at LHCb*, Tech. Rep. [LHCb-PUB-2014-039](#), CERN, Geneva, Switzerland (2014).
- [47] M. Calvo Gomez et al., *A tool for  $\gamma/\pi^0$  separation at high energies*, Tech. Rep. [LHCb-PUB-2015-016](#), CERN, Geneva, Switzerland (2015).
- [48] M. Pivk and F.R. Le Diberder, *sPlot: a statistical tool to unfold data distributions*, *Nucl. Instrum. Meth. A* **555** (2005) 356 [[physics/0402083](#)] [[INSPIRE](#)].
- [49] LHCb collaboration, *A precise measurement of the  $B^0$  meson oscillation frequency*, *Eur. Phys. J. C* **76** (2016) 412 [[arXiv:1604.03475](#)] [[INSPIRE](#)].
- [50] LHCb collaboration, *Determination of the quark coupling strength  $|V_{ub}|$  using baryonic decays*, *Nature Phys.* **11** (2015) 743 [[arXiv:1504.01568](#)] [[INSPIRE](#)].
- [51] G. Ciezarek, A. Lupato, M. Rotondo and M. Vesterinen, *Reconstruction of semileptonically decaying beauty hadrons produced in high energy pp collisions*, *JHEP* **02** (2017) 021 [[arXiv:1611.08522](#)] [[INSPIRE](#)].
- [52] R.J. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, *Comput. Phys. Commun.* **77** (1993) 219 [[INSPIRE](#)].
- [53] K. Cranmer et al., *HistFactory: a tool for creating statistical models for use with RooFit and RooStats*, Tech. Rep. [CERN-OPEN-2012-016](#), New York U., New York, NY, U.S.A. (2012).
- [54] L. Anderlini et al., *The PIDCalib package*, Tech. Rep. [LHCb-PUB-2016-021](#), CERN, Geneva, Switzerland (2016).
- [55] A. Höcker and V. Kartvelishvili, *SVD approach to data unfolding*, *Nucl. Instrum. Meth. A* **372** (1996) 469 [[hep-ph/9509307](#)] [[INSPIRE](#)].
- [56] T. Adye, *Unfolding algorithms and tests using RooUnfold*, in *PHYSTAT 2011*, [CERN-2011-006](#), CERN, Geneva, Switzerland (2011), pg. 313 [[arXiv:1105.1160](#)] [[INSPIRE](#)].
- [57] S. Calí, S. Klaver, M. Rotondo and B. Sciascia, *Impacts of radiative corrections on measurements of lepton flavour universality in  $B \rightarrow D\ell\nu_\ell$  decays*, *Eur. Phys. J. C* **79** (2019) 744 [[arXiv:1905.02702](#)] [[INSPIRE](#)].
- [58] BABAR collaboration, *Measurements of the semileptonic decays  $\bar{B} \rightarrow D\ell\bar{\nu}$  and  $\bar{B} \rightarrow D^*\ell\bar{\nu}$  using a global fit to  $DX\ell\bar{\nu}$  final states*, *Phys. Rev. D* **79** (2009) 012002 [[arXiv:0809.0828](#)] [[INSPIRE](#)].
- [59] HFLAV collaboration, *Averages of b-hadron, c-hadron, and  $\tau$ -lepton properties as of summer 2016*, *Eur. Phys. J. C* **77** (2017) 895 [[arXiv:1612.07233](#)] [[INSPIRE](#)].
- [60] M. Bordone, M. Jung and D. van Dyk, *Theory determination of  $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}$  form factors at  $O(1/m_c^2)$* , *Eur. Phys. J. C* **80** (2020) 74 [[arXiv:1908.09398](#)] [[INSPIRE](#)].
- [61] D. Bigi and P. Gambino, *Revisiting  $B \rightarrow D\ell\nu$* , *Phys. Rev. D* **94** (2016) 094008 [[arXiv:1606.08030](#)] [[INSPIRE](#)].

## The LHCb collaboration

R. Aaij<sup>31</sup>, C. Abellán Beteta<sup>49</sup>, T. Ackernley<sup>59</sup>, B. Adeva<sup>45</sup>, M. Adinolfi<sup>53</sup>, H. Afsharnia<sup>9</sup>, C.A. Aidala<sup>80</sup>, S. Aiola<sup>25</sup>, Z. Ajaltouni<sup>9</sup>, S. Akar<sup>64</sup>, P. Albicocco<sup>22</sup>, J. Albrecht<sup>14</sup>, F. Alessio<sup>47</sup>, M. Alexander<sup>58</sup>, A. Alfonso Albero<sup>44</sup>, Z. Aliouche<sup>61</sup>, G. Alkhazov<sup>37</sup>, P. Alvarez Cartelle<sup>60</sup>, A.A. Alves Jr<sup>45</sup>, S. Amato<sup>2</sup>, Y. Amhis<sup>11</sup>, L. An<sup>21</sup>, L. Anderlini<sup>21</sup>, G. Andreassi<sup>48</sup>, A. Andreianov<sup>37</sup>, M. Andreotti<sup>20</sup>, F. Archilli<sup>16</sup>, A. Artamonov<sup>43</sup>, M. Artuso<sup>67</sup>, K. Arzymatov<sup>41</sup>, E. Aslanides<sup>10</sup>, M. Atzeni<sup>49</sup>, B. Audurier<sup>11</sup>, S. Bachmann<sup>16</sup>, M. Bachmayer<sup>48</sup>, J.J. Back<sup>55</sup>, S. Baker<sup>60</sup>, P. Baladron Rodriguez<sup>45</sup>, V. Balagura<sup>11,b</sup>, W. Baldini<sup>20,47</sup>, J. Baptista Leite<sup>1</sup>, A. Baranov<sup>41</sup>, R.J. Barlow<sup>61</sup>, S. Barsuk<sup>11</sup>, W. Barter<sup>60</sup>, M. Bartolini<sup>23,47,h</sup>, F. Baryshnikov<sup>77</sup>, J.M. Basels<sup>13</sup>, G. Bassi<sup>28</sup>, V. Batozskaya<sup>35</sup>, B. Batsukh<sup>67</sup>, A. Battig<sup>14</sup>, A. Bay<sup>48</sup>, M. Becker<sup>14</sup>, F. Bedeschi<sup>28</sup>, I. Bediaga<sup>1</sup>, A. Beiter<sup>67</sup>, L.J. Bel<sup>31</sup>, V. Belavin<sup>41</sup>, S. Belin<sup>26</sup>, V. Bellec<sup>48</sup>, K. Belous<sup>43</sup>, I. Belyaev<sup>38</sup>, G. Bencivenni<sup>22</sup>, E. Ben-Haim<sup>12</sup>, S. Benson<sup>31</sup>, S. Beranek<sup>13</sup>, A. Berezhnoy<sup>39</sup>, R. Bernet<sup>49</sup>, D. Berninghoff<sup>16</sup>, H.C. Bernstein<sup>67</sup>, C. Bertella<sup>47</sup>, E. Bertholet<sup>12</sup>, A. Bertolin<sup>27</sup>, C. Betancourt<sup>49</sup>, F. Betti<sup>19,e</sup>, M.O. Bettler<sup>54</sup>, Ia. Bezshyiko<sup>49</sup>, S. Bhasin<sup>53</sup>, J. Bhom<sup>33</sup>, L. Bian<sup>72</sup>, M.S. Bieker<sup>14</sup>, S. Bifani<sup>52</sup>, P. Billoir<sup>12</sup>, F.C.R. Bishop<sup>54</sup>, A. Bizzeti<sup>21,u</sup>, M. Bjørn<sup>62</sup>, M.P. Blago<sup>47</sup>, T. Blake<sup>55</sup>, F. Blanc<sup>48</sup>, S. Blusk<sup>67</sup>, D. Bobulska<sup>58</sup>, V. Bocci<sup>30</sup>, J.A. Boelhauve<sup>14</sup>, O. Boente Garcia<sup>45</sup>, T. Boettcher<sup>63</sup>, A. Boldyrev<sup>78</sup>, A. Bondar<sup>42,x</sup>, N. Bondar<sup>37</sup>, S. Borghi<sup>61,47</sup>, M. Borisyak<sup>41</sup>, M. Borsato<sup>16</sup>, J.T. Borsuk<sup>33</sup>, S.A. Bouchiba<sup>48</sup>, T.J.V. Bowcock<sup>59</sup>, A. Boyer<sup>47</sup>, C. Bozzi<sup>20</sup>, M.J. Bradley<sup>60</sup>, S. Braun<sup>16</sup>, A. Brea Rodriguez<sup>45</sup>, M. Brodski<sup>47</sup>, J. Brodzicka<sup>33</sup>, A. Brossa Gonzalo<sup>55</sup>, D. Brundu<sup>26</sup>, E. Buchanan<sup>53</sup>, A. Büchler-Germann<sup>49</sup>, A. Buonaura<sup>49</sup>, C. Burr<sup>47</sup>, A. Bursche<sup>26</sup>, A. Butkevich<sup>40</sup>, J.S. Butter<sup>31</sup>, J. Buytaert<sup>47</sup>, W. Byczynski<sup>47</sup>, S. Cadeddu<sup>26</sup>, H. Cai<sup>72</sup>, R. Calabrese<sup>20,g</sup>, L. Calero Diaz<sup>22</sup>, S. Cali<sup>22</sup>, R. Calladine<sup>52</sup>, M. Calvi<sup>24,i</sup>, M. Calvo Gomez<sup>44,m</sup>, P. Camargo Magalhaes<sup>53</sup>, A. Camboni<sup>44,m</sup>, P. Campana<sup>22</sup>, D.H. Campora Perez<sup>47</sup>, A.F. Campoverde Quezada<sup>5</sup>, S. Capelli<sup>24,i</sup>, L. Capriotti<sup>19,e</sup>, A. Carbone<sup>19,e</sup>, G. Carboni<sup>29</sup>, R. Cardinale<sup>23,h</sup>, A. Cardini<sup>26</sup>, I. Carli<sup>6</sup>, P. Carniti<sup>24,i</sup>, K. Carvalho Akiba<sup>31</sup>, A. Casais Vidal<sup>45</sup>, G. Casse<sup>59</sup>, M. Cattaneo<sup>47</sup>, G. Cavallero<sup>47</sup>, S. Celani<sup>48</sup>, R. Cenci<sup>28</sup>, J. Cerasoli<sup>10</sup>, A.J. Chadwick<sup>59</sup>, M.G. Chapman<sup>53</sup>, M. Charles<sup>12,47</sup>, Ph. Charpentier<sup>47</sup>, G. Chatzikonstantinidis<sup>52</sup>, M. Chefdeville<sup>8</sup>, V. Chekalina<sup>41</sup>, C. Chen<sup>3</sup>, S. Chen<sup>26</sup>, A. Chernov<sup>33</sup>, S.-G. Chitic<sup>47</sup>, V. Chobanova<sup>45</sup>, S. Cholak<sup>48</sup>, M. Chruszcz<sup>33</sup>, A. Chubykin<sup>37</sup>, V. Chulikov<sup>37</sup>, P. Ciambone<sup>22</sup>, M.F. Cicala<sup>55</sup>, X. Cid Vidal<sup>45</sup>, G. Ciezarek<sup>47</sup>, P.E.L. Clarke<sup>57</sup>, M. Clemencic<sup>47</sup>, H.V. Cliff<sup>54</sup>, J. Closier<sup>47</sup>, J.L. Cobbledick<sup>61</sup>, V. Coco<sup>47</sup>, J.A.B. Coelho<sup>11</sup>, J. Cogan<sup>10</sup>, E. Cogneras<sup>9</sup>, L. Cojocariu<sup>36</sup>, P. Collins<sup>47</sup>, T. Colombo<sup>47</sup>, A. Comerma-Montells<sup>16</sup>, A. Contu<sup>26</sup>, N. Cooke<sup>52</sup>, G. Coombs<sup>58</sup>, S. Coquereau<sup>44</sup>, G. Corti<sup>47</sup>, C.M. Costa Sobral<sup>55</sup>, B. Couturier<sup>47</sup>, D.C. Craik<sup>63</sup>, J. Crkovská<sup>66</sup>, A. Crocombe<sup>55</sup>, M. Cruz Torres<sup>1,aa</sup>, R. Currie<sup>57</sup>, C.L. Da Silva<sup>66</sup>, E. Dall'Occo<sup>14</sup>, J. Dalseno<sup>45,53</sup>, C. D'Ambrosio<sup>47</sup>, A. Danilina<sup>38</sup>, P. d'Argent<sup>47</sup>, A. Davis<sup>61</sup>, O. De Aguiar Francisco<sup>47</sup>, K. De Bruyn<sup>47</sup>, S. De Capua<sup>61</sup>, M. De Cian<sup>48</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, M. De Serio<sup>18,d</sup>, D. De Simone<sup>49</sup>, P. De Simone<sup>22</sup>, J.A. de Vries<sup>31</sup>, C.T. Dean<sup>66</sup>, W. Dean<sup>80</sup>, D. Decamp<sup>8</sup>, L. Del Buono<sup>12</sup>, B. Delaney<sup>54</sup>, H.-P. Dembinski<sup>15</sup>, A. Dendek<sup>34</sup>, V. Denysenko<sup>49</sup>, D. Derkach<sup>78</sup>, O. Deschamps<sup>9</sup>, F. Desse<sup>11</sup>, F. Dettori<sup>26,f</sup>, B. Dey<sup>7</sup>, A. Di Canto<sup>47</sup>, P. Di Nezza<sup>22</sup>, S. Didenko<sup>77</sup>, H. Dijkstra<sup>47</sup>, V. Dobishuk<sup>51</sup>, A.M. Donohoe<sup>17</sup>, F. Dordei<sup>26</sup>, M. Dorigo<sup>28,y</sup>, A.C. dos Reis<sup>1</sup>, L. Douglas<sup>58</sup>, A. Dovbnya<sup>50</sup>, A.G. Downes<sup>8</sup>, K. Dreimanis<sup>59</sup>, M.W. Dudek<sup>33</sup>, L. Dufour<sup>47</sup>, G. Dujany<sup>12</sup>, P. Durante<sup>47</sup>, J.M. Durham<sup>66</sup>, D. Dutta<sup>61</sup>, M. Dziewiecki<sup>16</sup>, A. Dziurda<sup>33</sup>, A. Dzyuba<sup>37</sup>, S. Easo<sup>56</sup>, U. Egede<sup>69</sup>, V. Egorychev<sup>38</sup>, S. Eidelman<sup>42,x</sup>, S. Eisenhardt<sup>57</sup>, R. Ekelhof<sup>14</sup>, S. Ek-In<sup>48</sup>, L. Eklund<sup>58</sup>, S. Ely<sup>67</sup>, A. Ene<sup>36</sup>, E. Epple<sup>66</sup>, S. Escher<sup>13</sup>, J. Eschle<sup>49</sup>, S. Esen<sup>31</sup>, T. Evans<sup>47</sup>, A. Falabella<sup>19</sup>, J. Fan<sup>3</sup>, Y. Fan<sup>5</sup>, B. Fang<sup>72</sup>, N. Farley<sup>52</sup>, S. Farry<sup>59</sup>, D. Fazzini<sup>11</sup>, P. Fedin<sup>38</sup>, M. Féo<sup>47</sup>, P. Fernandez Declara<sup>47</sup>, A. Fernandez Prieto<sup>45</sup>, F. Ferrari<sup>19,e</sup>,

L. Ferreira Lopes<sup>48</sup>, F. Ferreira Rodrigues<sup>2</sup>, S. Ferreres Sole<sup>31</sup>, M. Ferrillo<sup>49</sup>, M. Ferro-Luzzi<sup>47</sup>, S. Filippov<sup>40</sup>, R.A. Fini<sup>18</sup>, M. Fiorini<sup>20,g</sup>, M. Firlej<sup>34</sup>, K.M. Fischer<sup>62</sup>, C. Fitzpatrick<sup>47</sup>, T. Fiutowski<sup>34</sup>, F. Fleuret<sup>11,b</sup>, M. Fontana<sup>47</sup>, F. Fontanelli<sup>23,h</sup>, R. Forty<sup>47</sup>, V. Franco Lima<sup>59</sup>, M. Franco Sevilla<sup>65</sup>, M. Frank<sup>47</sup>, E. Franzoso<sup>20</sup>, G. Frau<sup>16</sup>, C. Frei<sup>47</sup>, D.A. Friday<sup>58</sup>, J. Fu<sup>25,q</sup>, Q. Fuehring<sup>14</sup>, W. Funk<sup>47</sup>, E. Gabriel<sup>57</sup>, T. Gaintseva<sup>41</sup>, A. Gallas Torreira<sup>45</sup>, D. Galli<sup>19,e</sup>, S. Gallorini<sup>27</sup>, S. Gambetta<sup>57</sup>, Y. Gan<sup>3</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>25</sup>, Y. Gao<sup>4</sup>, M. Garau<sup>26</sup>, L.M. Garcia Martin<sup>46</sup>, P. Garcia Moreno<sup>44</sup>, J. García Pardiñas<sup>49</sup>, B. Garcia Plana<sup>45</sup>, F.A. Garcia Rosales<sup>11</sup>, L. Garrido<sup>44</sup>, D. Gascon<sup>44</sup>, C. Gaspar<sup>47</sup>, R.E. Geertsema<sup>31</sup>, D. Gerick<sup>16</sup>, E. Gersabeck<sup>61</sup>, M. Gersabeck<sup>61</sup>, T. Gershon<sup>55</sup>, D. Gerstel<sup>10</sup>, Ph. Ghez<sup>8</sup>, V. Gibson<sup>54</sup>, A. Gioventù<sup>45</sup>, O.G. Girard<sup>48</sup>, P. Gironella Gironell<sup>44</sup>, L. Giubega<sup>36</sup>, C. Giugliano<sup>20,g</sup>, K. Gizdov<sup>57</sup>, V.V. Gligorov<sup>12</sup>, C. Göbel<sup>70</sup>, E. Golobardes<sup>44,m</sup>, D. Golubkov<sup>38</sup>, A. Golutvin<sup>60,77</sup>, A. Gomes<sup>1,a</sup>, M. Goncerz<sup>33</sup>, P. Gorbounov<sup>38,6</sup>, I.V. Gorelov<sup>39</sup>, C. Gotti<sup>24,i</sup>, E. Govorkova<sup>31</sup>, J.P. Grabowski<sup>16</sup>, R. Graciani Diaz<sup>44</sup>, T. Grammatico<sup>12</sup>, L.A. Granado Cardoso<sup>47</sup>, E. Graugés<sup>44</sup>, E. Graverini<sup>48</sup>, G. Graziani<sup>21</sup>, A. Grecu<sup>36</sup>, L.M. Greeven<sup>31</sup>, R. Greim<sup>31</sup>, P. Griffith<sup>20,g</sup>, L. Grillo<sup>61</sup>, L. Gruber<sup>47</sup>, B.R. Gruber Cazon<sup>62</sup>, C. Gu<sup>3</sup>, M. Guarise<sup>20</sup>, P. A. Günther<sup>16</sup>, X. Guo<sup>71</sup>, E. Gushchin<sup>40</sup>, A. Guth<sup>13</sup>, Y. Guz<sup>43,47</sup>, T. Gys<sup>47</sup>, T. Hadavizadeh<sup>62</sup>, G. Haefeli<sup>48</sup>, C. Haen<sup>47</sup>, S.C. Haines<sup>54</sup>, P.M. Hamilton<sup>65</sup>, Q. Han<sup>7</sup>, X. Han<sup>16</sup>, T.H. Hancock<sup>62</sup>, S. Hansmann-Menzemer<sup>16</sup>, N. Harnew<sup>62</sup>, T. Harrison<sup>59</sup>, R. Hart<sup>31</sup>, C. Hasse<sup>14</sup>, M. Hatch<sup>47</sup>, J. He<sup>5</sup>, M. Hecker<sup>60</sup>, K. Heijhoff<sup>31</sup>, K. Heinicke<sup>14</sup>, A.M. Hennequin<sup>47</sup>, K. Hennessy<sup>59</sup>, L. Henry<sup>46</sup>, J. Heuel<sup>13</sup>, A. Hicheur<sup>68</sup>, D. Hill<sup>62</sup>, M. Hilton<sup>61</sup>, S.E. Hollitt<sup>14</sup>, P.H. Hopchev<sup>48</sup>, J. Hu<sup>16</sup>, J. Hu<sup>71</sup>, W. Hu<sup>7</sup>, W. Huang<sup>5</sup>, X. Huang<sup>72</sup>, W. Hulsbergen<sup>31</sup>, T. Humair<sup>60</sup>, R.J. Hunter<sup>55</sup>, M. Hushchyn<sup>78</sup>, D. Hutchcroft<sup>59</sup>, D. Hynds<sup>31</sup>, P. Ibis<sup>14</sup>, M. Idzik<sup>34</sup>, D. Ilin<sup>37</sup>, P. Ilten<sup>52</sup>, A. Inglessi<sup>37</sup>, K. Ivshin<sup>37</sup>, R. Jacobsson<sup>47</sup>, S. Jakobsen<sup>47</sup>, E. Jans<sup>31</sup>, B.K. Jashal<sup>46</sup>, A. Jawahery<sup>65</sup>, V. Jevtic<sup>14</sup>, F. Jiang<sup>3</sup>, M. John<sup>62</sup>, D. Johnson<sup>47</sup>, C.R. Jones<sup>54</sup>, T.P. Jones<sup>55</sup>, B. Jost<sup>47</sup>, N. Jurik<sup>62</sup>, S. Kandybei<sup>50</sup>, Y. Kang<sup>3</sup>, M. Karacson<sup>47</sup>, J.M. Kariuki<sup>53</sup>, N. Kazeev<sup>78</sup>, M. Kecke<sup>16</sup>, F. Keizer<sup>54,47</sup>, M. Kelsey<sup>67</sup>, M. Kenzie<sup>55</sup>, T. Ketel<sup>32</sup>, B. Khanji<sup>47</sup>, A. Kharisova<sup>79</sup>, S. Kholodenko<sup>43</sup>, K.E. Kim<sup>67</sup>, T. Kirn<sup>13</sup>, V.S. Kirsabom<sup>48</sup>, O. Kitouni<sup>63</sup>, S. Klaver<sup>22</sup>, K. Klimaszewski<sup>35</sup>, S. Koliiev<sup>51</sup>, A. Kondybayeva<sup>77</sup>, A. Konoplyannikov<sup>38</sup>, P. Kopciwicz<sup>34</sup>, R. Kopecna<sup>16</sup>, P. Koppenburg<sup>31</sup>, M. Korolev<sup>39</sup>, I. Kostiuik<sup>31,51</sup>, O. Kot<sup>51</sup>, S. Kotriakhova<sup>37,30</sup>, P. Kravchenko<sup>37</sup>, L. Kravchuk<sup>40</sup>, R.D. Krawczyk<sup>47</sup>, M. Krepis<sup>55</sup>, F. Kress<sup>60</sup>, S. Kretschmar<sup>13</sup>, P. Krokovny<sup>42,x</sup>, W. Krupa<sup>34</sup>, W. Krzemien<sup>35</sup>, W. Kucewicz<sup>33,l</sup>, M. Kucharczyk<sup>33</sup>, V. Kudryavtsev<sup>42,x</sup>, H.S. Kuindersma<sup>31</sup>, G.J. Kunde<sup>66</sup>, T. Kvaratskheliya<sup>38</sup>, D. Lacarrere<sup>47</sup>, G. Lafferty<sup>61</sup>, A. Lai<sup>26</sup>, A. Lampis<sup>26</sup>, D. Lancierini<sup>49</sup>, J.J. Lane<sup>61</sup>, R. Lane<sup>53</sup>, G. Lanfranchi<sup>22</sup>, C. Langenbruch<sup>13</sup>, O. Lantwin<sup>49</sup>, T. Latham<sup>55</sup>, F. Lazzari<sup>28,v</sup>, C. Lazzeroni<sup>52</sup>, R. Le Gac<sup>10</sup>, S.H. Lee<sup>80</sup>, R. Lefèvre<sup>9</sup>, A. Leflat<sup>39</sup>, S. Legotin<sup>77</sup>, O. Leroy<sup>10</sup>, T. Lesiak<sup>33</sup>, B. Leverington<sup>16</sup>, H. Li<sup>71</sup>, L. Li<sup>62</sup>, P. Li<sup>16</sup>, P.-R. Li<sup>5</sup>, X. Li<sup>66</sup>, Y. Li<sup>6</sup>, Y. Li<sup>6</sup>, Z. Li<sup>67</sup>, X. Liang<sup>67</sup>, T. Lin<sup>60</sup>, R. Lindner<sup>47</sup>, P. Ling<sup>71</sup>, V. Lisovskyi<sup>14</sup>, R. Litvinov<sup>26</sup>, G. Liu<sup>71</sup>, H. Liu<sup>5</sup>, S. Liu<sup>6</sup>, X. Liu<sup>3</sup>, D. Loh<sup>55</sup>, A. Loi<sup>26</sup>, J. Lomba Castro<sup>45</sup>, I. Longstaff<sup>58</sup>, J.H. Lopes<sup>2</sup>, G. Loustau<sup>49</sup>, G.H. Lovell<sup>54</sup>, Y. Lu<sup>6</sup>, D. Lucchesi<sup>27,o</sup>, S. Luchuk<sup>40</sup>, M. Lucio Martinez<sup>31</sup>, V. Lukashenko<sup>31</sup>, Y. Luo<sup>3</sup>, A. Lupato<sup>27</sup>, E. Luppi<sup>20,g</sup>, O. Lupton<sup>55</sup>, A. Lusiani<sup>28,t</sup>, X. Lyu<sup>5</sup>, L. Ma<sup>6</sup>, R. Ma<sup>71</sup>, S. Maccolini<sup>19,e</sup>, F. Machefert<sup>11</sup>, F. Maciuc<sup>36</sup>, V. Macko<sup>48</sup>, P. Mackowiak<sup>14</sup>, S. Maddrell-Mander<sup>53</sup>, L.R. Madhan Mohan<sup>53</sup>, O. Maev<sup>37,47</sup>, A. Maevskiy<sup>78</sup>, D. Maisuzenko<sup>37</sup>, M.W. Majewski<sup>34</sup>, S. Malde<sup>62</sup>, B. Malecki<sup>47</sup>, A. Malinin<sup>76</sup>, T. Maltsev<sup>42,x</sup>, H. Malygina<sup>16</sup>, G. Manca<sup>26,f</sup>, G. Mancinelli<sup>10</sup>, R. Manera Escalero<sup>44</sup>, D. Manuzzi<sup>19,e</sup>, D. Marangotto<sup>25,q</sup>, J. Maratas<sup>9,w</sup>, J.F. Marchand<sup>8</sup>, U. Marconi<sup>19</sup>, S. Mariani<sup>21</sup>, C. Marin Benito<sup>11</sup>, M. Marinangeli<sup>48</sup>, P. Marino<sup>48</sup>, J. Marks<sup>16</sup>, P.J. Marshall<sup>59</sup>, G. Martellotti<sup>30</sup>, L. Martinazzoli<sup>47</sup>, M. Martinelli<sup>24,i</sup>, D. Martinez Santos<sup>45</sup>, F. Martinez Vidal<sup>46</sup>, A. Massafferri<sup>1</sup>, M. Materok<sup>13</sup>, R. Matev<sup>47</sup>, A. Mathad<sup>49</sup>, Z. Mathe<sup>47</sup>, V. Matiunin<sup>38</sup>, C. Matteuzzi<sup>24</sup>, K.R. Mattioli<sup>80</sup>, A. Mauri<sup>49</sup>, E. Maurice<sup>11,b</sup>, M. Mazurek<sup>35</sup>, M. McCann<sup>60</sup>, L. McConnell<sup>17</sup>,

T.H. Mcgrath<sup>61</sup>, A. McNab<sup>61</sup>, R. McNulty<sup>17</sup>, J.V. Mead<sup>59</sup>, B. Meadows<sup>64</sup>, C. Meaux<sup>10</sup>, G. Meier<sup>14</sup>, N. Meinert<sup>74</sup>, D. Melnychuk<sup>35</sup>, S. Meloni<sup>24,i</sup>, M. Merk<sup>31</sup>, A. Merli<sup>25</sup>, L. Meyer Garcia<sup>2</sup>, M. Mikhasenko<sup>47</sup>, D.A. Milanes<sup>73</sup>, E. Millard<sup>55</sup>, M.-N. Minard<sup>8</sup>, O. Mineev<sup>38</sup>, L. Minzoni<sup>20,g</sup>, S.E. Mitchell<sup>57</sup>, B. Mitreska<sup>61</sup>, D.S. Mitzel<sup>47</sup>, A. Mödden<sup>14</sup>, A. Mogini<sup>12</sup>, R.A. Mohammed<sup>62</sup>, R.D. Moise<sup>60</sup>, T. Mombächer<sup>14</sup>, I.A. Monroy<sup>73</sup>, S. Monteil<sup>9</sup>, M. Morandin<sup>27</sup>, G. Morello<sup>22</sup>, M.J. Morello<sup>28,t</sup>, J. Moron<sup>34</sup>, A.B. Morris<sup>10</sup>, A.G. Morris<sup>55</sup>, R. Mountain<sup>67</sup>, H. Mu<sup>3</sup>, F. Muheim<sup>57</sup>, M. Mukherjee<sup>7</sup>, M. Mulder<sup>47</sup>, D. Müller<sup>47</sup>, K. Müller<sup>49</sup>, C.H. Murphy<sup>62</sup>, D. Murray<sup>61</sup>, P. Muzzetto<sup>26</sup>, P. Naik<sup>53</sup>, T. Nakada<sup>48</sup>, R. Nandakumar<sup>56</sup>, T. Nanut<sup>48</sup>, I. Nasteva<sup>27</sup>, M. Needham<sup>57</sup>, I. Neri<sup>20,g</sup>, N. Neri<sup>25,q</sup>, S. Neubert<sup>16</sup>, N. Neufeld<sup>47</sup>, R. Newcombe<sup>60</sup>, T.D. Nguyen<sup>48</sup>, C. Nguyen-Mau<sup>48,n</sup>, E.M. Niel<sup>11</sup>, S. Nieswand<sup>13</sup>, N. Nikitin<sup>39</sup>, N.S. Nolte<sup>47</sup>, C. Nunez<sup>80</sup>, A. Oblakowska-Mucha<sup>34</sup>, V. Obraztsov<sup>43</sup>, S. Ogilvy<sup>58</sup>, D.P. O'Hanlon<sup>53</sup>, R. Oldeman<sup>26,f</sup>, C.J.G. Onderwater<sup>75</sup>, J. D. Osborn<sup>80</sup>, A. Ossowska<sup>33</sup>, J.M. Otalora Goicochea<sup>2</sup>, T. Ovsianikova<sup>38</sup>, P. Owen<sup>49</sup>, A. Oyanguren<sup>46</sup>, B. Pagare<sup>55</sup>, P.R. Pais<sup>48</sup>, T. Pajero<sup>28,t</sup>, A. Palano<sup>18</sup>, M. Palutan<sup>22</sup>, Y. Pan<sup>61</sup>, G. Panshin<sup>79</sup>, A. Papanestis<sup>56</sup>, M. Pappagallo<sup>57</sup>, L.L. Pappalardo<sup>20,g</sup>, C. Pappenheimer<sup>64</sup>, W. Parker<sup>65</sup>, C. Parkes<sup>61</sup>, B. Passalacqua<sup>20</sup>, G. Passaleva<sup>21,47</sup>, A. Pastore<sup>18</sup>, M. Patel<sup>60</sup>, C. Patrignani<sup>19,e</sup>, A. Pearce<sup>47</sup>, A. Pellegrino<sup>31</sup>, M. Pepe Altarelli<sup>47</sup>, S. Perazzini<sup>19</sup>, D. Pereima<sup>38</sup>, P. Perret<sup>9</sup>, L. Pescatore<sup>48</sup>, K. Petridis<sup>53</sup>, A. Petrolini<sup>23,h</sup>, A. Petrov<sup>76</sup>, S. Petrucci<sup>57</sup>, M. Petruzzo<sup>25,q</sup>, A. Philippov<sup>41</sup>, L. Pica<sup>28</sup>, B. Pietrzyk<sup>8</sup>, G. Pietrzyk<sup>48</sup>, M. Pili<sup>62</sup>, D. Pinci<sup>30</sup>, J. Pinzino<sup>47</sup>, F. Pisani<sup>19</sup>, A. Piucci<sup>16</sup>, Resmi P.K<sup>10</sup>, V. Placinta<sup>36</sup>, S. Playfer<sup>57</sup>, J. Plews<sup>52</sup>, M. Plo Casasus<sup>45</sup>, F. Polci<sup>12</sup>, M. Poli Lener<sup>22</sup>, M. Poliakov<sup>67</sup>, A. Poluektov<sup>10</sup>, N. Polukhina<sup>77,c</sup>, I. Polyakov<sup>67</sup>, E. Polycarpo<sup>2</sup>, G.J. Pomery<sup>53</sup>, S. Ponce<sup>47</sup>, A. Popov<sup>43</sup>, D. Popov<sup>52</sup>, S. Popov<sup>41</sup>, S. Poslavskii<sup>43</sup>, K. Prasanth<sup>33</sup>, L. Promberger<sup>47</sup>, C. Prouve<sup>45</sup>, V. Pugatch<sup>51</sup>, A. Puig Navarro<sup>49</sup>, H. Pullen<sup>62</sup>, G. Punzi<sup>28,p</sup>, W. Qian<sup>5</sup>, J. Qin<sup>5</sup>, R. Quagliani<sup>12</sup>, B. Quintana<sup>8</sup>, N.V. Raab<sup>17</sup>, R.I. Rabadan Trejo<sup>10</sup>, B. Rachwal<sup>34</sup>, J.H. Rademacker<sup>53</sup>, M. Rama<sup>28</sup>, M. Ramos Pernas<sup>45</sup>, M.S. Rangel<sup>2</sup>, F. Ratnikov<sup>41,78</sup>, G. Raven<sup>32</sup>, M. Reboud<sup>8</sup>, F. Redi<sup>48</sup>, F. Reiss<sup>12</sup>, C. Remon Alepuz<sup>46</sup>, Z. Ren<sup>3</sup>, V. Renaudin<sup>62</sup>, R. Ribatti<sup>28</sup>, S. Ricciardi<sup>56</sup>, D.S. Richards<sup>56</sup>, S. Richards<sup>53</sup>, K. Rinnert<sup>59</sup>, P. Robbe<sup>11</sup>, A. Robert<sup>12</sup>, G. Robertson<sup>57</sup>, A.B. Rodrigues<sup>48</sup>, E. Rodrigues<sup>64</sup>, J.A. Rodriguez Lopez<sup>73</sup>, M. Roehrken<sup>47</sup>, S. Roiser<sup>47</sup>, A. Rollings<sup>62</sup>, V. Romanovskiy<sup>43</sup>, M. Romero Lamas<sup>45</sup>, A. Romero Vidal<sup>45</sup>, J.D. Roth<sup>80</sup>, M. Rotondo<sup>22</sup>, M.S. Rudolph<sup>67</sup>, T. Ruf<sup>47</sup>, J. Ruiz Vidal<sup>46</sup>, A. Ryzhikov<sup>78</sup>, J. Ryzka<sup>34</sup>, J.J. Saborido Silva<sup>45</sup>, N. Sagidova<sup>37</sup>, N. Sahoo<sup>55</sup>, B. Saitta<sup>26,f</sup>, C. Sanchez Gras<sup>31</sup>, C. Sanchez Mayordomo<sup>46</sup>, R. Santacesaria<sup>30</sup>, C. Santamarina Rios<sup>45</sup>, M. Santimaria<sup>22</sup>, E. Santovetti<sup>29,j</sup>, D. Saranin<sup>77</sup>, G. Sarpis<sup>61</sup>, M. Sarpis<sup>16</sup>, A. Sarti<sup>30</sup>, C. Satriano<sup>30,s</sup>, A. Satta<sup>29</sup>, M. Saur<sup>5</sup>, D. Savrina<sup>38,39</sup>, H. Sazak<sup>9</sup>, L.G. Scantlebury Smead<sup>62</sup>, S. Schael<sup>13</sup>, M. Schellenberg<sup>14</sup>, M. Schiller<sup>58</sup>, H. Schindler<sup>47</sup>, M. Schmelling<sup>15</sup>, T. Schmelzer<sup>14</sup>, B. Schmidt<sup>47</sup>, O. Schneider<sup>48</sup>, A. Schopper<sup>47</sup>, H.F. Schreiner<sup>64</sup>, M. Schubiger<sup>31</sup>, S. Schulte<sup>48</sup>, M.H. Schune<sup>11</sup>, R. Schwemmer<sup>47</sup>, B. Sciascia<sup>22</sup>, A. Sciubba<sup>30,k</sup>, S. Sellam<sup>68</sup>, A. Semennikov<sup>38</sup>, A. Sergi<sup>52,47</sup>, N. Serra<sup>49</sup>, J. Serrano<sup>10</sup>, L. Sestini<sup>27</sup>, A. Seuthe<sup>14</sup>, P. Seyfert<sup>47</sup>, D.M. Shangase<sup>80</sup>, M. Shapkin<sup>43</sup>, I. Shchemerov<sup>77</sup>, L. Shchutska<sup>48</sup>, T. Shears<sup>59</sup>, L. Shekhtman<sup>42,x</sup>, Z. Shen<sup>4</sup>, V. Shevchenko<sup>76,77</sup>, E.B. Shields<sup>24,i</sup>, E. Shmanin<sup>77</sup>, J.D. Shupperd<sup>67</sup>, B.G. Siddi<sup>20</sup>, R. Silva Coutinho<sup>49</sup>, L. Silva de Oliveira<sup>2</sup>, G. Simi<sup>27,o</sup>, S. Simone<sup>18,d</sup>, I. Skiba<sup>20,g</sup>, N. Skidmore<sup>16</sup>, T. Skwarnicki<sup>67</sup>, M.W. Slater<sup>52</sup>, J.C. Smallwood<sup>62</sup>, J.G. Smeaton<sup>54</sup>, A. Smetkina<sup>38</sup>, E. Smith<sup>13</sup>, I.T. Smith<sup>57</sup>, M. Smith<sup>60</sup>, A. Snoch<sup>31</sup>, M. Soares<sup>19</sup>, L. Soares Lavra<sup>9</sup>, M.D. Sokoloff<sup>64</sup>, F.J.P. Soler<sup>58</sup>, A. Solovev<sup>37</sup>, I. Solovyev<sup>37</sup>, F.L. Souza De Almeida<sup>2</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>14</sup>, E. Spadaro Norella<sup>25,q</sup>, P. Spradlin<sup>58</sup>, F. Stagni<sup>47</sup>, M. Stahl<sup>64</sup>, S. Stahl<sup>47</sup>, P. Stefko<sup>48</sup>, O. Steinkamp<sup>49</sup>, S. Stemmler<sup>16</sup>, O. Stenyakin<sup>43</sup>, M. Stepanova<sup>37</sup>, H. Stevens<sup>14</sup>, S. Stone<sup>67</sup>, M.E. Stramaglia<sup>48</sup>, M. Straticiu<sup>36</sup>, D. Strelakina<sup>77</sup>, S. Strokov<sup>79</sup>, F. Suljik<sup>62</sup>, J. Sun<sup>26</sup>, L. Sun<sup>72</sup>, Y. Sun<sup>65</sup>, P. Svihra<sup>61</sup>, P.N. Swallow<sup>52</sup>, K. Swientek<sup>34</sup>, A. Szabelski<sup>35</sup>, T. Szumlak<sup>34</sup>,

M. Szymanski<sup>47</sup>, S. Taneja<sup>61</sup>, Z. Tang<sup>3</sup>, T. Tekampe<sup>14</sup>, F. Teubert<sup>47</sup>, E. Thomas<sup>47</sup>, K.A. Thomson<sup>59</sup>, M.J. Tilley<sup>60</sup>, V. Tisserand<sup>9</sup>, S. T’Jampens<sup>8</sup>, M. Tobin<sup>6</sup>, S. Tolik<sup>47</sup>, L. Tomassetti<sup>20,g</sup>, D. Tonelli<sup>28</sup>, D. Torres Machado<sup>1</sup>, D.Y. Tou<sup>12</sup>, E. Tournefier<sup>8</sup>, M. Traill<sup>58</sup>, M.T. Tran<sup>48</sup>, E. Trifonova<sup>77</sup>, C. Trippel<sup>48</sup>, A. Trisovic<sup>54</sup>, A. Tsaregorodtsev<sup>10</sup>, G. Tuci<sup>28,47,p</sup>, A. Tully<sup>48</sup>, N. Tuning<sup>31</sup>, A. Ukleja<sup>35</sup>, D.J. Unverzagt<sup>16</sup>, A. Usachov<sup>31</sup>, A. Ustyuzhanin<sup>41,78</sup>, U. Uwer<sup>16</sup>, A. Vagner<sup>79</sup>, V. Vagnoni<sup>19</sup>, A. Valassi<sup>47</sup>, G. Valenti<sup>19</sup>, M. van Beuzekom<sup>31</sup>, H. Van Hecke<sup>66</sup>, E. van Herwijnen<sup>47</sup>, C.B. Van Hulse<sup>17</sup>, M. van Veghel<sup>75</sup>, R. Vazquez Gomez<sup>44,22</sup>, P. Vazquez Regueiro<sup>45</sup>, C. Vázquez Sierra<sup>31</sup>, S. Vecchi<sup>20</sup>, J.J. Velthuis<sup>53</sup>, M. Veltri<sup>21,r</sup>, A. Venkateswaran<sup>67</sup>, M. Vernet<sup>9</sup>, M. Veronesi<sup>31</sup>, M. Vesterinen<sup>55</sup>, J.V. Viana Barbosa<sup>47</sup>, D. Vieira<sup>64</sup>, M. Vieites Diaz<sup>48</sup>, H. Viemann<sup>74</sup>, X. Vilasis-Cardona<sup>44</sup>, E. Vilella Figueras<sup>59</sup>, P. Vincent<sup>12</sup>, G. Vitali<sup>28</sup>, A. Vitkovskiy<sup>31</sup>, A. Vollhardt<sup>49</sup>, D. Vom Bruch<sup>12</sup>, A. Vorobyev<sup>37</sup>, V. Vorobyev<sup>42,x</sup>, N. Voropaev<sup>37</sup>, R. Waldi<sup>74</sup>, J. Walsh<sup>28</sup>, C. Wang<sup>16</sup>, J. Wang<sup>3</sup>, J. Wang<sup>72</sup>, J. Wang<sup>4</sup>, J. Wang<sup>6</sup>, M. Wang<sup>3</sup>, R. Wang<sup>53</sup>, Y. Wang<sup>7</sup>, Z. Wang<sup>49</sup>, D.R. Ward<sup>54</sup>, H.M. Wark<sup>59</sup>, N.K. Watson<sup>52</sup>, S.G. Weber<sup>12</sup>, D. Websdale<sup>60</sup>, A. Weiden<sup>49</sup>, C. Weisser<sup>63</sup>, B.D.C. Westhenry<sup>53</sup>, D.J. White<sup>61</sup>, M. Whitehead<sup>13</sup>, D. Wiedner<sup>14</sup>, G. Wilkinson<sup>62</sup>, M. Wilkinson<sup>67</sup>, I. Williams<sup>54</sup>, M. Williams<sup>63</sup>, M.R.J. Williams<sup>61</sup>, T. Williams<sup>52</sup>, F.F. Wilson<sup>56</sup>, W. Wislicki<sup>35</sup>, M. Witek<sup>33</sup>, L. Witola<sup>16</sup>, G. Wormser<sup>11</sup>, S.A. Wotton<sup>54</sup>, H. Wu<sup>67</sup>, K. Wyllie<sup>47</sup>, Z. Xiang<sup>5</sup>, D. Xiao<sup>7</sup>, Y. Xie<sup>7</sup>, H. Xing<sup>71</sup>, A. Xu<sup>4</sup>, J. Xu<sup>5</sup>, L. Xu<sup>3</sup>, M. Xu<sup>7</sup>, Q. Xu<sup>5</sup>, Z. Xu<sup>4</sup>, D. Yang<sup>3</sup>, Y. Yang<sup>5</sup>, Z. Yang<sup>3</sup>, Z. Yang<sup>65</sup>, Y. Yao<sup>67</sup>, L.E. Yeomans<sup>59</sup>, H. Yin<sup>7</sup>, J. Yu<sup>7,z</sup>, X. Yuan<sup>67</sup>, O. Yushchenko<sup>43</sup>, K.A. Zarebski<sup>52</sup>, M. Zavertyaev<sup>15,c</sup>, M. Zdybal<sup>33</sup>, O. Zenaiev<sup>47</sup>, M. Zeng<sup>3</sup>, D. Zhang<sup>7</sup>, L. Zhang<sup>3</sup>, S. Zhang<sup>4</sup>, W.C. Zhang<sup>3</sup>, Y. Zhang<sup>47</sup>, A. Zhelezov<sup>16</sup>, Y. Zheng<sup>5</sup>, X. Zhou<sup>5</sup>, Y. Zhou<sup>5</sup>, X. Zhu<sup>3</sup>, V. Zhukov<sup>13,39</sup>, J.B. Zonneveld<sup>57</sup>, S. Zucchelli<sup>19,e</sup>, D. Zuliani<sup>27</sup> and G. Zunica<sup>61</sup>

<sup>1</sup> Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup> Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup> Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup> School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

<sup>5</sup> University of Chinese Academy of Sciences, Beijing, China

<sup>6</sup> Institute Of High Energy Physics (IHEP), Beijing, China

<sup>7</sup> Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

<sup>8</sup> Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

<sup>9</sup> Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>10</sup> Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

<sup>11</sup> Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

<sup>12</sup> LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

<sup>13</sup> I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

<sup>14</sup> Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>15</sup> Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>16</sup> Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>17</sup> School of Physics, University College Dublin, Dublin, Ireland

<sup>18</sup> INFN Sezione di Bari, Bari, Italy

<sup>19</sup> INFN Sezione di Bologna, Bologna, Italy

<sup>20</sup> INFN Sezione di Ferrara, Ferrara, Italy

<sup>21</sup> INFN Sezione di Firenze, Firenze, Italy

<sup>22</sup> INFN Laboratori Nazionali di Frascati, Frascati, Italy

<sup>23</sup> INFN Sezione di Genova, Genova, Italy

<sup>24</sup> INFN Sezione di Milano-Bicocca, Milano, Italy

<sup>25</sup> INFN Sezione di Milano, Milano, Italy

<sup>26</sup> INFN Sezione di Cagliari, Monserrato, Italy

- 27 *Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*
- 28 *INFN Sezione di Pisa, Pisa, Italy*
- 29 *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- 30 *INFN Sezione di Roma La Sapienza, Roma, Italy*
- 31 *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- 32 *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- 33 *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- 34 *AGH — University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- 35 *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- 36 *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- 37 *Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia*
- 38 *Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia*
- 39 *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- 40 *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*
- 41 *Yandex School of Data Analysis, Moscow, Russia*
- 42 *Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- 43 *Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia*
- 44 *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- 45 *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- 46 *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia — CSIC, Valencia, Spain*
- 47 *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- 48 *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- 49 *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- 50 *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- 51 *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- 52 *University of Birmingham, Birmingham, United Kingdom*
- 53 *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- 54 *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- 55 *Department of Physics, University of Warwick, Coventry, United Kingdom*
- 56 *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- 57 *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- 58 *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- 59 *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- 60 *Imperial College London, London, United Kingdom*
- 61 *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- 62 *Department of Physics, University of Oxford, Oxford, United Kingdom*
- 63 *Massachusetts Institute of Technology, Cambridge, MA, United States*
- 64 *University of Cincinnati, Cincinnati, OH, United States*
- 65 *University of Maryland, College Park, MD, United States*
- 66 *Los Alamos National Laboratory (LANL), Los Alamos, United States*
- 67 *Syracuse University, Syracuse, NY, United States*
- 68 *Laboratory of Mathematical and Subatomic Physics, Constantine, Algeria, associated to <sup>2</sup>*
- 69 *School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to <sup>55</sup>*
- 70 *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*

- <sup>71</sup> *Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to* <sup>3</sup>
- <sup>72</sup> *School of Physics and Technology, Wuhan University, Wuhan, China, associated to* <sup>3</sup>
- <sup>73</sup> *Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to* <sup>12</sup>
- <sup>74</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to* <sup>16</sup>
- <sup>75</sup> *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to* <sup>31</sup>
- <sup>76</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to* <sup>38</sup>
- <sup>77</sup> *National University of Science and Technology “MISIS”, Moscow, Russia, associated to* <sup>38</sup>
- <sup>78</sup> *National Research University Higher School of Economics, Moscow, Russia, associated to* <sup>41</sup>
- <sup>79</sup> *National Research Tomsk Polytechnic University, Tomsk, Russia, associated to* <sup>38</sup>
- <sup>80</sup> *University of Michigan, Ann Arbor, United States, associated to* <sup>67</sup>

- <sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*
- <sup>b</sup> *Laboratoire Leprince-Ringuet, Palaiseau, France*
- <sup>c</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>d</sup> *Università di Bari, Bari, Italy*
- <sup>e</sup> *Università di Bologna, Bologna, Italy*
- <sup>f</sup> *Università di Cagliari, Cagliari, Italy*
- <sup>g</sup> *Università di Ferrara, Ferrara, Italy*
- <sup>h</sup> *Università di Genova, Genova, Italy*
- <sup>i</sup> *Università di Milano Bicocca, Milano, Italy*
- <sup>j</sup> *Università di Roma Tor Vergata, Roma, Italy*
- <sup>k</sup> *Università di Roma La Sapienza, Roma, Italy*
- <sup>l</sup> *AGH — University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- <sup>m</sup> *DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- <sup>n</sup> *Hanoi University of Science, Hanoi, Vietnam*
- <sup>o</sup> *Università di Padova, Padova, Italy*
- <sup>p</sup> *Università di Pisa, Pisa, Italy*
- <sup>q</sup> *Università degli Studi di Milano, Milano, Italy*
- <sup>r</sup> *Università di Urbino, Urbino, Italy*
- <sup>s</sup> *Università della Basilicata, Potenza, Italy*
- <sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*
- <sup>u</sup> *Università di Modena e Reggio Emilia, Modena, Italy*
- <sup>v</sup> *Università di Siena, Siena, Italy*
- <sup>w</sup> *MSU — Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*
- <sup>x</sup> *Novosibirsk State University, Novosibirsk, Russia*
- <sup>y</sup> *INFN Sezione di Trieste, Trieste, Italy*
- <sup>z</sup> *Physics and Micro Electronic College, Hunan University, Changsha City, China*
- <sup>aa</sup> *Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras*