## ON THE LERAY-SCHAUDER DEGREE OF THE TODA SYSTEM ON COMPACT SURFACES

ANDREA MALCHIODI<sup>(1)</sup> AND DAVID RUIZ<sup>(2)</sup>

ABSTRACT. In this paper we consider the following Toda system of equations on a compact surface:

$$\begin{pmatrix} -\Delta u_1 = 2\rho_1 (h_1 e^{u_1} - 1) - \rho_2 (h_2 e^{u_2} - 1) \\ -\Delta u_2 = 2\rho_2 (h_2 e^{u_2} - 1) - \rho_1 (h_1 e^{u_1} - 1) \end{pmatrix}$$

Here  $h_1, h_2$  are smooth positive functions and  $\rho_1, \rho_2$  two positive parameters.

In this note we compute the Leray-Schauder degree mod  $\mathbb{Z}_2$  of the problem for  $\rho_i \in (4\pi k, 4\pi (k+1))$   $(k \in \mathbb{N})$ . Our main tool is a theorem of Krasnoselskii and Zabreiko on the degree of maps symmetric with respect to a subspace. This result yields new existence results as well as a new proof of previous results in literature.

## 1. INTRODUCTION

In this paper we consider the following problem on a compact orientable surface  $\Sigma$ :

(1) 
$$\begin{cases} -\Delta u_1 = 2\rho_1 \left( h_1 e^{u_1} - 1 \right) - \rho_2 \left( h_2 e^{u_2} - 1 \right), \\ -\Delta u_2 = 2\rho_2 \left( h_2 e^{u_2} - 1 \right) - \rho_1 \left( h_1 e^{u_1} - 1 \right). \end{cases}$$

Here  $h_1, h_2$  are smooth positive functions and  $\Delta$  is the Laplace-Beltrami operator.

Equation (1) is known as the Toda system, and has been extensively studied in the literature. This problem has a close relationship to geometry, since it describes the integrability of Frenet frames for holomorphic curves in  $\mathbb{CP}^2$  (see [5]). Moreover, it arises in the study of non-abelian Chern-Simons theory in the self-dual case, when a scalar Higgs field is coupled to a gauge potential, see [4, 17, 19].

Let us first discuss the scalar counterpart of (1), namely a Liouville equation in the form:

(2) 
$$-\Delta u = \rho \left(h e^u - 1\right),$$

where  $\rho \in \mathbb{R}$  and h(x) > 0. Equation (2) arises in the prescribed Gaussian curvature problem under conformal deformation of the metric, and also describes the abelian counterpart of (1) from the physical point of view. This equation has been very much studied in the literature; there are by now many results regarding existence, compactness of solutions, bubbling behavior, etc. We refer the interested reader to the reviews [14, 18].

Problem (2) presents a lack of compactness, as its solutions might blow-up. Indeed, take a blowing up sequence  $u_n$  of (1) with  $\rho_n \in \mathbb{R}$  bounded. Then it was proved in [2, 11, 12] that, up to a subsequence

$$\rho_n \to 8k\pi, \ k \in \mathbb{N}.$$

<sup>2000</sup> Mathematics Subject Classification. 35J47, 35J61, 58J20.

Key words and phrases. Geometric PDEs, Leray-Schauder degree.

A.M. is supported by the FIRB project Analysis and Beyond, the PRIN Variational Methods and Nonlinear PDE's and by the University fo Warwick. A. M. and D.R have been supported by the Spanish Ministry of Science and Innovation under Grant MTM2011-26717. D. R. has also been supported by J. Andalucia (FQM 116).

Moreover,  $e^{u_n}$  behaves like the conformal factor of the stereographic projection from  $S^2$  onto  $\mathbb{R}^2$ , composed with a dilation, and located at a finite number of points.

With that result at hand, one can define the Leray-Schauder degree associated to problem (1) and  $\rho \in (8k\pi, 8(k+1)\pi)$ . By the homotopy property of the degree, the latter is independent of the metric g and the function h, and will only vary with k and the topology of  $\Sigma$ . The computation of the degree has been accomplished in [3], where the following formula is given:

(3) 
$$d_{LS} = \frac{1}{k!} (-\chi(\Sigma) + 1) \cdots (-\chi(\Sigma) + k), \quad (\chi(\Sigma) \text{ is the Euler characteristic of } \Sigma).$$

In order to obtain this formula, in [3] a detailed study of all blowing-up solutions and their local degree is performed (see also [13] for a different approach).

Coming back to system (1), it was proved in [6, 8] that the set of solutions is compact for  $(\rho_1, \rho_2) \notin (4\pi \mathbb{N} \times \mathbb{R}) \cup (\mathbb{R} \times 4\pi \mathbb{N})$ . In other words, if blowing up occurs, at least one of the components is quantized.

Therefore, as above, the degree for (1) is well defined for  $(\rho_1, \rho_2)$  away from that set. It is easy to observe that this degree is equal to 1 if both  $\rho_i$  are smaller than  $4\pi$  (one can deform the parameters to  $\rho_1 = \rho_2 = 0$ ). Apart from that, there exists no formula for the Leray-Schauder degree for system (1) yet.

Because of that, most of the existence results for problem (1) have used variational methods so far. Indeed, it has been proved that there exists at least one solution in the following cases:

- (1) for both  $\rho_i < 4\pi$  (see [7]);
- (2) for any  $\rho_1 < 4\pi$ ,  $\rho_2 \in (4k\pi, 4(k+1)\pi)$ ,  $k \in \mathbb{N}$  (see [15]);
- (3) for both  $\rho_i \in (4\pi, 8\pi)$  (see [16]);
- (4) for  $\rho_1 \in (4k\pi, 4(k+1)\pi), \rho_2 \in (4m\pi, 4(m+1)\pi), k, m \in \mathbb{N}$  and  $\Sigma$  with positive genus (see [1]).

In (1), it is proved that the associated energy functional is coercive and hence a minimum is found. The rest of the results use min-max theory, as the functional is no longer bounded from below.

In this note we discuss the parity of the degree for  $\rho_i \in (4n\pi, 4(n+1)\pi)$ , see Proposition 2.2. Our result is consequence of a general theorem (recalled in the next section) concerning the degree of maps symmetric with respect to a subspace, see [9]. We will show that the degree of the Toda system has the same parity as the degree of the scalar case with  $\rho = \rho_i$ , which is given by (3).

In particular, the degree is always odd for  $\rho_i \in (4n\pi, 4(n+1)\pi)$  if n = 1, 2, 3. The case n = 1 implies a new, simpler proof, of the existence result of [16]. The cases n = 2 or 3 yield a new existence result:

**Theorem 1.1.** Assume that  $\Sigma$  is homeomorphic to  $\mathbb{S}^2$ , and  $\rho_i \in (4n\pi, 4(n+1)\pi)$ , with n = 2 or 3. Then there exists a solution to (1).

Acknowledgment: D. R. thanks Rafael Ortega, from the University of Granada, for several discussions on the degree for symmetric maps and his kind help in finding reference [9].

## 2. The parity of the Leray-Schauder degree

The main abstract tool we are going to use is the following one, which is a version of Theorem 21.12 of [9] (page 115).

**Theorem 2.1.** Let P be a continuous linear projection from a Banach space E onto a (closed) subspace  $E^0 \subseteq E$ , and define U = -Id + 2P the reflection with respect to  $E^0$ , which is assumed to be an isometry. Let  $A : E \to E$  be a compact operator equivariant with respect to U, that is

$$AU(x) = UA(x) \quad \forall x \in E.$$

Observe that in particular  $A(E^0) \subset E^0$ . Finally, assume that  $\Phi x = x - Ax$  does not vanish on the boundary of  $B_R = B(0, R)$ . Then,  $\deg(\Phi, B_R, 0)$  and  $\deg(\Phi|_{E^0}, B_R \cap E^0, 0)$  have the same parity.

**Remark 2.1.** Theorem 2.1 is easy to understand if we assume that  $\Phi$  is  $C^1$  and that all its zeroes are non-degenerate. In such case, all zeroes have index  $\pm 1$ , and the total degree is the sum of the indexes of all zeroes. Observe now that if x is a zero of  $\Phi$ , then also Ux is a zero. Moreover x = Ux if and only if  $x \in E^0$ . In other words, the zeroes outside  $E^0$  come in pairs, and give an even contribution to the total degree.

Furthermore, the index of  $x \in E^0$  a zero of  $\Phi$  could be different from its index as a zero of  $\Phi|_{E^0}$ , but in both cases it is  $\pm 1$ . So the difference is even.

The proof of [9] is topological and does not use these arguments.

Let us assume, for simplicity, that  $Vol_g(\Sigma) = 1$ . For  $\alpha \in (0, 1)$  we consider the functional framework

$$E = C_0^{2,\alpha}(\Sigma) \times C_0^{2,\alpha}(\Sigma),$$

where  $C_0^{2,\alpha}(\Sigma)$  stands for the class of  $C^{2,\alpha}$  functions with zero average. Define now the operator  $A = A_{\rho_1,\rho_2}^{h_1,h_2} : E \to E$  as

$$A \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} (-\Delta)^{-1} \left[ 2\rho_1 \left( h_1 \frac{e^{u_1}}{\int_{\Sigma} h_1 e^{u_1} dV_g} - 1 \right) - \rho_2 \left( h_2 \frac{e^{u_2}}{\int_{\Sigma} h_2 e^{u_2} dV_g} - 1 \right) \right] \\ (-\Delta)^{-1} \left[ 2\rho_2 \left( h_2 \frac{e^{u_2}}{\int_{\Sigma} h_2 e^{u_2} dV_g} - 1 \right) - \rho_1 \left( h_1 \frac{e^{u_1}}{\int_{\Sigma} h_1 e^{u_1} dV_g} - 1 \right) \right] \end{pmatrix}.$$

Here by  $(-\Delta)^{-1}f$ ,  $f \in C^{\alpha}(\Sigma)$ , we denote the unique solution u of  $-\Delta u = f$  with zero average. In the above formula solutions exist by Fredholm's theory. Notice also that zeroes of  $\Phi = Id - A$  give rise to solutions of (1). Indeed, it suffices to add proper constants to  $u_1, u_2$  in order to have  $\int_{\Sigma} h_i e^{u_i} dV_g = 1$ .

By elliptic regularity theory the operator A is compact. Moreover, if  $n \in \mathbb{N}$  and  $\rho_1, \rho_2 \in (4n\pi, 4(n+1)\pi)$ , the solutions are a priori bounded, see [6]. Therefore for such values of  $\rho_1, \rho_2$  and for R sufficiently large the degree deg( $\Phi, B_R, 0$ ) is well defined. The main result of this paper is the following:

**Proposition 2.2.** Consider  $n \in \mathbb{N}$  and let  $\rho_1$ ,  $\rho_2 \in (4n\pi, 4(n+1)\pi)$ . Then, for sufficiently large R, deg $(\Phi, B_R, 0)$  has the same parity as

$$d_k = \frac{1}{k!}(-\chi(\Sigma) + 1)\cdots(-\chi(\Sigma) + k),$$

where  $\chi(\Sigma)$  is the Euler characteristic of  $\Sigma$  and k = [n/2].

**PROOF.** Let us consider the homotopy

$$(1-s)(\rho_1,\rho_2) + s(\rho,\rho);$$
  $\rho = \frac{1}{2}(\rho_1 + \rho_2);$ 

$$(1-s)(h_1,h_2) + s(h,h);$$
  $h = \frac{1}{2}(h_1 + h_2).$ 

By the homotopy invariance of the degree we deduce that (for R sufficiently large)

(4) 
$$\deg(\Phi_{\rho_1,\rho_2}^{h_1,h_2}, B_R, 0) = \deg(\Phi_{\rho,\rho}^{h,h}, B_R, 0).$$

Because of that, it suffices to study the degree for  $h_1 = h_2 = h$ ,  $\rho_1 = \rho_2 = \rho$ . We choose  $E^0$  to be the couples of identical functions in E, namely

$$E^{0} = \left\{ \left( \begin{array}{c} u_{1} \\ u_{2} \end{array} \right) \in E : u_{1} = u_{2} \right\},$$

and we define the projection  $P: E \to E^0$  as

$$P\left(\begin{array}{c}u_1\\u_2\end{array}\right) = \frac{1}{2}\left(\begin{array}{c}u_1+u_2\\u_1+u_2\end{array}\right).$$

Observe that the reflection U is given by:

$$U\left(\begin{array}{c}u_1\\u_2\end{array}\right)=\left(\begin{array}{c}u_2\\u_1\end{array}\right).$$

With these definitions, we are in conditions to apply Theorem 2.1, and hence  $\deg(\Phi, B_R, 0)$  has the same parity as  $\deg(\Phi|_{E_0}, B_R \cap E_0, 0)$ .

We now define:

$$T: C_0^{2,\alpha}(\Sigma) \to E_0, \ T(u) = \begin{pmatrix} u \\ u \end{pmatrix}.$$

Clearly T is an homeomorphism, which implies that:

$$\deg(\Phi|_{E_0}, B_R \cap E_0, 0) = \deg(T^{-1} \circ \Phi|_{E_0} \circ T, B_R, 0),$$

where  $\tilde{B}_R = B(0, R) \subset C_0^{2,\alpha}(\Sigma)$ . Observe now that:

$$\tilde{\Phi}(u) := T^{-1} \circ \Phi|_{E_0} \circ T(u) = u - (-\Delta)^{-1} \left[ \rho \left( h \frac{e^u}{\int_{\Sigma} h e^u dV_g} - 1 \right) \right].$$

Moreover,  $\rho \in (4n\pi, 4(n+1)\pi) \subset (8k\pi, 8(k+1)\pi)$  for  $k = \lfloor n/2 \rfloor$ . Finally, deg $(\tilde{\Phi}, \tilde{B}_R, 0)$ has been computed in [3] and it is given by the formula (3). Notice that the Leray-Schauder degree in [3] has been computed in the  $H^1$  setting, but using elliptic regularity theory one can prove that this coincides with the degree in the  $C^{2,\alpha}$ setting, see Theorem B.1 in [10]. This concludes the proof.  $\blacksquare$ 

Observe that if n = 1, then k = 0 and hence the degree is odd. In this way we obtain the existence result of [16] with an alternative approach. There are other cases in which the degree is odd, so we recover some of the results of [1]. In particular, if n=2,3, then k=1 and the degree is odd for any compact orientable  $\Sigma$ . As a consequence, we obtain Theorem 1.1, which gives a new existence result.

## References

- Luca Battaglia, Aleks Jevnikar, Andrea Malchiodi and David Ruiz, A general existence result for the toda system on compact surfaces, preprint arXiv 1306.5404, 2013.
- [2] Haim Brezis and Frank Merle. Uniform estimates and blow-up behavior for solutions of  $-\Delta u = V(x)e^u$  in two dimensions. Comm. Partial Differential Equations, 16(8-9):1223–1253, 1991.
- [3] Chiun-Chuan Chen and Chang-Shou Lin, Topological degree for a mean field equation on Riemann surfaces, Comm. Pure Appl. Math., 56(12):1667-1727, 2003.
- [4] G. Dunne. Self-dual Chern-Simons Theories. Lecture notes in physics. New series m: Monographs. Springer, 1995.
- [5] M. A. Guest, Harmonic maps, loops groups, and integrable systems. London Mathematical Society Student Texts, 38. Cambridge University Press, Cambridge, 1997.
- [6] Jürgen Jost, Changshou Lin, and Guofang Wang. Analytic aspects of the Toda system. II. Bubbling behavior and existence of solutions. Comm. Pure Appl. Math., 59(4):526–558, 2006.
- [7] Jürgen Jost and Guofang Wang. Analytic aspects of the Toda system. I. A Moser-Trudinger inequality. Comm. Pure Appl. Math., 54(11):1289–1319, 2001.
- [8] Jürgen Jost and Guofang Wang. Classification of solutions of a Toda system in ℝ<sup>2</sup>. Int. Math. Res. Not., (6):277–290, 2002.
- [9] M. A. Krasnoselskii and P. P. Zabreiko, Geometrical methods of nonlinear anal- ysis, volume 263 Fundamental Principles of Mathematical Sciences. Springer-Verlag, Berlin, 1984. Translated from the Russian by Christian C. Fenske.
- [10] Yan Yan Li, Prescribing scalar curvature on  $S^n$  and related problems. I., J. Differential Equations, 120(2):319-410, 1995.
- [11] Yan Yan Li. Harnack type inequality: the method of moving planes. Comm. Math. Phys., 200(2):421-444, 1999.
- [12] Yan Yan Li and Itai Shafrir. Blow-up analysis for solutions of  $-\Delta u = Ve^u$  in dimension two. Indiana Univ. Math. J., 43(4):1255–1270, 1994.
- [13] Andrea Malchiodi. Morse theory and a scalar field equation on compact surfaces. Adv. Differential Equations, 13(11-12):1109–1129, 2008.
- [14] Andrea Malchiodi. Topological methods for an elliptic equation with exponential nonlinearities. Discrete Contin. Dyn. Syst., 21(1):277–294, 2008.
- [15] Andrea Malchiodi and Cheikh Birahim Ndiaye. Some existence results for the Toda system on closed surfaces. Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., 18(4):391–412, 2007.
- [16] Andrea Malchiodi and David Ruiz. A variational analysis of the Toda system on compact surfaces. Comm. Pure Appl. Math., 66(3):332–371, 2013.
- [17] Gabriella Tarantello. Selfdual gauge field vortices. Progress in Nonlinear Differential Equations and their Applications, 72. Birkhäuser Boston Inc., Boston, MA, 2008. An analytical approach.
- [18] Gabriella Tarantello. Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. Discrete Contin. Dyn. Syst., 28(3):931–973, 2010.
- [19] Yisong Yang. Solitons in field theory and nonlinear analysis. Springer Monographs in Mathematics. Springer-Verlag, New York, 2001.

<sup>(1)</sup> UNIVERSITY OF WARWICK, MATHEMATICS INSTITUTE, ZEEMAN BUILDING, COVENTRY CV4 7AL AND SISSA, VIA BONOMEA 265, 34136 TRIESTE (ITALY).

<sup>(2)</sup> Departamento de Análisis Matemático, University of Granada, 18071 Granada (Spain).

E-mail address: A.Malchiodi@warwick.ac.uk, malchiod@sissa.it, daruiz@ugr.es