



Supplement of

Variational techniques for a one-dimensional energy balance model

Gianmarco Del Sarto et al.

Correspondence to: Gianmarco Del Sarto (gianmarco.delsarto@sns.it)

The copyright of individual parts of the supplement might differ from the article licence.

Introduction

We are providing additional supplementary material to accompany our manuscript. Specifically, we present rigorous proofs for the results outlined in Section 3 of the paper. This supplementary material is organized as follows. In Section S1, we recall the definition of the functional associated with the elliptic problem arising from the study of the stationary solutions of the EBM depending on an additive positive parameter q representing the carbon dioxide concentration. In Section S3, we rigorously prove the existence of a global minimiser for the functional using the direct method from the calculus of variations. Furthermore, we establish the regularity, non-negativity, and boundedness of the minimiser. In Section S4, we characterize the uniqueness of the global minimiser in terms of the value function, i.e., the minimum, depending on q , among the values of the functional. Further, we show that the derivative of the value function is, up to the sign, the global mean temperature of the minimiser of the variational problem. Then, in Section S5 we prove that the value function is not only semiconcave but also concave. In particular, this last property implies that the global mean temperature of the variational problem minimiser is non-decreasing with respect to q . Lastly, in Section S6 we provide sufficient conditions to prove the existence of three steady-state solutions for the EBM. This relies on the use of the Mountain Pass Theorem and the direct method from the calculus of variations.

S1 Functional definition

The 1D-EBM we consider through the main paper has the form ([2]):

$$\begin{aligned} C_T \partial_t u &= \partial_x (\kappa(x) u_x) + Q_0(x) \beta(u) + q - \varepsilon_0 \sigma_0 u^4, \quad (t, x) \in [0, T] \times [-1, 1] \\ u_x(t, -1) &= u_x(t, 1) = 0, \quad t \geq 0, \\ u(0, x) &= \tilde{u}(x), \end{aligned}$$

where \tilde{u} denotes the initial condition. The steady-state solutions to the previous problem are associated with the functional ([3, 7]):

$$F_q(u) = \int_{-1}^1 \varepsilon_0 \sigma_0 \frac{(u^5)}{5} - Q_0(x) B(u) - qu \, dx + \frac{1}{2} \int_{-1}^1 \kappa(x) [u'(x)]^2 \, dx.$$

In the main manuscript, we have illustrated some properties of the minimiser of the variational problem

$$\inf \{ F_q(u) \mid u \in H^1, u \geq 0 \}.$$

But we can prove more. Indeed, we will extend F_q to a functional defined on H^1 , and then shows that there exists $u_0 \in H^1$ s.t. $u_0 \geq 0$ and

$$\tilde{F}_q(u_0) = \inf \{ \tilde{F}_q(u) \mid u \in H^1 \} = \inf \{ F_q(u) \mid u \in H^1, u \geq 0 \}.$$

The necessity to extend F_q to take into account also negative values for u comes from the fact that the natural space in which set the minimization problem is the Sobolev space $H^{1,2}(-1, 1)$. But, due to the presence of the odd polynomial term in F_q , we have

$$\inf \{F_q(u) \mid u \in H^1\} = -\infty.$$

In fact, choosing the constant function $u_\lambda \equiv -\lambda$, with $\lambda > 0$, we get $\lim_{\lambda \rightarrow +\infty} F_q(u_\lambda) = -\infty$. This is not surprising, since the term u^5 inside the functional F_q comes from the Stefan-Boltzmann law, which has no sense for negative values of the temperature. Further, in order to pick a primitive B of the co-albedo β , we need to extend the definition of the co-albedo also for the negative values of the Kelvin temperature. In any case, these extensions do not really affect the EBM, since are referred to negative values of u , which have no physical sense.

Let's turn to give details about the extensions of β and the Stefan-Boltzmann law. Let $\tilde{\beta}: \mathbb{R} \rightarrow \mathbb{R}$ be s.t. $\tilde{\beta} \in C^\infty$ and

- (i) $\tilde{\beta}$ is monotonically increasing
- (ii) $\tilde{\beta}(u) \geq 0 \forall u \in \mathbb{R}$
- (iii) $\tilde{\beta}(u) = 0$ for $u \leq -M$, for some $M > 0$
- (iv) $\tilde{\beta}$ extends β for $u \geq 0$.

Denote by $B(u) := B_0 + \int_0^u \tilde{\beta}(v)dv$ a primitive of β , with B_0 s.t. $B(u) \geq 0 \forall u \in \mathbb{R}$. It will be useful in future to note that

$$0 \leq B(u) \leq |B_0| + |u| \quad \forall u. \quad (1)$$

In a similar way, we consider the extension of the Stefan-Boltzmann law given by:

$$\psi(u) = \begin{cases} \varepsilon_0 \sigma_0 u^4, & \text{if } u \geq 0 \\ 0, & \text{if } u < 0. \end{cases}$$

A primitive is given by $\Psi(u) = \varepsilon_0 \sigma_0 \frac{(u^5)_+}{5}$, where $(x)_+ = \max(x, 0)$.

Lastly, since in our model we are assuming κ continuous and positive on $[-1, 1]$, we can also assume κ constant. Indeed, all the proofs, that in this manuscript are carried with constant κ , extended immediately to the non-constant case. In conclusion, the uniformly elliptic equation we are considering is given by:

$$\begin{aligned} 0 &= \kappa \Delta u + Q_0(x) \tilde{\beta}(u) + q - \psi(u), \\ u'(-1) &= u'(1) = 0, \end{aligned} \quad (2)$$

and the functional associated with its solution is

$$\tilde{F}_q(u) = \int_{-1}^1 \Psi(u) - Q_0(x)B(u) - qu \, dx + \frac{\kappa}{2} \int_{-1}^1 [u'(x)]^2 \, dx.$$

S2 Gibbs invariant measure and functional minimum point

In this section, we make rigorous the relation between F_q and the invariant measure ν of the stochastic EBM. In particular, we prove a result giving information about the concentration of ν around minimum points for F_q . We start by recalling the notation and some useful results.

First, we set

$$I(u) := \int_{-1}^1 \left(\varepsilon_0 \sigma_0 \frac{(u^5)_+}{5} - Q_0(x)B(u) - qu \right) dx.$$

Consider $H = L^2(-1, 1)$ and $E = C([-1, 1])$. Then, following the theory of stochastic partial differential equation (SPDE), the stochastic equation obtained by adding a cylindrical Brownian motion is a gradient SPDE of the form

$$dX_t = [AX_t + f(x, X_t)] dt + \varepsilon dW_t, \quad X|_{t=0} = x_0 \quad (3)$$

where $(W_t)_t$ is a cylindrical Wiener process on H and $A = \kappa \Delta$ is the Neumann Laplacian with constant viscosity $\kappa > 0$, i.e. $A: D(A) \subset H \rightarrow H$,

$$\begin{aligned} D(A) &= \{u \in H^2(-1, 1) \mid u'(-1) = u'(1) = 0\} \\ Au &= \kappa u'', \end{aligned} \quad (4)$$

and

$$f(x, u) = Q_0(x)\tilde{\beta}(u) + q - \varepsilon_0 \sigma_0 (u^4)_+$$

We refer to [5] for details about the properties of the previous SPDE. The mild solution X_t of (3) is \mathbb{P} -a.s. valued in E . Further, applying the theory of invariant measure developed in [8], we get the following property.

Proposition S1 *The SPDE (3) has a unique Gibbs invariant measure ν . Further, $\nu \ll \mu$ with explicit formula:*

$$\nu(du) = \frac{1}{Z} \exp\left(-\frac{2}{\varepsilon^2} I(u)\right) \mu(du), \quad u \in H \quad (5)$$

where $\mu \sim \mathcal{N}\left(0, -\frac{\varepsilon^2}{2} A^{-1}\right)$ is a Gaussian measure on H .

Remark S2 *The Neumann Laplacian Δ is not invertible on $H = L^2(-1, 1)$ and the invariant measure theory applies for a strictly negative definite operator A on H . For this reason, we should consider the strictly negative operator*

$$\tilde{A} := \lambda Id - A, \quad \lambda > 0.$$

In this way, the functional takes the form:

$$\tilde{I}(u) = I(u) - \lambda \|u\|_2^2,$$

and the reaction term in the SPDE is given by:

$$\tilde{f}(x, u) = f(x, u) - \lambda u.$$

In conclusion, the invariant measure for the SPDE

$$d\tilde{X}_t = \left[\tilde{A}\tilde{X}_t + \tilde{f}(x, X_t) \right] dt + \varepsilon dW_t, \quad \tilde{X}_{|t=0} = \tilde{x}_0$$

is given by:

$$\tilde{\nu}(du) = \frac{1}{Z} \exp\left(-\frac{2}{\varepsilon^2} \tilde{I}(u)\right) \tilde{\mu}(du), \quad u \in H,$$

where $\tilde{\mu} \sim \mathcal{N}(0, -\frac{\varepsilon^2}{2} \tilde{A}^{-1})$ is a Gaussian measure on H . Since this change only complicates the notation in the proofs, we will keep writing A^{-1} but the reader should interpret the Laplacian with the shift described above, in order to get the rigorous meaning.

Second, keeping in mind the previous remark, we adopt from now on the notation:

$$\mathcal{Q} = -\frac{\varepsilon^2}{2\kappa} \Delta^{-1}, \quad \mu(du; v, \mathcal{Q}) \sim \mathcal{N}(v, \mathcal{Q}).$$

The following statement is a classical result about the equivalence of Gaussian measures. See [8] for more details.

Theorem S3 (Cameron-Martin) *The Gaussian measures $\mu(du; 0, \mathcal{Q})$ and $\mu(du; v, \mathcal{Q})$ on H are equivalent if and only if $v \in \mathcal{Q}^{1/2}(H)$. In this case:*

$$\frac{\mu(du; 0, \mathcal{Q})}{\mu(du; v, \mathcal{Q})} = \exp\left(-\langle \mathcal{Q}^{-1/2}u, \mathcal{Q}^{-1/2}v \rangle + \frac{1}{2} \left\| \mathcal{Q}^{-1/2}v \right\|_2^2\right). \quad (6)$$

In the following, we are going to recall the rigorous meaning for

$$\langle \mathcal{Q}^{-1/2}u, \mathcal{Q}^{-1/2}v \rangle_{L^2}.$$

Consider

$$W_z: \mathcal{Q}^{1/2}(H) \subset H \rightarrow L^2(H, \mu), \quad W_z(u) := \langle u, \mathcal{Q}^{-1/2}z \rangle_{L^2}.$$

It can be shown that:

- (i) W_z is an isometry,
- (ii) $\mathcal{Q}^{1/2}(H)$ is dense in H (here it is fundamental $\ker(\mathcal{Q}) = \{0\}$.)

In this way, W_z can be extended in a unique way to a map $W_z: H \rightarrow L^2(H, \mu)$. So, it should be interpreted as:

$$\langle \mathcal{Q}^{-1/2}u, \mathcal{Q}^{-1/2}v \rangle_{L^2} = W_{\mathcal{Q}^{-1/2}v}(u), \quad u \in H.$$

Remark S4 For our choice of the operator \mathcal{Q} , we have that the Cameron-Martin space is $\mathcal{Q}^{1/2}(L^2) = H^1$.

At this point, we move to prove the main result of this section. Given a Banach space X , we denote by

$$B_X(x_0, \rho) = \{x \in X \mid \|x - x_0\|_X < \rho\}$$

the open ball with center $x_0 \in X$ and radius $\rho > 0$.

Proposition S5 Let $C > 0$, $r > 5$ and $v \in H^{2,2}(-1, 1)$. Consider the set

$$B_C(v, \eta) := B_{L^2}(v, \eta) \cap B_{L^r}(v, C).$$

Then,

$$(i) \nu(B_C(v, \eta)) \xrightarrow{C \rightarrow +\infty} \nu(B_{L^2}(v, \eta))$$

$$(ii) \mu(B_C(0, \eta)) \xrightarrow{C \rightarrow +\infty} \mu(B_{L^2}(0, \eta))$$

(iii) For each $C > 0$, it holds

$$\frac{\nu(B_C(v, \eta))}{\mu(B_C(0, \eta))} = \frac{1}{Z} \exp\left(-\frac{2}{\varepsilon^2} \left(\tilde{F}_q(v) + O(\eta^\theta)\right)\right),$$

where $\theta \in (0, 1)$ satisfies

$$\frac{1}{5} = \frac{\theta}{2} + \frac{1 - \theta}{r}.$$

Proof. Assume for simplicity $\kappa = 1$.

(i)-(ii) Observe that for $v_1 \in L^2 \cap L^r$, we have:

$$B_{C_1}(v_1, \eta) \subset B_{C_2}(v_1, \eta), \quad \text{if } C_1 \leq C_2$$

and

$$B_{L^2}(v_1, \eta) = \left(\bigcup_{C>0} B_C(v_1, \eta) \right) \cup \{u \in L^2 \mid \|u\|_r = \infty\}.$$

Denote by $B = \{u \in L^2 \mid \|u\|_r = \infty\}$. If we are able to prove:

$$\mu(B) = 0,$$

then we get (i) and (ii) thanks to the continuity of measures on an increasing sequence of sets. Since $\mu \sim \mathcal{N}(0, -\frac{\varepsilon^2}{2} \Delta^{-1})$, then

$$\mu = M \sum_{n=1}^{\infty} \frac{Z_n}{n} e_n$$

where $\{Z_n\}_n$ are i.i.d. $\mathcal{N}(0, 1)$ defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, $\{e_n\}_n$ is an orthonormal basis of $H = L^2(-1, 1)$ and $M > 0$ is a constant. The previous series is convergent in $L^2((\Omega, \mathcal{F}, \mathbb{P}); H^s)$, for all $s < 1/2$; indeed

$$\|\mu\|_{H^s}^2 = \left\| M \sum_n \frac{Z_n}{n} (-\Delta)^s e_n \right\|_2^2 = \left\| M \sum_n \frac{Z_n}{n} n^s e_n \right\|_2^2 = M^2 \sum_n M \frac{Z_n^2}{n^2} n^{2s} = \sum_n \frac{Z_n^2}{n^{2-2s}}.$$

Taking the expected values, we get

$$\mathbb{E} \|\mu\|_{H^s}^2 = M^2 \sum_n \frac{1}{n^{2-2s}} < \infty \leftrightarrow s < 1/2.$$

The Sobolev embedding $H^s(-1, 1) \hookrightarrow L^p$, holds for each $p < \infty$ if s is close to $1/2$. This leads to $\mu(B) = 0$.

(iii) We start by using the explicit formula (5) in order to get

$$\frac{\nu(B_C(v, \eta))}{\mu(B_C(0, \eta))} = \frac{1}{Z} \frac{\int_{B_C(v, \eta)} \exp\left(-\frac{2}{\varepsilon^2} I(u)\right) \mu(du; 0, \mathcal{Q})}{\mu(B_C(0, \eta); 0, \mathcal{Q})}. \quad (7)$$

Using the Cameron-Martin formula (6), we have

$$\mu(du; 0, \mathcal{Q}) = \exp\left(-W_{\mathcal{Q}^{-1/2}v} + \frac{1}{2} \|\mathcal{Q}^{-1/2}v\|_2^2\right) \mu(du; v, \mathcal{Q})$$

for each $v \in Q^{1/2}(L^2) = H^1$. Since

$$\mathcal{Q}^{-1/2}u = \frac{\sqrt{2}}{\varepsilon} (-\Delta)^{1/2}u,$$

and for $v \in H^2$

$$W_{\mathcal{Q}^{-1/2}v}(u) = \langle u, \mathcal{Q}^{-1/2}\mathcal{Q}^{-1/2}v \rangle = \frac{2}{\varepsilon^2} \langle u, v'' \rangle$$

we arrive to

$$\mu(du; 0, \mathcal{Q}) = \exp\left(-\frac{2}{\varepsilon^2} \langle u', v' \rangle + \frac{1}{\varepsilon^2} \|v'\|_2^2\right) \mu(du; v, \mathcal{Q}).$$

Plugging the previous identity into (7), we deduce:

$$\frac{\nu(B_C(v, \eta))}{\mu(B_C(0, \eta))} = \frac{1}{Z} \frac{\int_{B_C(v, \eta)} \exp\left[-\frac{2}{\varepsilon^2} \left(I(u) + \langle u', v' \rangle - \frac{1}{2} \|v'\|_2^2\right)\right] \mu(du; v, \mathcal{Q})}{\mu(B_C(0, \eta); 0, \mathcal{Q})}$$

Assume for a moment that we are able to prove:

$$-\frac{2}{\varepsilon^2} \left(I(u) + \langle u', v' \rangle - \frac{1}{2} \|v'\|_2^2\right) = -\frac{2}{\varepsilon^2} \left(\tilde{F}_q(v) + O(\eta^\theta)\right), \quad u \in B_C(v, \eta), \quad (8)$$

for $\theta \in (0, 1)$. Then,

$$\begin{aligned} \frac{\nu(B_C(v, \eta))}{\mu(B_C(0, \eta))} &= \frac{1}{Z} \exp\left(-\frac{2}{\varepsilon^2} \left(\tilde{F}_q(v) + O(\eta^\theta)\right)\right) \frac{\mu(B_C(v, \eta); v, \mathcal{Q})}{\mu(B_C(0, \eta); 0, \mathcal{Q})} \\ &= \frac{1}{Z} \exp\left(-\frac{2}{\varepsilon^2} \left(\tilde{F}_q(v) + O(\eta^\theta)\right)\right), \end{aligned}$$

where we have used

$$\mu(B_C(v, \eta); v, \mathcal{Q}) = \mu(B_C(0, \eta); 0, \mathcal{Q}).$$

This concludes the proof. ■

It remains to prove (8).

Lemma S6 *If $u \in B_C(v, \eta)$ and $v \in H^2$ then there exists $\theta \in (0, 1)$ s.t.*

$$-\frac{2}{\varepsilon^2} \left(I(u) + W_{\mathcal{Q}^{-1/2}v}(u) - \frac{1}{2} \|v'\|_2^2 \right) = -\frac{2}{\varepsilon^2} \left(\tilde{F}_q(v) + O(\eta^\theta) \right).$$

Proof. Assume for simplicity $\kappa = 1$. We divide the proof into steps.

Step 1: $W_{\mathcal{Q}^{-1/2}v}(u) = \langle v', v' \rangle + O(\eta)$, if $u \in B_C(v, \eta)$ and $v \in H^2$.

Indeed, since $v \in H^2$, we have $W_{\mathcal{Q}^{-1/2}v}(u) = -\langle u, v'' \rangle$ and

$$|-\langle u, v'' \rangle - \langle v', v' \rangle| = |-\langle u, v'' \rangle + \langle v, v'' \rangle| = |\langle v - u, v'' \rangle| \leq \|v - u\|_2 \|v\|_{H^2} \leq \eta \|v\|_{H^2}.$$

Step 2: $I(u) = I(v) + O(\eta^\theta)$, if $u \in B_C(v, \eta)$.

Observe that:

$$|I(u) - I(v)| \leq \frac{\varepsilon_0 \sigma_0}{5} \int_{-1}^1 |(u^5)_+ - (v^5)_+| dx + \int_{-1}^1 Q_0(x) |B(u) - B(v)| dx + q \int_{-1}^1 |u - v| dx$$

By the properties of B and Q_0 , we get that there exists $M, M' > 0$ s.t.

$$\int_{-1}^1 Q_0(x) |B(u) - B(v)| dx + q \int_{-1}^1 |u - v| dx \leq M \int_{-1}^1 |u - v| dx \leq M' \|u - v\|_2 \leq M' \eta.$$

By the mean value theorem, we get that if $u, v \geq 0$ and $p \geq 1$, then

$$|u^p - v^p| \leq p \max\{|u|, |v|\}^{p-1} |u - v| \leq p (|u| + |v|)^{p-1} |u - v|.$$

By this inequality, we get

$$\int_{-1}^1 |(u^p)_+ - (v^p)_+| dx \leq p \int_{-1}^1 (u_+ + v_+)^{p-1} |u_+ - v_+| dx.$$

Let q_1 s.t. $\frac{1}{p} + \frac{1}{q_1} = 1$. By Holder's inequality, we deduce:

$$\begin{aligned} \int_{-1}^1 (u_+ + v_+)^{p-1} |(u)_+ - (v)_+| dx &\leq \left[\int_{-1}^1 (u_+ + v_+)^{q_1(p-1)} dx \right]^{1/q_1} \|u_+ - v_+\|_p \\ &\leq \|u_+ + v_+\|_p^{p/q_1} \|u_+ - v_+\|_p \\ &\leq \left(\|u\|_p + \|v\|_p \right)^{p/q_1} \|u - v\|_p. \end{aligned}$$

Choosing $p = 5$, by interpolation inequality there exists $\theta \in (0, 1)$ s.t.

$$\|u - v\|_p \leq \|u - v\|_2^\theta \|u - v\|_r^{1-\theta} \leq \eta^\theta C^{1-\theta}, \quad u \in B_C(v, \eta).$$

In this way, for $u \in B_C(v, \eta)$, we deduce

$$\int_{-1}^1 |u_+^5 - v_+^5| dx \leq p \left(\|u\|_p + \|v\|_p \right)^{p/q_1} \|u - v\|_p \leq p \left(C + 2\|v\|_p \right)^{p/q_1} \eta^\theta C^{1-\theta} = O(\eta^\theta).$$

■

S3 Variational problem - existence

Given a Banach space X and a sequence $\{u_n\}_n \subseteq X$, we denote by $u_n \rightharpoonup u$ the weak convergence, while we reserve the symbol $u_n \rightarrow u$ for strong convergence. Further, $H^1 = H^{1,2}(-1, 1)$ will denote the Sobolev Space on $[-1, 1]$ with order 1 and exponent 2. The main result of this section is the following.

Proposition S7 *Assume $q > 0$. Then, the variational problem*

$$\inf \left\{ \tilde{F}_q(u) \mid u \in H^{1,2} \right\} \quad (9)$$

admits a minimiser u_0 . Further, $u_0 \in C^\infty$, $u_0'(-1) = u_0'(1) = 0$ and $u_0 \geq 0$.

Proof. Let assume for simplicity $Q_0(x) = 1 \forall x$. This is not restrictive and the proof can be carried on in a similar way since

$$Q_0(x) > \delta > 0.$$

We divide the proof into steps.

Step 1: compactness. We consider the notion of convergence on \mathbb{X} given by:

$$u_n \xrightarrow{\mathbb{X}} u_\infty \quad \text{if and only if } u_n \rightarrow u_\infty \text{ uniformly in } [-1, 1] \text{ and } u_n' \rightharpoonup u_\infty' \text{ in } L^2.$$

We want to verify the compactness of the sublevel sets of \tilde{F}_q . Let $\{u_n\}_n \subset \mathbb{X}$ and $M > 0$ s.t $M \geq \tilde{F}_q(u_n) \forall n$. First, we observe

$$\begin{aligned} M \geq \tilde{F}_q(u_n) &\geq \int_{-1}^1 \varepsilon_0 \sigma_0 \frac{(u_n^5)_+}{5} - B(u_n) - q u_n dx \stackrel{\text{Lagrange Thm 2}}{\geq} 2 \left[\varepsilon_0 \sigma_0 \frac{(u_n^5)_+}{5}(\xi_n) - B(u_n)(\xi_n) - q u_n(\xi_n) \right] \\ &\geq 2 \left[\varepsilon_0 \sigma_0 \frac{(u_n^5)_+}{5}(\xi_n) - B_0 - |u_n(\xi_n)| - q u_n(\xi_n) \right], \end{aligned}$$

where $\xi_n \in [-1, 1]$. Since $v \mapsto \varepsilon_0 \sigma_0 (v^5)_+ / 5 - B(v) - qv$ explodes for $v \rightarrow \pm\infty$, we get the existence of $C_1 > 0$ s.t. $|u_n(\xi_n)| \leq C_1 \forall n$. Second, we get $\forall x \in [-1, 1]$

$$|u_n(x)| \leq |u_n(\xi_n)| + |u_n(x) - u_n(\xi_n)| \leq C_1 + \|u_n'\|_2 |x - \xi_n|^{1/2}, \quad (10)$$

the second inequality follows from the fact that a function in H^1 is Holder-continuous. Third, since (1) holds, we have

$$M \geq \tilde{F}_q(u_n) \geq \int_{-1}^1 -B(u_n) - u_n dx + \frac{1}{2} \|u_n'\|_2 \gtrsim -\|u_n\|_1 + \frac{1}{2} \|u_n'\|_2 \gtrsim -C_1 - \|u_n'\|_2 + \frac{1}{2} \|u_n'\|_2^2,$$

where $a \gtrsim b$ if and only if exists $c > 0$ s.t. $a \geq c \cdot b$ and the last inequality follows from $\|u_n\|_1 \lesssim \|u_n\|_2$. The previous inequality of second order in the unknown $\|u_n'\|_2$ is verified if and only if

$$\|u_n'\|_2 \leq C_2, \quad (11)$$

for some $C_2 > 0$. Up to remaining the subsequence, we have $u'_n \rightharpoonup v$, for a $v \in H^1$. It remains to prove the uniform converge of u_n in $[-1, 1]$. Let's do it using the Ascoli-Arzelà theorem. We get equi-continuity from the properties of the Sobolev space. Indeed

$$|u_n(x) - u_n(y)| \leq \|u'_n\|_{L^2} |x - y|^{1/2} \quad \forall x, y \in [-1, 1]$$

and $\|u'_n\|_{L^2}$ is bounded thanks to weak convergence. Since (10) holds, we get also equi-boundedness. Then (up to remaining) $u_n \rightarrow u_\infty$ uniformly in $[-1, 1]$.

It remains to prove $u'_\infty = v$ in weak sense. Let $\phi \in C_c^\infty([-1, 1])$. Then, by weak derivative definition,

$$\int_{-1}^1 u_n \phi' dx = - \int_{-1}^1 u'_n \phi dx \quad \forall n,$$

and taking the limit on both sides of the equality (we use uniform convergence at LHS, and weak convergence at RHS)

$$\int_{-1}^1 u_\infty \phi' dx = - \int v \phi dx$$

Step 2: lower semi-continuity of \tilde{F}_q . Let $\{u_n\} \subset \mathbb{X}$ be s.t. $u_n \xrightarrow{\mathbb{X}} u$. Let F_1, F_2 be s.t.

$$\tilde{F}_q(u) = F_1(u) + F_2(u), \quad F_1(u) := \int_{-1}^1 \frac{\varepsilon_0 \sigma_0 (u^5)_+ - B(u) - qu}{5} dx, \quad F_2(u) := \frac{\kappa}{2} \int_{-1}^1 (u')^2 dx$$

By uniform convergence, we have $\lim_{n \rightarrow \infty} F_1(u_n) = F_1(u)$; by lower semi-continuity of the L^2 norm w.r.t. weak convergence, we have $\liminf_{n \rightarrow \infty} F_2(u_n) \geq F_2(u)$.

In conclusion, \tilde{F}_q is lower semi-continuous and coercive. Then, $\exists u_0 \in \mathbb{X}$ minimum point for \tilde{F}_q in \mathbb{X} .

Step 3: regularity for u_0 The first variation of \tilde{F}_q in the point u in direction h is given by:

$$\delta \tilde{F}_q(u, h) = \int_{-1}^1 (\psi(u) - \tilde{\beta}(u) - q) h dx + \kappa \int_{-1}^1 u' h' dx.$$

Choosing $h \in C_c^\infty([-1, 1])$ and setting $\phi(t) := F_q(u_0 + th)$, it holds $\phi'(0) = 0$. So:

$$0 = \phi'(0) = \delta \tilde{F}_q(u_0, h),$$

from which it follows

$$\int_{-1}^1 (\psi(u_0) - \tilde{\beta}(u_0) - q) h dx = -\kappa \int_{-1}^1 u'_0 h' dx. \quad (12)$$

Then

$$\kappa u''_0 = \psi(u_0) - \tilde{\beta}(u_0) - q \quad (13)$$

in the weak sense. The RHS is C^0 because $u_0 \in H^1$. Then $u'_0 \in C^1$ and $u_0 \in C^2$. Repeating the bootstrap argument, we get $u_0 \in C^\infty$.

Step 4: Neumann boundary conditions. Let $h \in C^\infty$. Following the same arguments above, we get to (12). Integrating by parts the RHS, we have:

$$\int_{-1}^1 \left(\psi(u_0) - \tilde{\beta}(u_0) - q - u''_0 \right) h \, dx = -\kappa (h(1)u'_0(1) - h(-1)u'_0(-1))$$

But the LHS of the previous equation is null thanks to (13). Choosing h s.t. $h(1) = 0$ and $h(-1) \neq 0$, it follows $u'_0(-1) = 0$. In a similar way, we can get $u'_0(1) = 0$.

Step 5: $u_0 \geq 0$. This can be proved by the following truncation argument. Assume there exists $x_0 \in [-1, 1]$ s.t. $u_0(x_0) < 0$. Consider the following points

$$\tau_1 := \sup \{x < x_0 \mid u_0(x) = 0\}, \quad \tau_2 := \inf \{x > x_0 \mid u_0(x) = 0\}.$$

Let \tilde{u}_0 be the truncation to 0 of u_0 in $[\tau_1, \tau_2]$, i.e.

$$\tilde{u}_0(x) := \begin{cases} u_0(x) & \text{if } x \in [\tau_1, \tau_2]^c, \\ 0 & \text{if } x \in [\tau_1, \tau_2]. \end{cases} \quad (14)$$

Then

$$\begin{aligned} \tilde{F}_q(u_0) - \tilde{F}_q(\tilde{u}_0) &= \int_{\tau_1}^{\tau_2} \varepsilon_0 \sigma_0 \frac{(u_0^5)_+}{5} - Q_0(x)B(u_0) - qu_0(x) \, dx + \frac{\kappa}{2} \int_{\tau_1}^{\tau_2} [u'_0]^2 \, dx + \int_{\tau_1}^{\tau_2} Q_0(x)B(0) \, dx \\ &\geq \int_{\tau_1}^{\tau_2} Q_0(x) (B(0) - B(u_0)) - qu_0(x) \, dx > 0 \end{aligned}$$

where the last inequality follows from the fact that $B(0) \geq -B(u_0(x))$ and $q > 0$. This is a contradiction and thus $u_0(x) \geq 0 \, \forall x \in [-1, 1]$. ■

Repeating again a truncation argument similar to the one used in the last part of the previous proof, we can prove that the minimiser is bounded from above.

Lemma S8 *Assume $q \in (0, b)$. Then, there exists $M > 0$ s.t. if $u_0 = u_0(q)$ is the minimiser for \tilde{F}_q , then $u_0 \leq M$.*

Proof. Indeed, set $\mathcal{R}(x, u) := \varepsilon_0 \sigma_0 \frac{(u^5)_+}{5} - Q_0(x)B(u) - qu$. Note that the following inequalities hold

$$\begin{aligned} \mathcal{R}(x, u) &\geq \varepsilon_0 \sigma_0 \frac{u^5}{5} - \|Q_0\|_\infty B(u) - bu, \quad u \geq 0, \\ \mathcal{R}(x, v) &\leq \varepsilon_0 \sigma_0 \frac{v^5}{5}, \quad v \geq 0. \end{aligned}$$

So, if we set $G(x, u, v) := \mathcal{R}(x, u) - \mathcal{R}(x, v)$, we have, uniformly in x

$$G(x, u, v) \geq \varepsilon_0 \sigma_0 \left(\frac{u^5}{5} - \frac{v^5}{5} \right) - \|Q_0\|_\infty B(u) - bu.$$

Note that for each $v \geq 0$ fixed the term on the right-hand side diverges to $+\infty$ for $u \rightarrow +\infty$. So, given $v \geq 0$, there exists $M = M(v)$ s.t if $u \geq M(v)$, then $G(x, u, v) \geq 1 \forall x \in [-1, 1]$.

Let us pick $v = 1$. We want to prove that $u_0(x) \leq M \forall x \in [-1, 1]$. Assume by contradiction that $u_0(x_0) > M$ for some $x_0 \in [-1, 1]$. Set

$$\tau_1 := \sup \{x < x_0 \mid u_0(x) = M\}, \quad \tau_2 := \inf \{x > x_0 \mid u_0(x) = M\},$$

and consider the truncated minimiser

$$\tilde{u}_0(x) := \begin{cases} u_0(x) & \text{if } x \in [\tau_1, \tau_2]^c, \\ M & \text{if } x \in [\tau_1, \tau_2]. \end{cases} \quad (15)$$

Then

$$\tilde{F}_q(u_0) - \tilde{F}_q(\tilde{u}_0) \geq \int_{\tau_1}^{\tau_2} \mathcal{R}(x, u_*(x)) - \mathcal{R}(x, M) dx = \int_{\tau_1}^{\tau_2} G(x, u_*(x), M) dx.$$

Since by construction $u_0(x) \geq M \forall x \in [\tau_1, \tau_2]$, then we conclude

$$\tilde{F}_q(u_0) - \tilde{F}_q(\tilde{u}_0) > 0.$$

This concludes the proof. ■

S4 Variational problem - uniqueness

In this section we are going to characterize the uniqueness for the solution of the variational problem in terms of the value function, i.e. the minimum value attained by \tilde{F}_q on H^1 .

The *value function* V is defined as follows:

$$V(q) = \inf \left\{ \tilde{F}_q(u) : u \in H^1 \right\}.$$

First of all, from the last result in Section S3 follows the Lipschitz property for V .

Lemma S9 *Assume $q \in (0, b)$. Then, the value function $q \mapsto V(q)$ is Lipschitz continuous.*

Proof. First of all, observe that thanks to the non-negativity of the minimiser and Lemma S8 there exists $M > 0$ s.t.

$$V(q) = \inf \left\{ \tilde{F}_q(u) \mid 0 \leq u \leq M, u \in H^1 \right\}.$$

Second, given a family $\{f_i\}_{i \in I}$ of L_i -Lipschitz function f_i , we know that the infimum $\inf_{i \in I} f_i$ is Lipschitz as long as we can bound uniformly the constants L_i .

In our case this is true. Indeed, given $u \in H^1$, $0 \leq u \leq M$, we have:

$$\left| \tilde{F}_{q_1}(u) - \tilde{F}_{q_2}(u) \right| = |q_2 - q_1| \int_1^1 |u(x)| dx \leq 2M|q_2 - q_1|.$$

This concludes the proof. ■

The main result of this section is the following. We immediately give its proof and postpone to the remaining part of the section the proof of auxiliary results.

Proposition S10 *Assume $q \in (0, b)$. Then, V is differentiable in μ if and only if there exists a unique minimiser for \tilde{F}_μ in H^1 . Further, if V is differentiable, then*

$$V'(\mu) = - \int_{-1}^1 u_* dx,$$

with $u_* \in \operatorname{argmin} \{ \tilde{F}_\mu(u) : u \in H^1 \}$.

Proof. \Rightarrow) Let's consider the auxiliary function

$$W : \mathbb{R} \times (H^1 \cap \{u \geq 0\}) \rightarrow \mathbb{R}$$

given by:

$$W(q, u) := \tilde{F}_q(u) - V(q).$$

As preliminary remarks, note that $W(q, u) \geq 0$ and, if $\tilde{u} \in \operatorname{argmin} \{ \tilde{F}_q(u) : u \in H^1 \}$, then $W(q, \tilde{u}) = 0$. The existence of minimiser has been proved in Proposition S7. To prove uniqueness, assume that u_1, u_2 are two minimisers. Since (i) \tilde{F}_q is differentiable for each q , (ii) V is differentiable in μ by hypothesis and (iii) $W(\mu, u_i) = 0$, $i = 1, 2$, then

$$0 = \partial_q W(\mu, u_i) = - \int_{-1}^1 u_i(x) dx - V'(\mu).$$

Thus

$$- \int_{-1}^1 u_1(x) dx = - \int_{-1}^1 u_2(x) dx = V'(\mu).$$

Using Lemma S10, we know that also $u_1 \wedge u_2$ is a minimiser. With the same reasoning above, it holds

$$- \int_{-1}^1 u_1(x) dx = - \int_{-1}^1 u_2(x) dx = - \int_{-1}^1 u_1 \wedge u_2(x) dx$$

Since $u_i \geq u_1 \wedge u_2 \geq 0$, the previous identities can hold only if $u_1 = u_2$.

\Leftarrow) Assume that, given μ , $\exists!$ u_* minimiser for \tilde{F}_μ . Let $\{h_n\}_n$ be a sequence s.t. $h_n \rightarrow 0$. Let's denote by u_q a minimiser for \tilde{F}_q , i.e.

$$u_q \in \operatorname{argmin} \{ \tilde{F}_q(u) : u \in H^1 \},$$

Then, set $u_n := u_{\mu+h_n}$. We are going to show that $V'(\mu) = - \int_{-1}^1 u_* dx$, i.e.

$$\lim_{n \rightarrow \infty} \frac{V(\mu + h_n) - V(\mu)}{h_n} = - \int_{-1}^1 u_* dx.$$

First, observe that by definition of the value function, we have

$$\frac{V(\mu + h_n) - V(\mu)}{h_n} = \frac{\tilde{F}_{\mu+h_n}(u_n) - \tilde{F}_\mu(u_*)}{h_n} \leq \frac{\tilde{F}_{\mu+h_n}(u_*) - \tilde{F}_\mu(u_*)}{h_n} = - \int_{-1}^1 u_* dx.$$

Hence

$$\lim_{n \rightarrow \infty} \frac{V(\mu + h_n) - V(\mu)}{h_n} \leq - \int_{-1}^1 u_* dx.$$

On the other hand,

$$\begin{aligned} \frac{V(\mu + h_n) - V(\mu)}{h_n} &= \frac{\tilde{F}_{\mu+h_n}(u_n) - \tilde{F}_\mu(u_*)}{h_n} \geq \frac{\tilde{F}_{\mu+h_n}(u_n) - \tilde{F}_\mu(u_n)}{h_n} \\ &= - \int_{-1}^1 u_n dx. \end{aligned}$$

It follows that:

$$\frac{V(\mu + h_n) - V(\mu)}{h_n} \geq - \int_{-1}^1 u_* dx + \int_{-1}^1 (u_* - u_n) dx$$

But the second integral on the right-hand side converges to zero as $n \rightarrow +\infty$ thanks to Lemma S12. This concludes the proof. \blacksquare

In order to complete the proof of the previous result, we need to verify some auxiliaries lemmas. First, let's prove that the infimum of two minimisers for \tilde{F}_q is still a minimiser.

Lemma S11 *If u_1, u_2 are minimisers for \tilde{F}_q in H^1 , then also $u_1 \vee u_2$ and $u_1 \wedge u_2$ are minimisers.*

Proof. For simplicity of notation, set $\mathcal{R}(x, u) := \varepsilon_0 \sigma_0 \frac{u^5}{5} - Q_0(x) \beta(u) - qu$. Further, we divide the proof into steps.

Step 1: $\tilde{F}_q(u_1 \wedge u_2) \geq \tilde{F}_q(u_1 \vee u_2)$.

We start observing that:

$$\begin{aligned} \tilde{F}_q(u_1 \wedge u_2) &= \frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_2)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_2) dx + \frac{\kappa}{2} \int_{u_1 < u_2} (u'_1)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_1) dx \\ &\geq \tilde{F}_q(u_1) = \frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_1)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_1) dx + \frac{\kappa}{2} \int_{u_1 < u_2} (u'_1)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_1) dx \end{aligned}$$

where the inequality holds since u_1 is a minimiser. So, we deduce

$$\frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_2)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_2) dx \geq \frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_1)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_1) dx. \quad (16)$$

In a similar way, we get

$$\frac{\kappa}{2} \int_{u_1 < u_2} (u'_1)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_1) dx \geq \frac{\kappa}{2} \int_{u_1 < u_2} (u'_2)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_2) dx. \quad (17)$$

Indeed, the previous inequality follows bounding from below $\tilde{F}_q(u_1 \wedge u_2)$ with $\tilde{F}_q(u_2)$ and comparing the terms on both sides of the inequality. Now, adding together equation (16) and equation (17), we obtain the claimed relation between $\tilde{F}_q(u_1 \wedge u_2)$ and $\tilde{F}_q(u_1 \vee u_2)$:

$$\begin{aligned}\tilde{F}_q(u_1 \wedge u_2) &= \frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_2)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_2) dx + \frac{\kappa}{2} \int_{u_1 < u_2} (u'_1)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_1) dx \\ &\geq \frac{\kappa}{2} \int_{u_1 \geq u_2} (u'_1)^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, u_1) dx + \frac{\kappa}{2} \int_{u_1 < u_2} (u'_2)^2 dx + \int_{u_1 < u_2} \mathcal{R}(x, u_2) dx \\ &= \tilde{F}_q(u_1 \vee u_2).\end{aligned}$$

Step 2: $\tilde{F}_q(u_1 \vee u_2) \geq \tilde{F}_q(u_1 \wedge u_2)$.

This inequality can be obtained by repeating the step above starting with $\tilde{F}_q(u_1 \vee u_2)$ instead of $\tilde{F}_q(u_1 \wedge u_2)$.

Step 3: $\tilde{F}_q(u_1) = \tilde{F}_q(u_1 \wedge u_2) = \tilde{F}_q(u_1 \vee u_2)$.

The last identity follows from Step 1 and Step 2. To get the first identity, let's observe that in our case

$$\tilde{F}_q(u_1) + \tilde{F}_q(u_2) = \tilde{F}_q(u_1 \wedge u_2) + \tilde{F}_q(u_1 \vee u_2). \quad (18)$$

The previous identity can be verified by writing

$$\tilde{F}_q(v) = \frac{\kappa}{2} \int_{u_1 \geq u_2} (v')^2 dx + \frac{\kappa}{2} \int_{u_1 < u_2} (v')^2 dx + \int_{u_1 \geq u_2} \mathcal{R}(x, v) dx + \int_{u_1 < u_2} \mathcal{R}(x, v) dx,$$

for $v = u_1, u_2, u_1 \wedge u_2, u_1 \vee u_2$. Then, it just consists in checking that equation (18) holds. At this point, since $\tilde{F}_q(u_1) = \tilde{F}_q(u_2)$ (because u_1, u_2 are minimisers) and $\tilde{F}_q(u_1 \wedge u_2) = \tilde{F}_q(u_1 \vee u_2)$ (thanks to Step 1 and Step 2), equation (18) can be rewritten as:

$$2\tilde{F}_q(u_1) = 2\tilde{F}_q(u_1 \wedge u_2).$$

■

Second, we need to verify that the space integral of a sequence of minimisers behaves in a continuous way as the parameter q approaches a point where the uniqueness hold for the variational problem.

Lemma S12 *Assume $q \in (0, b)$ and that there exists an unique minimiser u_* for \tilde{F}_μ in H^1 . Consider a sequence q_n s.t. $q_n \rightarrow \mu$. Then,*

$$\int_{-1}^1 u_n dx \rightarrow \int_{-1}^1 u_* dx,$$

with $u_n \in \operatorname{argmin} \left\{ \tilde{F}_{q_n} : u \in H^1 \right\}$.

Proof. We divide the proof into several steps. Some of them will involve repeating part of the direct method used to solve the variational problem considered in Proposition S7.

Step 1: there exists $u_\infty \in H^1$ and a subsequence $(n_k)_k$ s.t. $u_{n_k} \rightarrow u_\infty$ uniformly in $[-1, 1]$ and $u'_{n_k} \rightharpoonup u'_\infty$ weakly in L_2 .

Indeed, since V is continuous thanks to Lemma S9, we have:

$$\tilde{F}_{q_n}(u_n) = V(q_n) \rightarrow V(\mu) = \tilde{F}_\mu(u_*).$$

In this way, we infer the existence of $M > 0$ s.t. $\tilde{F}_{q_n}(u_n) \leq M \forall n$. At this point, we are in the same hypothesis of the proof for Proposition S7 - Step 1. Following that reasoning, we get the claimed statement.

Step 2: $\tilde{F}_\mu(u_n) \rightarrow \tilde{F}_\mu(u_)$.*

Since u_* is a minimiser for \tilde{F}_μ , we have:

$$\tilde{F}_\mu(u_n) \geq \tilde{F}_\mu(u_*) \quad \forall n.$$

To get the thesis, we fix $\varepsilon > 0$ and we will verify that for n large enough it holds

$$\tilde{F}_\mu(u_n) \leq \tilde{F}_\mu(u_*) + \varepsilon.$$

Indeed

$$\begin{aligned} \left| \tilde{F}_\mu(u_n) - \tilde{F}_\mu(u_*) \right| &\leq \left| \tilde{F}_\mu(u_n) - \tilde{F}_{q_n}(u_n) \right| + \left| \tilde{F}_{q_n}(u_n) - \tilde{F}_\mu(u_*) \right| \\ &\leq |\mu - q_n| \|u_n\|_1 + |V(q_n) - V(\mu)|. \end{aligned}$$

Thanks to Lemma S8, the term $\|u_n\|_1$ is bounded uniformly in n . Further, by the continuity of V we conclude that the right-hand side converges to 0 for $n \rightarrow \infty$.

Step 3: $u_\infty = u_$.*

By proof of Proposition S7 - Step 2, we know that if $u_{n_k} \rightarrow u_\infty$ uniformly in $[-1, 1]$ and $u'_{n_k} \rightharpoonup u'_\infty$ weakly in L^2 , then

$$\liminf_k \tilde{F}_\mu(u_{n_k}) \geq \tilde{F}_\mu(u_\infty).$$

Further, since the sequence $\tilde{F}_\mu(u_n)$ is convergent to $\tilde{F}_\mu(u_*)$, we get

$$\tilde{F}_\mu(u_*) = \lim_n \tilde{F}_\mu(u_n) = \liminf_k \tilde{F}_\mu(u_{n_k}) \geq \tilde{F}_\mu(u_\infty).$$

Thus, by the uniqueness of the minimiser for \tilde{F}_μ , we conclude $u_* = u_\infty$.

Step 4: $u_n \rightarrow u_$ unif. in $[-1, 1]$. In particular, $\lim_{n \rightarrow \infty} \int_{-1}^1 u_n dx = \int_{-1}^1 u_* dx$.*

Take a subsequence u_{n_k} of u_n . We can use the same reasoning of Step 1 and get that there exists a subsubsequence $u_{n_{k_h}}$ s.t. $u_{n_{k_h}} \rightarrow u_\infty$ uniformly in $[-1, 1]$. But with the same reasoning in Step 3, it follows $u_\infty = u_*$. Since the limit does not depend on n_{k_h} , we get the claimed statement. ■

Remark S13 *Note that the previous result is equivalent to say that V' is continuous, where it is defined.*

S5 Value function - semiconcavity, concavity and non-increasing derivative

In the previous section, we introduced the value function V as a tool for characterizing the uniqueness of the variational problem through its differentiability. In this section, our objective is to establish the semiconcavity of V . Subsequently, we will demonstrate that V is also concave. This, in turn, implies that its derivative is non-increasing wherever it is defined. The explicit expression for V' provided in Proposition S10 allows us to say that this last property is equivalent to state that the map $q \mapsto \int_{-1}^1 u_*(x)dx$, where $u_* \in \operatorname{argmin} \{ \tilde{F}_q(u) \mid u \in H^1 \}$, is non-decreasing on the set where V' is well-defined. This set is the complement of a countable set.

First, we start by recalling the definition and the basic properties of semiconcave functions. More details can be found in [4].

Definition S14 *Let $A \subseteq \mathbb{R}$. A function $g: A \rightarrow \mathbb{R}$ is semiconcave if there exists a non-decreasing upper semicontinuous function $\omega: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $\lim_{\rho \rightarrow 0^+} \omega(\rho) = 0$ and*

$$\lambda g(q_1) + (1 - \lambda)g(q_2) - g(\lambda q_1 + (1 - \lambda)q_2) \leq \lambda(1 - \lambda)|q_1 - q_2|\omega(|q_1 - q_2|),$$

for any pair $q_1, q_2 \in A$, such that the segment $[q_1, q_2]$ is contained in A and for any $\lambda \in [0, 1]$. We call ω a modulus of semicontinuity for g in A .

As in the case of Lipschitz functions, taking the infimum of a family of semiconcave functions, the semiconcavity is preserved, provided that the functions have the same modulus.

Lemma S15 [4, Proposition 2.1.5] *Let $\{g_i\}_{i \in I}$ be a family of functions defined on A and semiconcave with the same modulus ω . Then the function $g := \inf_{i \in I} g_i$ is also semiconcave in A with the same modulus of ω , provided $g > -\infty$.*

If a semiconcave function has a modulus satisfying a limiting property, we retrieve the classical notion of concavity.

Lemma S16 [4, Proposition 2.1.9] *Let $g: A \rightarrow \mathbb{R}$ be semiconcave with A open and with a modulus ω such that*

$$\lim_{\rho \rightarrow 0^+} \frac{\omega(\rho)}{\rho} = 0.$$

Then g is concave on all convex subsets of A .

Second, we recall some elementary properties from convex analysis. More details can be found in [9]. Given $g: A \rightarrow \mathbb{R}$, with $A \subset \mathbb{R}$ convex, we denote by:

$$g'_-(q) = \lim_{h \rightarrow 0^-} \frac{g(q+h) - g(q)}{h}, \quad g'_+(q) = \lim_{h \rightarrow 0^+} \frac{g(q+h) - g(q)}{h},$$

respectively the *left* and *right derivative* of g in q . The following lemma summarizes some basic results for a concave function.

Lemma S17 *Let $g: A \rightarrow \mathbb{R}$ be a concave function. Then the following statements hold.*

- (i) *g is differentiable at all but at most countably many points.*
- (ii) *g'_- and g'_+ are well defined and, respectively, left and right continuous.*
- (iii) *For each $q_1 < q_2 \in \text{Int}(A)$, we have:*

$$g'_+(q_2) \leq g'_-(q_2) \leq g'_+(q_1) \leq g'_-(q_1).$$

Lastly, we are able to present the main result of the section.

Proposition S18 *Let $b > 0$ and consider the value function $V: (0, b) \rightarrow \mathbb{R}$. The following statements hold.*

- (i) *Let ω be a non-decreasing upper semicontinuous function $\omega: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $\lim_{\rho \rightarrow 0_+} \omega(\rho) = 0$. Then the value function is semiconcave with modulus ω .*
- (ii) *The value function is concave.*
- (iii) *There exists a countable set $S \subseteq (0, b)$ such that V is differentiable on $S^c \cap (0, b)$ and its derivative V' is non-increasing on $S^c \cap (0, b)$.*

Proof. (a) Since $V(q) = \inf_{u \in H^1} \tilde{F}_q(u)$, thanks to Lemma S15 it is sufficient to prove that, given $u \in H^1$, the map $q \mapsto \tilde{F}_q(u) = \tilde{F}(q)$ is semiconcave. This is true since \tilde{F}_q is an affine function which respect to q and thus its modulus is also independent of u .

(b) It is a direct application of Lemma S16.

(c) Since V is concave, it is sufficient to recall the properties expressed in Lemma S17 and remember that V is differentiable in q if and only if $V'_-(q) = V'_+(q)$, and in that case the derivative is equal to the side derivatives. ■

We conclude the section highlighting how the properties of the value function's derivative reflect on the global mean temperature of the variational problem's minimiser.

Remark S19 *By Froda's theorem, a monotone function can have only jump discontinuities and further they can be at most countable. Since V'_-, V'_+ are monotone thanks to Lemma S17 and*

$$V'(q) = - \int_{-1}^1 u_*(x) dx,$$

where $u_* \in \text{argmin} \{ \tilde{F}_q(u) \mid u \in H^1 \}$, we get that the global mean temperature, depending on q , is monotone increasing and continuous, except for at most a countable number of jumps. Further, the jumps coincide with the points in the set S from Proposition S18 where the value function V is not differentiable.

S6 Mountain Pass Theorem and existence of at least three steady-state solutions

In this section, we are going to use the Mountain pass theorem (MPT) from the calculus of variation to show the existence of at least three solutions to the elliptic problem (2). First, we start by checking that the functional \tilde{F}_q satisfies the compactness condition (Palais-Smale) needed in the hypothesis of the MPT. Second, we are going to show how numerical simulations suggest the existence of two (local) minimum points for \tilde{F}_q corresponding to u_S and u_W ; thus, the MPT gives us the existence of a third critical point, that corresponds to u_M thanks to numerical simulations. Third, we are giving sufficient hypotheses in order to prove the existence of the two local minimum points mentioned before; the existence of these two local minimum points is again obtained using the direct method.

Let $(X, \|\cdot\|)$ be a reflexive Banach space, $\Phi \in C^1(X, \mathbb{R})$ be a functional and Φ' denote the first variation of Φ .

Definition S20 *The functional Φ satisfies the Palais-Smale condition ((PS)-condition) if any sequence $\{u_n\}_n \subset X$ s.t.*

$$\Phi(u_n) \text{ is bounded and } \Phi'(u_n) \rightarrow 0,$$

admits a convergent subsequence.

The previous one is a compactness condition needed in order to use the Mountain pass theorem, which loosely speaking affirms the existence of a mountain pass between two valleys. See [6] for more details.

Theorem S21 (Mountain pass) *If Φ satisfies the (PS)-condition, $\Phi(0) = 0$ and*

$$\begin{aligned} \exists \rho, \alpha > 0 \text{ s.t. } \Phi(x) &\geq \alpha \quad \forall x \text{ with } \|x\| = \rho, \\ \exists x_1 \text{ s.t. } \|x_1\| &> \rho \text{ and } \Phi(x_1) \leq 0 \end{aligned}$$

then, $\exists x_2$ s.t. $\Phi(x_2) = c \geq \alpha$ and x_2 is a stationary point for Φ . Further,

$$c = \inf_{\gamma \in \Gamma} \max_{u \in \gamma([0,1])} \Phi(u)$$

where:

$$\Gamma = \{\gamma \in C([0,1], X) : \gamma(0) = 0, \gamma(1) = x_2\}.$$

At this point, we already know from Proposition S7 the existence of a global minimiser for \tilde{F}_q . On the other hand, numerical simulations suggest the existence of a second local minimiser. Indeed, the second variation $\delta^2 \tilde{F}_q$ of the functional \tilde{F}_q in the point u in direction h is given by:

$$\delta^2 \tilde{F}_q(u, h) = \int_{-1}^1 [4u^3 h - Q_0(x)\beta'(u)h - \kappa h''] h dx.$$

We denote by:

$$\lambda_1(u) \leq \lambda_2(u) \leq \dots,$$

the eigenvalues of the second variation

$$h \mapsto 4u^3h - Q_0(x)\beta'(u)h - \partial_x(\kappa(x)h').$$

We numerically evaluate the eigenvalues of the second variations in the three steady-state points $u_S \leq u_M \leq u_W$. The results, which are shown in Figure S1, tell us that u_S and u_W are strict local minimum points, except at the bifurcations points. This is because the smallest eigenvalues λ_1 for u_S and u_W is positive, hence the second variation in u_S and u_W is positive definite. From this, we get numerical evidence for the existence of a second minimiser. At this point, the Mountain pass theorem guarantees the existence of a third steady-state point, that from our numerical simulations corresponds to u_M , if we are able to prove the (PS)-property for \tilde{F}_q . This is what we are going to check in the following.




Figure S1: Smallest eigenvalues λ_1 for the second variation in u_S, u_M, u_W .

Let X be a reflexive Banach space and X^* its dual space. Given $x_n, x \in X$, denote by $x_n \rightharpoonup x$ the weak convergence in X .

Definition S22 $A: X \rightarrow X^*$ is of type $(S)_+$ if any $\{x_n\}_n \subset X$ s.t. $x_n \rightharpoonup x$ and $\limsup_{n \rightarrow +\infty} \langle A(x_n), x_n - x \rangle \leq 0$ imply $x_n \rightarrow x$ in X .

Let $X = W_n^{1,2} = W_n^{1,2}(-1, 1)$, i.e. the Banach space

$$W_n^{1,2}(-1, 1) = \left\{ u \in W^{1,2}(-1, 1) : u = \lim_{n \rightarrow \infty} u_n \text{ in } W^{1,2}, u_n \in C^\infty([-1, 1]), u'_n(-1) = u'_n(1) = 0 \right\}.$$

Further, let $A: X \rightarrow X^*$ be the operator given by:

$$\langle A(u), v \rangle = \int_{-1}^1 u'v' dx,$$

where $\langle \cdot, \cdot \rangle =_{X^*} \langle \cdot, \cdot \rangle_X$ denotes the duality pairing. We need to recall the following property of the operator A .

Proposition S23 ([1]) *The operator A is of type $(S)_+$.*

At this point, we are able to check the Palais-Smale condition for the functional \tilde{F}_q .

Proposition S24 *The functional $\tilde{F}_q: W_n^{1,2} \rightarrow \mathbb{R}$ satisfies the (PS)-condition.*

Proof. Consider $\{u_n\}_n \subseteq W_n^{1,2}$ and assume there exist $M > 0$ and a sequence $\{\varepsilon_n\}_n$ s.t.

$$\left| \tilde{F}_q(u_n) \right| \leq M, \quad \left\| \tilde{F}'_q(u_n) \right\|_{(W_n^{1,2})^*} \leq \varepsilon_n, \quad (19)$$

where $\varepsilon_n > 0$, $\varepsilon_n \rightarrow 0$ and

$$\langle \tilde{F}'_q(u), v \rangle = \int_{-1}^1 \left(\psi(u) \varepsilon_0 \sigma_0 - q - \tilde{\beta}(u) \right) v \, dx - \langle A(u), v \rangle := \int_{-1}^1 f(x, u) v \, dx - \langle A(u), v \rangle.$$

The proof is divided in two steps.

Step 1: u_n is bounded in $W_n^{1,2}$.

This first step is a corollary of the proof of Proposition S7. Indeed, in that proof, we assume $\tilde{F}_q(u_n) \leq M$ and we prove the boundedness of $\|u_n\|_\infty$ thanks to (10) and the boundedness of $\|u'_n\|_2$ thanks to (11).

Step 2: $\exists n_k$ s.t. $u_{n_k} \rightarrow u$ in $W_n^{1,2}$.

Up to subsequence, by the previous point we get $u_n \rightharpoonup u$ for some $u \in W_n^{1,2}$. Since the embedding $W_n^{1,2} \hookrightarrow L^2$ is compact, we deduce, again up to a subsequence, that $u_n \rightarrow u$ in L^2 . By (19), we have:

$$|\langle A(u_n), v \rangle - \langle f, v \rangle_{L^2}| \leq \varepsilon_n \quad \forall v \in W_n^{1,2}.$$

So, we can choose $v = u_n - u \in W_n^{1,2}$ and get

$$|\langle A(u_n), u - u_n \rangle| \leq \left| \langle \tilde{F}'_q(u_n), u - u_n \rangle \right| + |\langle f, u - u_n \rangle_{L^2}| \leq \varepsilon_n + \|f\|_2 \|u_n - u\|_2.$$

Taking the limits on both sides of the previous inequality we get:

$$\lim_{n \rightarrow \infty} \langle A(u_n), u - u_n \rangle = 0.$$

Since the operator A is of type $(S)_+$, we conclude $u_n \rightarrow u$ in $W_n^{1,2}$.

■

We conclude the section by giving sufficient conditions in order to have at least three solutions for the elliptic PDE (2). We introduce

$$\bar{\mathcal{R}}: \mathbb{R} \rightarrow \mathbb{R}, \quad \bar{\mathcal{R}}(u) = \frac{1}{2} \int_{-1}^1 \mathcal{R}(x, u) dx.$$

where we recall that \mathcal{R} is such that:

$$\tilde{F}_q(u) = \frac{k}{2} \|u'\|_2^2 + \int_{-1}^1 \mathcal{R}(x, u(x)) dx.$$

The assumptions we need in order to get our results are basically three: (1) the space averaged EBM with potential $\bar{\mathcal{R}}$ has (at least) two stable steady-state solutions, (2) the viscosity $\kappa > 0$ of the 1D-EBM is sufficiently large, (3) the two wells in the potential functional $\bar{\mathcal{R}}$ corresponding to the two minimum points are sufficiently deep.

Theorem S25 *Assume $\bar{\mathcal{R}}$ has two minimum points $u_1 \neq u_2$, with $\tilde{F}_q(u_1) \geq \tilde{F}_q(u_2)$. There exist $\omega > 0$ and $f, g \in O(\varepsilon^{-1})$ as $\varepsilon \rightarrow 0^+$ s.t. if $\bar{\varepsilon} > 0$ satisfies:*

$$(i) \quad \bar{\mathcal{R}}''(u_i) > f(\bar{\varepsilon}) \text{ for } i = 1, 2,$$

$$(ii) \quad \kappa > g(\bar{\varepsilon}),$$

$$(iii) \quad \bar{\varepsilon} \leq \omega,$$

then \tilde{F}_q has two local minimum points \tilde{u}_1, \tilde{u}_2 such that:

$$(a) \quad B_{H^1}(u_1, \bar{\varepsilon}) \cap B_{H^1}(u_2, \bar{\varepsilon}) = \emptyset.$$

$$(b) \quad \tilde{u}_i \in B_{H^1}(u_i, \bar{\varepsilon}), \text{ for } i = 1, 2,$$

$$(c) \quad \text{If } \|u - u_1\|_{H^1} = \bar{\varepsilon}, \text{ then } \tilde{F}_q(u) \geq \tilde{F}_q(u_1) + \delta, \text{ with } \delta = \delta(\bar{\varepsilon}) > 0.$$

Proof. The proof consists in repeating the direct method used to prove Proposition S7 and applying it to the set $H^1 \cap B_{H^1}(u_i, \bar{\varepsilon})$. Indeed, thanks to Lemma S26, we can find $\bar{\varepsilon} > 0$ s.t. $B_{H^1}(u_1, \bar{\varepsilon}) \cap B_{H^1}(u_2, \bar{\varepsilon}) = \emptyset$, $\tilde{u}_i \in B_{H^1}(u_i, \bar{\varepsilon})$ and

$$\|u - u_i\|_{H^1} = \bar{\varepsilon} \implies \tilde{F}_q(u) - \tilde{F}_q(u_i) \geq \delta > 0, \quad \delta = \delta(\bar{\varepsilon}). \quad (20)$$

Now, we consider the set

$$\mathbb{X}_i = H^1 \cap \overline{B_{H^1}(u_i, \bar{\varepsilon})},$$

where we stress the fact that $\overline{B_{H^1}(u_i, \bar{\varepsilon})}$ denotes the closed ball in H^1 . Considering a sequence $\{u_{n,i}\}_n$, we want to show that the sublevel sets of \tilde{F}_q are compact in \mathbb{X}_i under the following notion of convergence:

$$u_{n,i} \xrightarrow{\mathbb{X}_i} u_\infty \quad \text{if and only if } u_{n,i} \rightarrow u_\infty \text{ uniformly in } [-1, 1] \text{ and } u'_{n,i} \rightharpoonup u'_\infty \text{ in } L^2.$$

Repeating the argument in the proof of Proposition S7, we get the existence of $u_{\infty,i} \in H^1$ s.t. $u_{n,i} \xrightarrow{\mathbb{X}_i} u_{\infty,i}$. Thanks to the uniform convergence, we have

$$\|u_{\infty,i} - u_i\|_{H^1}^2 = \|u_{\infty,i} - u_i\|_2^2 + \|u'_{\infty,i}\|_2^2 = \lim_n \|u_{n,i} - u_i\|_2^2 + \|u'_{\infty,i}\|_2^2.$$

Second, for each n it holds

$$\bar{\varepsilon}^2 \geq \|u_{n,i} - u_i\|_{H^1}^2 = \|u_{n,i} - u_i\|_2^2 + \|u'_{n,i}\|_2^2.$$

Then, using the inferior lower-semi-continuity of the norm, we get:

$$\begin{aligned}\bar{\varepsilon}^2 &\geq \liminf_n \left(\|u_{n,i} - u_i\|_2^2 + \|u'_{n,i}\|_2^2 \right) = \|u_{\infty,i} - u_i\|_2^2 + \liminf_n \|u'_{n,i}\|_2^2 \\ &\geq \|u_{\infty,i} - u_i\|_2^2 \|u'_{\infty,i}\|_2^2 = \|u_{\infty,i} - u_i\|_{H^1}^2.\end{aligned}$$

Hence $u_{\infty,i} \in \overline{B_{H^1}(u_i, \bar{\varepsilon})}$. Since \tilde{F}_q is lower semi-continuous, we get the existence of \tilde{u}_i minimum point for \tilde{F}_q in \mathbb{X}_i . But thanks to the property (20), we deduce $\|\tilde{u}_i - u_i\|_{H^1} < \bar{\varepsilon}$. Hence \tilde{u}_i are local minimum points for \tilde{F}_q in H^1 .
■

Lemma S26 *Assume $\bar{\mathcal{R}}$ as a minimum point u_0 . Then, there exists $\omega > 0$ and $f, g \in O(\varepsilon^{-1})$ as $\varepsilon \rightarrow 0^+$ s.t. if $\varepsilon > 0$ satisfies:*

- (i) $\bar{\mathcal{R}}''(u_0) > f(\varepsilon)$,
- (ii) $\kappa > g(\varepsilon)$,
- (iii) $\varepsilon = \|u - u_0\|_{H^1} \leq \omega$,

then $\tilde{F}_q(u) \geq \tilde{F}_q(u_0) + \delta$, with $\delta = \delta(\varepsilon) > 0$.

Proof. Let $u \in H^1$ s.t. $\|u - u_0\|_{H^1} = \varepsilon$. Since $\varepsilon^2 = \|u - u_0\|_2^2 + \|u'\|_2^2$, we divide the proof in two cases according to the magnitude of $\|u - u_0\|_2^2$ and $\|u'\|_2^2$.

Case 1: $\|u'\|_2^2 \geq \varepsilon^2/2$ and $\|u - u_0\|_2^2 \leq \varepsilon^2/2$.

Since $\mathcal{R}(x, v)$ is locally Lipschitz in v uniformly in x , there exists $L = L(\mathcal{R}, u_0) > 0$ and $\omega = \omega(\mathcal{R}, u_0) > 0$ s.t.

$$|\mathcal{R}(x, v) - \mathcal{R}(x, u_0)| \leq L|v - u_0|, \quad |u - u_0| \leq \omega, \quad x \in [-1, 1].$$

Thus, if $\|u - u_0\|_\infty \leq \|u - u_0\|_{H^1} = \varepsilon \leq \omega$, it holds

$$\int_{-1}^1 |\mathcal{R}(x, u(x)) - \mathcal{R}(x, u_0)| dx \leq L\|u - u_0\|_1 \leq \sqrt{2}L\|u - u_0\|_2 \leq L\varepsilon$$

Using the previous inequality and the bound on $\|u'\|_2^2$, we have

$$\tilde{F}_q(u) - \tilde{F}_q(u_0) \geq - \left| \int_{-1}^1 \mathcal{R}(x, u(x)) - \mathcal{R}(x, u_0) dx \right| + \frac{k}{2} \|u'\|_2^2 \geq -L\varepsilon + \kappa \frac{\varepsilon^2}{4},$$

and thus $g(\varepsilon) := \frac{4L}{\varepsilon}$.

Case 2: $\|u'\|_2^2 \leq \varepsilon^2/2$ and $\|u - u_0\|_2^2 \geq \varepsilon^2/2$.

Let's consider $\bar{u} = \frac{1}{2} \int_{-1}^1 u(x) dx$. We start by pointing out two useful inequalities. First

$$|\bar{u} - u_0| = \left| \frac{1}{2} \int_{-1}^1 u(x) dx - u_0 \right| = \left| \frac{1}{2} \int_{-1}^1 u(x) - u_0 dx \right| \leq \|u - u_0\|_\infty \quad (21)$$

Second, for each $x \in [-1, 1]$, it holds:

$$|u(x) - \bar{u}| = \left| \frac{1}{2} \int_{-1}^1 (u(x) - u(y)) dx \right| \leq \frac{1}{2} \|u'\|_2 \cdot \int_{-1}^1 |x - y|^{1/2} dy \leq \sqrt{2} \|u'\|_2, \quad (22)$$

where the first inequality follows from the Holder properties of the Sobolev space H^1 . At this point, the estimate on the value of the functional at u can be done considering

$$\tilde{F}_q(u) - \tilde{F}_q(u_0) \geq \int_{-1}^1 \mathcal{R}(x, u(x)) - \mathcal{R}(x, u_0) dx$$

and using a Taylor expansion for \mathcal{R} . Indeed,

$$\mathcal{R}(x, u(x)) = \mathcal{R}(x, u_0) + \mathcal{R}_u(x, u_0)(u(x) - u_0) + \mathcal{R}_{uu}(x, u_0) \frac{(u(x) - u_0)^2}{2} + O(\|u - u_0\|_\infty^3).$$

Performing the decomposition

$$u(x) - u_0 = (u(x) - \bar{u}) + (\bar{u} - u_0),$$

we observe that:

$$\int_{-1}^1 \mathcal{R}_u(x, u_0)(\bar{u} - u_0) dx = (\bar{u} - u_0) \int_{-1}^1 \mathcal{R}_u(x, u_0) dx = (\bar{u} - u_0) 2\bar{\mathcal{R}}'(u_0) = 0$$

and thus

$$\begin{aligned} \int_{-1}^1 \mathcal{R}(x, u(x)) - \mathcal{R}(x, u_0) dx &= \int_{-1}^1 \mathcal{R}_u(x, u_0)(u(x) - \bar{u}) dx \\ &\quad + \int_{-1}^1 \mathcal{R}_{uu}(x, u_0) \frac{(u(x) - u_0)^2}{2} dx + O(\|u - u_0\|_\infty^3) \end{aligned} \quad (23)$$

The absolute value of the first term on the right-hand side can be bounded thanks to (22) and Holder's inequality. Indeed

$$\begin{aligned} \left| \int_{-1}^1 \mathcal{R}_u(x, u_0)(u(x) - \bar{u}) dx \right| &\leq \|\mathcal{R}_u(\cdot, u_0)\|_\infty \|u - \bar{u}\|_1 \leq 2\sqrt{2} \|\mathcal{R}_u(\cdot, u_0)\|_\infty \|u'\|_2 \\ &\leq 2 \|\mathcal{R}_u(\cdot, u_0)\|_\infty \varepsilon. \end{aligned}$$

Now, need to estimate the second term on the RHS of (23). Adding and subtracting $\bar{\mathcal{R}}''(u_0)$, we have:

$$\begin{aligned} \int_{-1}^1 \mathcal{R}_{uu}(x, u_0) \frac{(u(x) - u_0)^2}{2} dx &= \frac{1}{2} \int_{-1}^1 \bar{\mathcal{R}}''(u_0)(u - u_0)^2 dx \\ &\quad + \frac{1}{2} \int_{-1}^1 (\mathcal{R}_{uu}(x, u_0) - \bar{\mathcal{R}}''(u_0))(u - u_0)^2 dx. \end{aligned} \quad (24)$$

The first term on the RHS is large thanks to the central assumption of Case 2. In fact

$$\frac{1}{2} \int_{-1}^1 \bar{\mathcal{R}}''(u_0)(u - u_0)^2 dx = \frac{\bar{\mathcal{R}}''(u_0)}{2} \|u - u_0\|_2^2 \geq \bar{\mathcal{R}}''(u_0) \frac{\varepsilon^2}{4}.$$

The absolute value of the second term on the RHS of (24) can be bounded using Holder's inequality as follows:

$$\begin{aligned} \left| \frac{1}{2} \int_{-1}^1 (\mathcal{R}_{uu}(x, u_0) - \bar{\mathcal{R}}''(u_0)) (u - u_0)^2 dx \right| &\leq \frac{1}{2} \|\bar{\mathcal{R}}''(u_0) - \mathcal{R}_{uu}(\cdot, u_0)\|_\infty \|u - u_0\|_2^2 \\ &\leq \|\bar{\mathcal{R}}''(u_0) - \mathcal{R}_{uu}(\cdot, u_0)\|_\infty \frac{\varepsilon^2}{2}. \end{aligned}$$

Summing up, we have shown

$$\begin{aligned} \tilde{F}_q(u) - \tilde{F}_q(u_0) &\geq -2\varepsilon \|\mathcal{R}_u(\cdot, u_0)\|_\infty + \frac{\varepsilon^2}{4} \bar{\mathcal{R}}''(u_0) - \frac{\varepsilon^2}{2} \left\| \bar{\mathcal{R}}''(u_0) - \frac{\varepsilon^2}{2} \mathcal{R}_{uu}(\cdot, u_0) \right\|_\infty + O(\varepsilon^3), \\ &= \frac{\varepsilon^2}{2} \bar{\mathcal{R}}''(u_0) + f_1(\varepsilon) =: \delta(\varepsilon) \end{aligned}$$

with $f_1 \in O(\varepsilon)$, f_1 negative for small ε and

$$f(\varepsilon) := -\frac{2}{\varepsilon^2} f_1(\varepsilon) \in O(\varepsilon^{-1}).$$

■

In conclusion, applying the MPT we get the following result.

Corollary S27 *If the hypothesis of Theorem S25 are satisfied, then the elliptic problem (2) has at least three solutions.*

References

- [1] Sergiu Aizicovici, Nikolaos S. Papageorgiou, and Vasile Staicu. Existence of multiple solutions with precise sign information for superlinear neumann problems. *Annali di Matematica Pura ed Applicata*, 188(4):679–719, February 2009.
- [2] Robbin Bastiaansen, Henk A Dijkstra, and Anna S von der Heydt. Fragmented tipping in a spatially heterogeneous world. *Environmental Research Letters*, 17(4):045006, mar 2022.
- [3] Haim Brezis. *Functional analysis, Sobolev spaces and partial differential equations*, volume 2. Springer, 2011.
- [4] Piermarco Cannarsa and Carlo Sinestrari. *Semiconcave Functions, Hamilton—Jacobi Equations, and Optimal Control*. Birkhäuser Boston, 2004.

- [5] Giuseppe Da Prato. *Kolmogorov Equations for Stochastic PDEs*. Birkhäuser Basel, 2004.
- [6] Youssef Jabri. *The Mountain Pass Theorem*. Cambridge University Press, September 2003.
- [7] Gerald R. North, Louis Howard, David Pollard, and Bruce Wielicki. Variational formulation of budyko-sellers climate models. *Journal of Atmospheric Sciences*, 36(2):255 – 259, 1979.
- [8] Giuseppe Da Prato. *An Introduction to Infinite-Dimensional Analysis*. Springer Berlin Heidelberg, 2006.
- [9] Ralph Tyrell Rockafellar. *Convex Analysis*, volume 28. Princeton University Press, 1970.